



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2023 – 05:38 AM JST

PDB ID : 6KX9  
Title : Crystal structure of 8-mer peptide from avian influenza H5N1 virus in complex with BF2\*1501  
Authors : Xiao, L.; Zhang, L.  
Deposited on : 2019-09-10  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	 76% 21% .
2	B	100	 73% 23% ..
3	C	8	 50% 38% 12%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3108 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	272	2180	1363	382	428	7	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLU	-	expression tag	UNP Q9GIP6
A	3	PHE	-	expression tag	UNP Q9GIP6

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	785	503	127	150	5	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	GLU	-	expression tag	UNP P21611
B	3	PHE	-	expression tag	UNP P21611

- Molecule 3 is a protein called 8-peptide (ARG-ARG-ALA-LEU-ARG-GLU-GLY-TYR).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	8	72	43	17	12	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	54	Total	O	0	0
			54	54		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	B	17	Total	O	0	0
			17	17		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MHC class I



- Molecule 2: Beta-2-microglobulin



- Molecule 3: 8-peptide (ARG-ARG-ALA-LEU-ARG-GLU-GLY-TYR)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.19Å 124.19Å 80.57Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.73 – 2.90 37.73 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (37.73-2.90) 100.0 (37.73-2.90)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	0.14	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	10.80 (at 2.90Å)	Xtrriage
Refinement program	PHENIX 1.14	Depositor
R, $R_{free}$	0.193 , 0.246 0.194 , 0.244	Depositor DCC
$R_{free}$ test set	809 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.7	Xtrriage
Anisotropy	0.749	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 36.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3108	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/2239	0.61	5/3045 (0.2%)
2	B	0.23	0/810	0.69	4/1100 (0.4%)
3	C	0.23	0/72	0.33	0/92
All	All	0.29	0/3121	0.63	9/4237 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	226	ASP	CB-CA-C	-19.09	72.23	110.40
2	B	4	ASP	N-CA-C	11.94	143.22	111.00
2	B	4	ASP	CB-CA-C	-9.29	91.83	110.40
1	A	198	LEU	N-CA-CB	-8.15	94.09	110.40
2	B	5	LEU	N-CA-C	7.46	131.15	111.00
1	A	227	ALA	N-CA-C	-6.93	92.29	111.00
1	A	198	LEU	N-CA-C	-6.64	93.08	111.00
2	B	5	LEU	N-CA-CB	-6.11	98.19	110.40
1	A	227	ALA	N-CA-CB	-6.02	101.67	110.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2180	0	2021	44	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	785	0	738	20	0
3	C	72	0	75	6	0
4	A	54	0	0	2	0
4	B	17	0	0	0	0
All	All	3108	0	2834	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:ASP:O	1:A:246:ILE:HG13	1.88	0.73
1:A:159:TYR:CZ	3:C:3:ALA:HB2	2.27	0.70
1:A:9:ARG:HD3	1:A:113:TYR:OH	1.92	0.70
2:B:51:ALA:HA	2:B:70:ALA:HB2	1.78	0.66
1:A:25:TYR:HB3	1:A:41:SER:HB3	1.79	0.64
1:A:8:LEU:HB2	1:A:168:LEU:HD13	1.79	0.62
1:A:133:PHE:O	1:A:144:LYS:HE2	2.01	0.61
2:B:56:MET:CE	2:B:56:MET:HA	2.32	0.60
2:B:48:MET:HE1	2:B:72:PHE:HB3	1.84	0.58
1:A:6:HIS:CG	1:A:172:VAL:HG21	2.39	0.57
1:A:197:ILE:HG22	1:A:248:ALA:O	2.04	0.57
1:A:210:ARG:HG3	1:A:240:TYR:CD2	2.42	0.55
1:A:38:HIS:HE1	1:A:40:ASN:HD22	1.52	0.55
2:B:31:GLY:HA2	2:B:63:THR:HB	1.89	0.54
2:B:4:ASP:O	2:B:88:THR:HG21	2.08	0.54
1:A:98:LEU:HD12	1:A:99:MET:H	1.73	0.54
2:B:56:MET:HA	2:B:56:MET:HE3	1.88	0.54
1:A:192:LYS:O	1:A:198:LEU:HA	2.09	0.52
1:A:26:VAL:HG21	2:B:56:MET:HB2	1.91	0.52
1:A:213:ALA:HB3	1:A:261:GLU:HB2	1.91	0.52
1:A:185:PRO:HB3	1:A:207:PHE:HB3	1.93	0.51
1:A:193:GLU:HA	1:A:198:LEU:HA	1.92	0.51
2:B:13:SER:HB2	2:B:23:ASN:ND2	2.26	0.51
2:B:34:PRO:O	2:B:86:HIS:HE1	1.94	0.50
2:B:13:SER:HB2	2:B:23:ASN:HD21	1.77	0.49
1:A:6:HIS:CE1	1:A:172:VAL:HG11	2.47	0.48
1:A:37:THR:OG1	3:C:2:ARG:NH2	2.46	0.48
1:A:36:PHE:C	1:A:49:ARG:HB2	2.34	0.48
1:A:31:VAL:HG23	1:A:36:PHE:CE1	2.49	0.48

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:PRO:HG3	1:A:24:TRP:HA	1.96	0.47
1:A:196:GLY:O	1:A:250:PRO:CD	2.62	0.47
1:A:17:ASP:OD2	2:B:36:LYS:HE3	2.15	0.46
1:A:196:GLY:O	1:A:250:PRO:HD3	2.15	0.45
1:A:210:ARG:HG3	1:A:240:TYR:CG	2.52	0.45
1:A:31:VAL:HG23	1:A:36:PHE:CD1	2.52	0.45
1:A:6:HIS:NE2	1:A:172:VAL:HG11	2.32	0.45
2:B:86:HIS:HD2	2:B:88:THR:OG1	2.00	0.44
1:A:47:VAL:HG12	1:A:48:PRO:O	2.18	0.44
2:B:42:MET:HG2	2:B:47:PRO:HA	1.99	0.44
1:A:4:GLU:H	1:A:4:GLU:HG2	1.56	0.44
1:A:208:TYR:CG	1:A:209:PRO:HA	2.52	0.44
1:A:27:ASP:OD1	3:C:2:ARG:NH1	2.51	0.43
1:A:210:ARG:N	1:A:211:PRO:CD	2.80	0.43
1:A:65:GLU:OE2	3:C:1:ARG:HB2	2.18	0.43
1:A:190:TRP:CD1	2:B:100:GLU:HG2	2.53	0.43
1:A:196:GLY:O	1:A:250:PRO:CG	2.66	0.43
2:B:72:PHE:CZ	2:B:74:PRO:HG3	2.52	0.43
1:A:107:ASP:OD1	1:A:109:THR:HG23	2.19	0.43
1:A:5:LEU:HG	1:A:6:HIS:CD2	2.53	0.43
2:B:11:VAL:HG11	2:B:97:TRP:HB2	2.00	0.42
1:A:159:TYR:CE2	3:C:3:ALA:HB2	2.55	0.42
1:A:38:HIS:CE1	1:A:40:ASN:HD22	2.35	0.42
1:A:130:THR:HG23	1:A:132:THR:OG1	2.20	0.42
1:A:186:GLU:HB2	1:A:205:HIS:HB2	2.01	0.41
2:B:21:THR:O	2:B:74:PRO:HD2	2.20	0.41
1:A:151:ASP:HA	4:A:344:HOH:O	2.21	0.41
2:B:101:PHE:CD1	2:B:101:PHE:C	2.93	0.41
2:B:5:LEU:H	2:B:5:LEU:HG	1.45	0.41
1:A:49:ARG:HD3	1:A:49:ARG:HA	1.85	0.40
3:C:4:LEU:HD12	3:C:4:LEU:HA	1.91	0.40
1:A:145:ARG:HD2	4:A:337:HOH:O	2.21	0.40
2:B:7:PRO:HA	2:B:32:PHE:HB3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/272 (99%)	262 (97%)	8 (3%)	0	100	100
2	B	97/100 (97%)	94 (97%)	3 (3%)	0	100	100
3	C	6/8 (75%)	6 (100%)	0	0	100	100
All	All	373/380 (98%)	362 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/225 (100%)	211 (94%)	14 (6%)	18	47
2	B	86/88 (98%)	84 (98%)	2 (2%)	50	80
3	C	6/6 (100%)	5 (83%)	1 (17%)	2	6
All	All	317/319 (99%)	300 (95%)	17 (5%)	22	54

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
1	A	9	ARG
1	A	27	ASP
1	A	45	ARG
1	A	116	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	151	ASP
1	A	182	ARG
1	A	193	GLU
1	A	197	ILE
1	A	198	LEU
1	A	201	SER
1	A	215	SER
1	A	243	TRP
1	A	259	ARG
2	B	6	THR
2	B	72	PHE
3	C	4	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	HIS
1	A	84	GLN
1	A	115	GLN
1	A	225	GLN
2	B	33	HIS
2	B	86	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	272/272 (100%)	-0.29	3 (1%) 80 80	18, 31, 67, 89	0
2	B	99/100 (99%)	-0.03	4 (4%) 38 33	26, 38, 63, 96	0
3	C	8/8 (100%)	-0.51	0 100 100	22, 27, 40, 42	0
All	All	379/380 (99%)	-0.23	7 (1%) 68 67	18, 33, 66, 96	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	3	PHE	6.1
2	B	4	ASP	4.2
1	A	273	TRP	2.8
1	A	197	ILE	2.8
2	B	90	LYS	2.8
2	B	101	PHE	2.5
1	A	226	ASP	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.