



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 16, 2023 – 02:07 AM JST

PDB ID : 6KXW
Title : Crystal structure of human aquaporin AQP7 in bound to glycerol
Authors : Zhang, L.; Yao, D.; Zhou, F.; Zhang, Q.; Zhou, L.; Cao, Y.
Deposited on : 2019-09-13
Resolution : 3.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1049 (3.88-3.52)
Clashscore	141614	1027 (3.86-3.54)
Ramachandran outliers	138981	1069 (3.88-3.52)
Sidechain outliers	138945	1065 (3.88-3.52)
RSRZ outliers	127900	1578 (3.90-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	
1	B	342	
1	C	342	
1	D	342	

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 7586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aquaporin-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	252	Total 1925	C 1278	N 312	O 322	S 13	0	0	0
1	B	249	Total 1890	C 1253	N 308	O 316	S 13	0	0	0
1	C	249	Total 1894	C 1258	N 306	O 317	S 13	0	0	0
1	D	249	Total 1865	C 1234	N 302	O 316	S 13	0	0	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C O		
2	A	1	Total 6	C O 3 3	0	0
2	C	1	Total 6	C O 3 3	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aquaporin-7





- Molecule 1: Aquaporin-7






• Molecule 1: Aquaporin-7



• Molecule 1: Aquaporin-7



L267
 G268
 G269
 I270
 I271
 Y272
 L273
 Y274
 F275
 I276
 G277
 S278
 THR
 ILE
 PRO
 ARG
 ARG
 GLU
 PRO
 PRO
 LEU
 LEU
 LYS
 PRO
 LEU
 LEU
 ASP
 ASP
 SER
 VAL
 VAL
 ALA
 ALA
 TYR
 GLU
 ASP
 HIS
 HIS
 GLY
 ILE
 ILE
 THR
 THR
 VAL
 LEU
 LEU
 PRO
 PRO
 LYS
 LYS
 MET
 MET
 GLY
 SER
 SER
 HIS
 HIS
 GLU
 GLU
 PRO
 THR
 THR
 ILE
 ILE
 SER
 SER
 PRO
 PRO
 LEU
 LEU
 THR
 THR
 PRO
 PRO
 VAL
 VAL
 SER
 SER
 VAL
 VAL
 SER
 SER
 PRO
 PRO
 ALA
 ALA
 ASN
 ASN
 ARG
 ARG
 SER
 SER

VAL
 HIS
 PRO
 ALA
 PRO
 PRO
 LEU
 LEU
 HIS
 HIS
 GLU
 SER
 MET
 MET
 ALA
 ALA
 LEU
 LEU
 GLU
 HIS
 HIS
 PHE

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	93.51Å 106.29Å 214.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.61 – 3.70 50.08 – 3.70	Depositor EDS
% Data completeness (in resolution range)	71.9 (47.61-3.70) 72.1 (50.08-3.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 3.67Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.268 , 0.322 0.268 , 0.322	Depositor DCC
R_{free} test set	883 reflections (5.21%)	wwPDB-VP
Wilson B-factor (Å ²)	55.0	Xtrriage
Anisotropy	0.443	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 132.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.78	EDS
Total number of atoms	7586	wwPDB-VP
Average B, all atoms (Å ²)	109.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.82	5/1982 (0.3%)	0.99	6/2702 (0.2%)
1	B	0.56	1/1946 (0.1%)	0.78	1/2652 (0.0%)
1	C	0.81	3/1951 (0.2%)	1.07	5/2661 (0.2%)
1	D	0.52	0/1918	0.81	1/2617 (0.0%)
All	All	0.69	9/7797 (0.1%)	0.92	13/10632 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
1	C	0	1
1	D	0	2
All	All	0	6

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	89	SER	CA-CB	8.38	1.65	1.52
1	C	256	VAL	C-N	6.59	1.46	1.34
1	A	40	GLU	CG-CD	6.09	1.61	1.51
1	C	164	ASP	CB-CG	5.97	1.64	1.51
1	B	185	CYS	CB-SG	-5.35	1.73	1.81

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	231	LEU	CB-CG-CD2	7.76	124.20	111.00
1	C	267	LEU	CA-CB-CG	-6.08	101.31	115.30
1	A	60	LEU	CA-CB-CG	5.87	128.79	115.30
1	C	231	LEU	CA-CB-CG	5.81	128.67	115.30
1	A	121	LEU	CB-CG-CD1	5.79	120.84	111.00

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	120	PHE	Peptide
1	A	251	GLU	Peptide
1	B	119	GLN	Peptide
1	C	224	ALA	Peptide
1	D	86	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1925	0	1930	184	0
1	B	1890	0	1888	173	0
1	C	1894	0	1891	230	0
1	D	1865	0	1856	199	0
2	A	6	0	8	1	0
2	C	6	0	8	1	0
All	All	7586	0	7581	694	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 46.

The worst 5 of 694 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:TRP:CZ3	1:D:129:THR:HB	1.89	1.07
1:B:38:LEU:HD23	1:D:275:PHE:HA	1.34	1.07

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:ARG:O	1:A:32:LYS:HG2	1.62	1.00
1:D:41:PHE:HZ	1:D:125:LEU:HD12	1.27	0.99
1:D:33:MET:HA	1:D:36:GLU:HG3	1.42	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/342 (73%)	185 (74%)	60 (24%)	5 (2%)	7	39
1	B	247/342 (72%)	176 (71%)	66 (27%)	5 (2%)	7	39
1	C	247/342 (72%)	174 (70%)	67 (27%)	6 (2%)	6	35
1	D	247/342 (72%)	189 (76%)	55 (22%)	3 (1%)	13	48
All	All	991/1368 (72%)	724 (73%)	248 (25%)	19 (2%)	8	40

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	95	ALA
1	A	86	GLY
1	C	86	GLY
1	A	95	ALA
1	B	254	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/278 (70%)	182 (93%)	13 (7%)	16	47
1	B	189/278 (68%)	178 (94%)	11 (6%)	20	52
1	C	191/278 (69%)	178 (93%)	13 (7%)	16	47
1	D	187/278 (67%)	173 (92%)	14 (8%)	13	43
All	All	762/1112 (68%)	711 (93%)	51 (7%)	16	47

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	124	PHE
1	C	174	ASN
1	D	266	TYR
1	C	131	TYR
1	C	147	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	71	ASN
1	D	71	ASN
1	D	195	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	C	400	-	5,5,5	1.68	2 (40%)	5,5,5	0.67	0
2	GOL	A	401	-	5,5,5	1.50	1 (20%)	5,5,5	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	400	-	-	2/4/4/4	-
2	GOL	A	401	-	-	1/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	GOL	C3-C2	2.67	1.62	1.51
2	C	400	GOL	C3-C2	2.41	1.61	1.51
2	C	400	GOL	C1-C2	2.39	1.61	1.51

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	400	GOL	C1-C2-C3-O3
2	C	400	GOL	O2-C2-C3-O3
2	A	401	GOL	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	400	GOL	1	0
2	A	401	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	252/342 (73%)	-0.09	8 (3%) 47 35	17, 71, 132, 174	0
1	B	249/342 (72%)	0.38	27 (10%) 5 4	60, 147, 218, 244	0
1	C	249/342 (72%)	-0.41	2 (0%) 86 78	8, 55, 121, 166	0
1	D	249/342 (72%)	0.41	22 (8%) 10 7	74, 153, 211, 252	0
All	All	999/1368 (73%)	0.07	59 (5%) 22 15	8, 102, 201, 252	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	278	SER	8.4
1	A	89	SER	6.5
1	D	277	GLY	6.4
1	B	238	PHE	4.9
1	B	246	VAL	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	401	6/6	0.91	0.35	69,73,76,82	0
2	GOL	C	400	6/6	0.95	0.22	23,29,48,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.