



# wwPDB X-ray Structure Validation Summary Report

Jan 3, 2024 – 06:02 pm GMT

PDB ID : 5L6V  
Title : Crystal structure of E. coli ADP-glucose pyrophosphorylase (AGPase) in complex with a negative allosteric regulator adenosine monophosphate (AMP) - AGPase\*AMP  
Authors : Cifuentes, J.O.; Albesa-Jove, D.; Comino, N.; Madariaga-Marcos, J.; Agirre, J.; Lopez-Fernandez, S.; Garcia-Alija, M.; Guerin, M.E.  
Deposited on : 2016-05-31  
Resolution : 2.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






















Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	431	
1	B	431	
1	C	431	
1	D	431	
1	E	431	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	431	 % 88% 7% 5%
1	G	431	 7% 87% 8% 5%
1	H	431	 3% 89% 8%
1	I	431	 7% 85% 7% 7%
1	J	431	 3% 90% 6%
1	K	431	 % 89% 8%
1	L	431	 2% 87% 8%
1	M	431	 % 88% 7%
1	N	431	 % 90% 7%
1	O	431	 2% 91% 5%
1	P	431	 % 92% 6%
2	Q	2	 100%
2	R	2	 50% 50%
2	S	2	 100%
2	T	2	 100%
2	U	2	 100%
2	V	2	 100%
2	W	2	 100%
2	X	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FRU	U	2	-	-	-	X



## 2 Entry composition


There are 5 unique types of molecules in this entry. The entry contains 53530 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucose-1-phosphate adenylyltransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	422	Total 3325	C 2100	N 583	O 619	S 23	0	3	0
1	B	421	Total 3293	C 2081	N 579	O 611	S 22	0	0	0
1	C	421	Total 3325	C 2105	N 585	O 612	S 23	0	2	0
1	D	413	Total 3210	C 2028	N 563	O 599	S 20	0	0	0
1	E	411	Total 3202	C 2024	N 561	O 596	S 21	0	1	0
1	F	411	Total 3225	C 2035	N 567	O 602	S 21	0	0	0
1	G	411	Total 3127	C 1985	N 543	O 579	S 20	0	0	0
1	H	416	Total 3162	C 2008	N 545	O 589	S 20	0	0	0
1	I	400	Total 2984	C 1893	N 522	O 549	S 20	0	0	0
1	J	412	Total 3158	C 2002	N 555	O 581	S 20	0	0	0
1	K	421	Total 3292	C 2080	N 578	O 612	S 22	0	0	0
1	L	412	Total 3195	C 2022	N 561	O 591	S 21	0	0	0
1	M	415	Total 3250	C 2056	N 572	O 601	S 21	0	0	0
1	N	422	Total 3285	C 2079	N 575	O 609	S 22	0	0	0
1	O	415	Total 3237	C 2044	N 568	O 604	S 21	0	0	0
1	P	423	Total 3305	C 2090	N 578	O 615	S 22	0	0	0

- Molecule 2 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	Q	2	Total	C	O	0	0	0
			23	12	11			
2	R	2	Total	C	O	0	0	0
			23	12	11			
2	S	2	Total	C	O	0	0	0
			23	12	11			
2	T	2	Total	C	O	0	0	0
			23	12	11			
2	U	2	Total	C	O	0	0	0
			23	12	11			
2	V	2	Total	C	O	0	0	0
			23	12	11			
2	W	2	Total	C	O	0	0	0
			23	12	11			
2	X	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C<sub>10</sub>H<sub>14</sub>N<sub>5</sub>O<sub>7</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	D	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	E	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	F	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	G	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	H	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	I	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	J	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	K	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	L	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	M	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	N	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	O	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	P	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	O P	0	0
			5	4 1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	104	Total	O	0	0
			104	104		
5	B	85	Total	O	0	0
			85	85		
5	C	148	Total	O	0	0
			148	148		
5	D	97	Total	O	0	0
			97	97		
5	E	92	Total	O	0	0
			92	92		
5	F	101	Total	O	0	0
			101	101		
5	G	62	Total	O	0	0
			62	62		
5	H	42	Total	O	0	0
			42	42		
5	I	57	Total	O	0	0
			57	57		
5	J	74	Total	O	0	0
			74	74		
5	K	75	Total	O	0	0
			75	75		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	L	81	Total O 81 81	0	0
5	M	111	Total O 111 111	0	0
5	N	142	Total O 142 142	0	0
5	O	67	Total O 67 67	0	0
5	P	60	Total O 60 60	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucose-1-phosphate adenylyltransferase




- Molecule 1: Glucose-1-phosphate adenylyltransferase



- Molecule 1: Glucose-1-phosphate adenylyltransferase





- Molecule 1: Glucose-1-phosphate adenylyltransferase










- Molecule 1: Glucose-1-phosphate adenylyltransferase






- Molecule 1: Glucose-1-phosphate adenylyltransferase




- Molecule 1: Glucose-1-phosphate adenylyltransferase




- Molecule 1: Glucose-1-phosphate adenylyltransferase




- Molecule 1: Glucose-1-phosphate adenylyltransferase




- Molecule 1: Glucose-1-phosphate adenylyltransferase




- Molecule 1: Glucose-1-phosphate adenylyltransferase




- Molecule 1: Glucose-1-phosphate adenylyltransferase




- Molecule 1: Glucose-1-phosphate adenylyltransferase









• Molecule 1: Glucose-1-phosphate adenylyltransferase



• Molecule 1: Glucose-1-phosphate adenylyltransferase



• Molecule 1: Glucose-1-phosphate adenylyltransferase



• Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



• Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



GLC1  
FRU2

- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain S:  100%

GLC1  
FRU2

- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain T:  100%

GLC1  
FRU2

- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain U:  100%

GLC1  
FRU2

- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain V:  100%

GLC1  
FRU2

- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain W:  100%

GLC1  
FRU2

- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain X:  100%

GLC1  
FRU2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	166.88Å 148.19Å 180.16Å 90.00° 115.48° 90.00°	Depositor
Resolution (Å)	45.27 – 2.67 45.27 – 2.67	Depositor EDS
% Data completeness (in resolution range)	95.1 (45.27-2.67) 99.8 (45.27-2.67)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 2.65Å)	Xtrriage
Refinement program	PHENIX (dev_2219: ???)	Depositor
R, $R_{free}$	0.239 , 0.263 0.223 , 0.251	Depositor DCC
$R_{free}$ test set	11273 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.6	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 30.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.006 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	53530	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, GLC, FRU, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/3396	0.45	0/4605
1	B	0.26	0/3363	0.45	0/4560
1	C	0.25	0/3404	0.44	0/4613
1	D	0.25	0/3277	0.45	0/4445
1	E	0.25	0/3269	0.44	0/4432
1	F	0.25	0/3292	0.46	0/4459
1	G	0.25	0/3194	0.44	0/4341
1	H	0.25	0/3232	0.44	0/4398
1	I	0.25	0/3047	0.43	0/4148
1	J	0.26	0/3225	0.46	0/4379
1	K	0.25	0/3362	0.44	0/4556
1	L	0.25	0/3262	0.45	0/4422
1	M	0.25	0/3319	0.45	0/4497
1	N	0.25	0/3356	0.45	0/4552
1	O	0.25	0/3304	0.45	0/4480
1	P	0.25	0/3376	0.44	0/4577
All	All	0.25	0/52678	0.45	0/71464

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3325	0	3262	15	0
1	B	3293	0	3234	21	0
1	C	3325	0	3282	32	0
1	D	3210	0	3141	29	0
1	E	3202	0	3141	17	0
1	F	3225	0	3180	19	0
1	G	3127	0	3027	19	0
1	H	3162	0	3030	19	0
1	I	2984	0	2800	19	0
1	J	3158	0	3078	12	0
1	K	3292	0	3233	22	0
1	L	3195	0	3141	19	0
1	M	3250	0	3187	20	0
1	N	3285	0	3215	19	0
1	O	3237	0	3177	14	0
1	P	3305	0	3240	15	0
2	Q	23	0	21	0	0
2	R	23	0	21	0	0
2	S	23	0	21	0	0
2	T	23	0	21	0	0
2	U	23	0	21	0	0
2	V	23	0	21	0	0
2	W	23	0	21	0	0
2	X	23	0	21	0	0
3	A	23	0	12	0	0
3	B	23	0	12	0	0
3	C	23	0	12	0	0
3	D	23	0	12	0	0
3	E	23	0	12	0	0
3	F	23	0	12	0	0
3	G	23	0	12	0	0
3	H	23	0	12	0	0
3	I	23	0	12	1	0
3	J	23	0	12	0	0
3	K	23	0	12	0	0
3	L	23	0	12	0	0
3	M	23	0	12	0	0
3	N	23	0	12	0	0
3	O	23	0	12	0	0
3	P	23	0	12	0	0
4	D	5	0	0	0	0
5	A	104	0	0	1	0
5	B	85	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	148	0	0	0	0
5	D	97	0	0	0	0
5	E	92	0	0	0	0
5	F	101	0	0	0	0
5	G	62	0	0	3	0
5	H	42	0	0	1	0
5	I	57	0	0	2	0
5	J	74	0	0	0	0
5	K	75	0	0	0	0
5	L	81	0	0	1	0
5	M	111	0	0	3	0
5	N	142	0	0	0	0
5	O	67	0	0	0	0
5	P	60	0	0	1	0
All	All	53530	0	50728	284	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 284 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:37:THR:O	1:I:75:TYR:OH	2.00	0.79
1:M:13:ALA:HA	1:M:154:ILE:HD11	1.63	0.78
1:B:235:ASN:OD1	1:K:423:ARG:NH1	2.20	0.73
1:B:274:TRP:CZ2	1:B:274:TRP:CZ3	2.76	0.70
1:J:63:ASN:O	1:K:14:ARG:NH2	2.29	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/431 (98%)	414 (98%)	9 (2%)	0	100	100
1	B	419/431 (97%)	409 (98%)	9 (2%)	1 (0%)	47	64
1	C	421/431 (98%)	409 (97%)	12 (3%)	0	100	100
1	D	409/431 (95%)	401 (98%)	8 (2%)	0	100	100
1	E	408/431 (95%)	398 (98%)	9 (2%)	1 (0%)	47	64
1	F	407/431 (94%)	397 (98%)	10 (2%)	0	100	100
1	G	407/431 (94%)	397 (98%)	10 (2%)	0	100	100
1	H	412/431 (96%)	400 (97%)	11 (3%)	1 (0%)	47	64
1	I	390/431 (90%)	377 (97%)	12 (3%)	1 (0%)	41	56
1	J	408/431 (95%)	397 (97%)	10 (2%)	1 (0%)	47	64
1	K	419/431 (97%)	410 (98%)	9 (2%)	0	100	100
1	L	408/431 (95%)	396 (97%)	11 (3%)	1 (0%)	47	64
1	M	409/431 (95%)	396 (97%)	12 (3%)	1 (0%)	47	64
1	N	420/431 (97%)	410 (98%)	10 (2%)	0	100	100
1	O	411/431 (95%)	398 (97%)	13 (3%)	0	100	100
1	P	421/431 (98%)	411 (98%)	10 (2%)	0	100	100
All	All	6592/6896 (96%)	6420 (97%)	165 (2%)	7 (0%)	51	69

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	74	GLN
1	J	74	GLN
1	B	75	TYR
1	H	74	GLN
1	M	74	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/373 (96%)	355 (99%)	2 (1%)	86	92
1	B	354/373 (95%)	350 (99%)	4 (1%)	73	85
1	C	358/373 (96%)	356 (99%)	2 (1%)	86	92
1	D	343/373 (92%)	342 (100%)	1 (0%)	92	96
1	E	343/373 (92%)	339 (99%)	4 (1%)	71	84
1	F	349/373 (94%)	348 (100%)	1 (0%)	92	96
1	G	326/373 (87%)	323 (99%)	3 (1%)	78	87
1	H	328/373 (88%)	323 (98%)	5 (2%)	65	80
1	I	297/373 (80%)	293 (99%)	4 (1%)	69	82
1	J	331/373 (89%)	329 (99%)	2 (1%)	86	92
1	K	353/373 (95%)	350 (99%)	3 (1%)	81	89
1	L	341/373 (91%)	334 (98%)	7 (2%)	53	72
1	M	347/373 (93%)	343 (99%)	4 (1%)	71	84
1	N	350/373 (94%)	348 (99%)	2 (1%)	86	92
1	O	348/373 (93%)	347 (100%)	1 (0%)	92	96
1	P	354/373 (95%)	353 (100%)	1 (0%)	92	96
All	All	5479/5968 (92%)	5433 (99%)	46 (1%)	81	89

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	125	LEU
1	L	201	SER
1	K	154	ILE
1	L	29	ARG
1	L	430	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	106	GLN
1	H	123	GLN
1	K	112	ASN
1	P	106	GLN



### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	Q	1	2	11,11,12	0.60	0	15,15,17	0.81	0
2	FRU	Q	2	2	11,12,12	0.62	0	10,18,18	0.77	0
2	GLC	R	1	2	11,11,12	0.52	0	15,15,17	0.92	1 (6%)
2	FRU	R	2	2	11,12,12	0.71	0	10,18,18	0.80	0
2	GLC	S	1	2	11,11,12	0.57	0	15,15,17	0.82	0
2	FRU	S	2	2	11,12,12	0.65	0	10,18,18	0.99	0
2	GLC	T	1	2	11,11,12	0.57	0	15,15,17	0.75	0
2	FRU	T	2	2	11,12,12	0.63	0	10,18,18	0.73	0
2	GLC	U	1	2	11,11,12	0.53	0	15,15,17	0.96	1 (6%)
2	FRU	U	2	2	11,12,12	0.71	0	10,18,18	1.15	1 (10%)
2	GLC	V	1	2	11,11,12	0.56	0	15,15,17	0.91	0
2	FRU	V	2	2	11,12,12	0.68	0	10,18,18	0.61	0
2	GLC	W	1	2	11,11,12	0.56	0	15,15,17	0.72	0
2	FRU	W	2	2	11,12,12	0.69	0	10,18,18	0.63	0
2	GLC	X	1	2	11,11,12	0.57	0	15,15,17	0.88	0
2	FRU	X	2	2	11,12,12	0.70	0	10,18,18	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	Q	1	2	-	0/2/19/22	0/1/1/1
2	FRU	Q	2	2	-	0/5/24/24	0/1/1/1
2	GLC	R	1	2	-	0/2/19/22	0/1/1/1
2	FRU	R	2	2	-	5/5/24/24	0/1/1/1
2	GLC	S	1	2	-	0/2/19/22	0/1/1/1
2	FRU	S	2	2	-	0/5/24/24	0/1/1/1
2	GLC	T	1	2	-	0/2/19/22	0/1/1/1
2	FRU	T	2	2	-	5/5/24/24	0/1/1/1
2	GLC	U	1	2	-	2/2/19/22	0/1/1/1
2	FRU	U	2	2	-	3/5/24/24	0/1/1/1
2	GLC	V	1	2	-	2/2/19/22	0/1/1/1
2	FRU	V	2	2	-	2/5/24/24	0/1/1/1
2	GLC	W	1	2	-	0/2/19/22	0/1/1/1
2	FRU	W	2	2	-	2/5/24/24	0/1/1/1
2	GLC	X	1	2	-	0/2/19/22	0/1/1/1
2	FRU	X	2	2	-	5/5/24/24	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	U	1	GLC	C1-O5-C5	2.30	115.31	112.19
2	U	2	FRU	O2-C2-O5	2.29	113.93	109.50
2	R	1	GLC	O5-C5-C6	2.23	110.70	107.20

There are no chirality outliers.


5 of 26 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	T	2	FRU	O1-C1-C2-O2
2	T	2	FRU	O1-C1-C2-O5
2	X	2	FRU	O1-C1-C2-C3
2	X	2	FRU	O1-C1-C2-O2
2	X	2	FRU	O1-C1-C2-O5


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









## 5.6 Ligand geometry

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	AMP	G	501	-	22,25,25	0.66	0	25,38,38	0.87	2 (8%)
3	AMP	I	501	-	22,25,25	0.72	0	25,38,38	0.91	2 (8%)
3	AMP	P	501	-	22,25,25	0.79	1 (4%)	25,38,38	0.94	2 (8%)
3	AMP	N	501	-	22,25,25	0.79	1 (4%)	25,38,38	0.96	2 (8%)
3	AMP	D	501	-	22,25,25	0.70	0	25,38,38	0.90	2 (8%)
3	AMP	F	501	-	22,25,25	0.68	0	25,38,38	0.89	2 (8%)
3	AMP	M	501	-	22,25,25	0.71	0	25,38,38	0.88	2 (8%)
3	AMP	L	501	-	22,25,25	0.76	1 (4%)	25,38,38	0.91	2 (8%)
3	AMP	A	501	-	22,25,25	0.58	0	25,38,38	0.87	2 (8%)
3	AMP	E	501	-	22,25,25	0.75	0	25,38,38	0.92	2 (8%)
3	AMP	B	501	-	22,25,25	0.75	0	25,38,38	0.95	2 (8%)
4	PO4	D	502	-	4,4,4	0.88	0	6,6,6	0.42	0
3	AMP	J	501	-	22,25,25	0.74	1 (4%)	25,38,38	0.90	2 (8%)
3	AMP	K	501	-	22,25,25	0.68	0	25,38,38	0.89	2 (8%)
3	AMP	H	501	-	22,25,25	0.66	0	25,38,38	0.89	2 (8%)
3	AMP	C	501	-	22,25,25	0.77	1 (4%)	25,38,38	0.93	2 (8%)
3	AMP	O	501	-	22,25,25	0.70	0	25,38,38	0.95	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMP	G	501	-	-	5/6/26/26	0/3/3/3
3	AMP	I	501	-	-	5/6/26/26	0/3/3/3
3	AMP	P	501	-	-	5/6/26/26	0/3/3/3

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMP	N	501	-	-	0/6/26/26	0/3/3/3
3	AMP	D	501	-	-	3/6/26/26	0/3/3/3
3	AMP	F	501	-	-	0/6/26/26	0/3/3/3
3	AMP	M	501	-	-	3/6/26/26	0/3/3/3
3	AMP	L	501	-	-	0/6/26/26	0/3/3/3
3	AMP	A	501	-	-	2/6/26/26	0/3/3/3
3	AMP	E	501	-	-	5/6/26/26	0/3/3/3
3	AMP	B	501	-	-	5/6/26/26	0/3/3/3
3	AMP	J	501	-	-	3/6/26/26	0/3/3/3
3	AMP	K	501	-	-	3/6/26/26	0/3/3/3
3	AMP	H	501	-	-	5/6/26/26	0/3/3/3
3	AMP	C	501	-	-	5/6/26/26	0/3/3/3
3	AMP	O	501	-	-	5/6/26/26	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	N	501	AMP	P-O2P	-2.28	1.46	1.54
3	P	501	AMP	P-O2P	-2.27	1.46	1.54
3	L	501	AMP	P-O2P	-2.14	1.46	1.54
3	C	501	AMP	P-O2P	-2.12	1.46	1.54
3	J	501	AMP	P-O2P	-2.02	1.47	1.54

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	O	501	AMP	P-O5'-C5'	3.00	126.56	118.30
3	B	501	AMP	P-O5'-C5'	2.96	126.46	118.30
3	N	501	AMP	P-O5'-C5'	2.92	126.33	118.30
3	F	501	AMP	P-O5'-C5'	2.91	126.32	118.30
3	K	501	AMP	P-O5'-C5'	2.82	126.07	118.30

There are no chirality outliers.

5 of 54 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	501	AMP	C5'-O5'-P-O2P
3	B	501	AMP	C5'-O5'-P-O3P
3	C	501	AMP	C5'-O5'-P-O1P

*Continued on next page...*



Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	C	501	AMP	C5'-O5'-P-O2P
3	C	501	AMP	C5'-O5'-P-O3P



There are no ring outliers.



1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	501	AMP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



























## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	422/431 (97%)	-0.34	3 (0%) 87 87	17, 25, 40, 53	0
1	B	421/431 (97%)	-0.19	4 (0%) 82 81	20, 30, 47, 62	0
1	C	421/431 (97%)	-0.32	4 (0%) 82 81	14, 22, 36, 63	0
1	D	413/431 (95%)	-0.08	6 (1%) 73 71	15, 30, 56, 65	0
1	E	411/431 (95%)	-0.06	10 (2%) 59 54	16, 33, 56, 71	0
1	F	411/431 (95%)	-0.16	4 (0%) 82 81	16, 28, 48, 61	0
1	G	411/431 (95%)	0.32	31 (7%) 14 11	23, 42, 65, 76	0
1	H	416/431 (96%)	0.30	11 (2%) 56 52	28, 48, 67, 73	0
1	I	400/431 (92%)	0.41	30 (7%) 14 11	23, 45, 77, 85	0
1	J	412/431 (95%)	0.10	14 (3%) 45 41	25, 37, 61, 76	0
1	K	421/431 (97%)	-0.10	5 (1%) 79 77	20, 34, 48, 60	0
1	L	412/431 (95%)	0.15	10 (2%) 59 54	19, 36, 62, 76	0
1	M	415/431 (96%)	-0.24	1 (0%) 95 96	14, 26, 44, 55	0
1	N	422/431 (97%)	-0.27	3 (0%) 87 87	15, 25, 45, 64	0
1	O	415/431 (96%)	-0.18	8 (1%) 66 63	16, 29, 50, 63	0
1	P	423/431 (98%)	-0.35	1 (0%) 95 96	14, 23, 38, 59	0
All	All	6646/6896 (96%)	-0.07	145 (2%) 62 57	14, 31, 60, 85	0

The worst 5 of 145 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	116	GLY	6.2
1	J	104	ALA	6.2
1	H	206	PRO	4.8
1	I	255	TYR	4.6
1	B	204	ASN	4.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FRU	U	2	12/12	0.77	0.41	69,69,69,69	0
2	FRU	T	2	12/12	0.83	0.19	60,70,77,85	0
2	GLC	V	1	11/12	0.83	0.16	37,52,56,56	0
2	GLC	T	1	11/12	0.86	0.17	46,50,55,57	0
2	FRU	R	2	12/12	0.87	0.19	44,58,61,70	0
2	GLC	U	1	11/12	0.87	0.28	69,69,69,69	0
2	FRU	S	2	12/12	0.88	0.24	51,64,73,76	0
2	FRU	V	2	12/12	0.90	0.17	61,63,66,67	0
2	FRU	X	2	12/12	0.90	0.15	34,37,40,41	0
2	GLC	X	1	11/12	0.91	0.16	25,30,34,34	0
2	FRU	W	2	12/12	0.92	0.17	37,47,54,54	0
2	GLC	S	1	11/12	0.92	0.19	36,42,47,52	0
2	GLC	R	1	11/12	0.92	0.15	40,47,51,52	0
2	GLC	Q	1	11/12	0.93	0.15	22,25,27,27	0
2	GLC	W	1	11/12	0.94	0.12	28,41,46,48	0
2	FRU	Q	2	12/12	0.95	0.13	27,30,34,34	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain Q:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around Chain R:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain S:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain T:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain U:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









**Electron density around Chain V:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain W:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	PO4	D	502	5/5	0.91	0.24	55,56,57,58	0
3	AMP	H	501	23/23	0.94	0.20	40,51,53,54	0
3	AMP	K	501	23/23	0.94	0.17	25,41,47,48	0
3	AMP	B	501	23/23	0.94	0.16	28,35,36,36	0
3	AMP	G	501	23/23	0.95	0.14	34,40,44,46	0
3	AMP	I	501	23/23	0.95	0.19	33,38,41,43	0
3	AMP	J	501	23/23	0.96	0.14	28,32,34,35	0
3	AMP	E	501	23/23	0.96	0.19	22,32,34,37	0
3	AMP	L	501	23/23	0.96	0.15	24,25,26,26	0
3	AMP	O	501	23/23	0.96	0.13	22,24,26,28	0

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	AMP	P	501	23/23	0.96	0.15	23,24,25,25	0
3	AMP	A	501	23/23	0.96	0.15	20,26,30,33	0
3	AMP	M	501	23/23	0.97	0.14	21,23,24,25	0
3	AMP	N	501	23/23	0.97	0.14	21,21,22,22	0
3	AMP	D	501	23/23	0.97	0.15	26,33,34,36	0
3	AMP	C	501	23/23	0.97	0.12	18,20,21,21	0
3	AMP	F	501	23/23	0.97	0.13	24,29,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around AMP K 501:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around AMP B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AMP G 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around AMP I 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)













**Electron density around AMP E 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AMP L 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AMP O 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AMP P 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AMP A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around AMP M 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)










**Electron density around AMP D 501:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around AMP C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around AMP F 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.