



wwPDB NMR Structure Validation Summary Report ⓘ

Jun 5, 2023 – 02:35 PM JST

PDB ID : 6L8V
BMRB ID : 36294
Title : membrane-bound Bax helix2-helix5 domain
Authors : OuYang, B.; Lv, F.
Deposited on : 2019-11-07

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.33


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 18%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	81	
1	B	81	

2 Ensemble composition and analysis i

This entry contains 15 models. Model 15 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:55-A:128, B:56-B:128 (147)	1.22	15

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 5, 6, 7, 8, 9, 11, 14, 15
2	1, 12
Single-model clusters	10; 13

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2520 atoms, of which 1262 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Apoptosis regulator BAX.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	81	1260	398	631	107	120	4	0
1	B	81	1260	398	631	107	120	4	0

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	48	GLY	-	expression tag	UNP Q07812
A	49	PRO	-	expression tag	UNP Q07812
A	50	GLY	-	expression tag	UNP Q07812
A	51	SER	-	expression tag	UNP Q07812
A	52	MET	-	expression tag	UNP Q07812
A	62	SER	CYS	engineered mutation	UNP Q07812
A	126	SER	CYS	engineered mutation	UNP Q07812
B	48	GLY	-	expression tag	UNP Q07812
B	49	PRO	-	expression tag	UNP Q07812
B	50	GLY	-	expression tag	UNP Q07812
B	51	SER	-	expression tag	UNP Q07812
B	52	MET	-	expression tag	UNP Q07812
B	62	SER	CYS	engineered mutation	UNP Q07812
B	126	SER	CYS	engineered mutation	UNP Q07812

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Apoptosis regulator BAX




- Molecule 1: Apoptosis regulator BAX




4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 15. Colouring as in section 4.1 above.

- Molecule 1: Apoptosis regulator BAX



- Molecule 1: Apoptosis regulator BAX



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 15 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure calculation	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	387
Number of shifts mapped to atoms	387
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	18%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	587	593	593	4±2
1	B	581	588	588	4±2
All	All	17520	17715	17715	98

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

5 of 54 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:111:VAL:HG12	1:B:111:VAL:HG13	0.72	1.61	10	2
1:A:111:VAL:CG1	1:B:111:VAL:HG13	0.67	2.20	10	2
1:A:80:ILE:O	1:A:83:VAL:HG13	0.63	1.93	13	3
1:A:125:LEU:O	1:A:125:LEU:HD13	0.60	1.94	3	1
1:A:111:VAL:HG13	1:B:111:VAL:CG1	0.56	2.30	6	3

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	73/81 (90%)	67±2 (92±2%)	4±2 (5±3%)	2±1 (2±1%)	9	48
1	B	72/81 (89%)	66±1 (92±2%)	4±1 (6±1%)	1±1 (1±1%)	14	59
All	All	2175/2430 (90%)	2007 (92%)	127 (6%)	41 (2%)	11	53

5 of 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	84	ASP	9
1	A	83	VAL	6
1	B	84	ASP	6
1	B	73	ASN	4
1	A	87	SER	3

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	64/68 (94%)	54±3 (84±4%)	10±3 (16±4%)	5	41
1	B	63/68 (93%)	52±3 (82±4%)	11±3 (18±4%)	4	38
All	All	1905/2040 (93%)	1582 (83%)	323 (17%)	5	40

5 of 99 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	109	ARG	13
1	B	115	TYR	12
1	B	105	PHE	12
1	A	105	PHE	11
1	A	115	TYR	10

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 18% for the well-defined parts and 18% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *bax2-5CS_V3.str*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	387
Number of shifts mapped to atoms	387
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	80	0.14 ± 0.16	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	73	1.35 ± 0.16	Should be checked
$^{13}\text{C}'$	80	0.05 ± 0.22	None needed (< 0.5 ppm)
^{15}N	77	-0.09 ± 0.25	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 18%, i.e. 360 atoms were assigned a chemical shift out of a possible 2044. 0 out of 28 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	292/737 (40%)	72/298 (24%)	148/294 (50%)	72/145 (50%)
Sidechain	68/1145 (6%)	0/744 (0%)	68/351 (19%)	0/50 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	0/162 (0%)	0/80 (0%)	0/80 (0%)	0/2 (0%)
Overall	360/2044 (18%)	72/1122 (6%)	216/725 (30%)	72/197 (37%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1464
Intra-residue ($ i-j =0$)	110
Sequential ($ i-j =1$)	850
Medium range ($ i-j >1$ and $ i-j <5$)	362
Long range ($ i-j \geq 5$)	142
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	9.0
Number of long range restraints per residue ¹	0.9

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	84.1	0.2
0.2-0.5 (Medium)	60.9	0.5
>0.5 (Large)	51.2	30.32

8.2.2 Average number of dihedral-angle violations per model

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis i


9.1 Summary of distance violations i

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	110	7.5	12	10.9	0.8	0	0.0	0.0
Backbone-Backbone	40	2.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	66	4.5	12	18.2	0.8	0	0.0	0.0
Sidechain-Sidechain	4	0.3	0	0.0	0.0	0	0.0	0.0
Sequential ($i-j =1$)	850	58.1	228	26.8	15.6	24	2.8	1.6
Backbone-Backbone	496	33.9	64	12.9	4.4	12	2.4	0.8
Backbone-Sidechain	314	21.4	140	44.6	9.6	12	3.8	0.8
Sidechain-Sidechain	40	2.7	24	60.0	1.6	0	0.0	0.0
Medium range ($i-j >1$ & $i-j <5$)	362	24.7	98	27.1	6.7	8	2.2	0.5
Backbone-Backbone	208	14.2	24	11.5	1.6	0	0.0	0.0
Backbone-Sidechain	96	6.6	40	41.7	2.7	6	6.2	0.4
Sidechain-Sidechain	58	4.0	34	58.6	2.3	2	3.4	0.1
Long range ($i-j \geq 5$)	142	9.7	96	67.6	6.6	50	35.2	3.4
Backbone-Backbone	4	0.3	4	100.0	0.3	4	100.0	0.3
Backbone-Sidechain	42	2.9	34	81.0	2.3	26	61.9	1.8
Sidechain-Sidechain	96	6.6	58	60.4	4.0	20	20.8	1.4
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1464	100.0	434	29.6	29.6	82	5.6	5.6
Backbone-Backbone	748	51.1	92	12.3	6.3	16	2.1	1.1
Backbone-Sidechain	518	35.4	226	43.6	15.4	44	8.5	3.0
Sidechain-Sidechain	198	13.5	116	58.6	7.9	22	11.1	1.5

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	4	84	30	66	0	184	3.89	28.53	7.21	0.22
2	0	92	26	56	0	174	3.97	29.45	7.36	0.25
3	0	116	26	66	0	208	3.43	28.88	6.9	0.24
4	0	100	18	62	0	180	3.97	30.32	7.39	0.24
5	0	124	22	68	0	214	3.34	29.2	6.85	0.21
6	0	112	42	74	0	228	3.16	28.17	6.63	0.21
7	0	100	22	62	0	184	3.78	28.91	7.2	0.24
8	0	112	26	62	0	200	3.46	29.6	6.94	0.2
9	0	96	26	68	0	190	3.78	29.97	7.25	0.25
10	0	88	46	62	0	196	3.63	27.45	6.92	0.2
11	0	120	30	58	0	208	3.37	29.63	6.91	0.23


Continued on next page...

Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
12	4	92	32	60	0	188	3.82	28.16	7.19	0.22
13	0	92	36	62	0	190	3.74	27.66	7.0	0.22
14	0	104	30	76	0	210	3.34	29.54	6.84	0.24
15	4	96	22	68	0	190	3.65	29.56	7.06	0.23

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble [i](#)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints

for a given fraction of the ensemble. In total, 1030(IR:98, SQ:622, MR:264, LR:46, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
12	48	28	6	0	94	1	6.7
0	20	24	0	0	44	2	13.3
0	20	16	6	0	42	3	20.0
0	8	0	6	0	14	4	26.7
0	16	4	12	0	32	5	33.3
0	4	4	4	0	12	6	40.0
0	16	0	8	0	24	7	46.7
0	16	4	4	0	24	8	53.3
0	12	4	0	0	16	9	60.0
0	12	0	0	0	12	10	66.7
0	0	0	0	0	0	11	73.3
0	12	0	0	0	12	12	80.0
0	8	6	0	0	14	13	86.7
0	12	0	0	0	12	14	93.3
0	24	8	50	0	82	15	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG21	15	29.0	0.82	29.2
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG22	15	29.0	0.82	29.2
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG23	15	29.0	0.82	29.2
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG21	15	29.0	0.82	29.2
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG22	15	29.0	0.82	29.2
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG23	15	29.0	0.82	29.2
(1,37)	1:A:95:VAL:H	1:A:56:THR:HG21	15	28.37	0.83	28.53
(1,37)	1:A:95:VAL:H	1:A:56:THR:HG22	15	28.37	0.83	28.53
(1,37)	1:A:95:VAL:H	1:A:56:THR:HG23	15	28.37	0.83	28.53
(1,38)	1:A:95:VAL:H	1:A:56:THR:HG21	15	28.37	0.83	28.53
(1,38)	1:A:95:VAL:H	1:A:56:THR:HG22	15	28.37	0.83	28.53
(1,38)	1:A:95:VAL:H	1:A:56:THR:HG23	15	28.37	0.83	28.53
(1,5)	1:A:60:SER:H	1:A:95:VAL:HG21	15	22.57	0.46	22.72
(1,5)	1:A:60:SER:H	1:A:95:VAL:HG22	15	22.57	0.46	22.72
(1,5)	1:A:60:SER:H	1:A:95:VAL:HG23	15	22.57	0.46	22.72
(1,6)	1:A:60:SER:H	1:A:95:VAL:HG21	15	22.57	0.46	22.72

Continued on next page...

Continued from previous page...


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,6)	1:A:60:SER:H	1:A:95:VAL:HG22	15	22.57	0.46	22.72
(1,6)	1:A:60:SER:H	1:A:95:VAL:HG23	15	22.57	0.46	22.72
(1,3)	1:A:60:SER:H	1:A:95:VAL:HG11	15	21.58	0.88	21.41
(1,3)	1:A:60:SER:H	1:A:95:VAL:HG12	15	21.58	0.88	21.41
(1,3)	1:A:60:SER:H	1:A:95:VAL:HG13	15	21.58	0.88	21.41
(1,4)	1:A:60:SER:H	1:A:95:VAL:HG11	15	21.58	0.88	21.41
(1,4)	1:A:60:SER:H	1:A:95:VAL:HG12	15	21.58	0.88	21.41
(1,4)	1:A:60:SER:H	1:A:95:VAL:HG13	15	21.58	0.88	21.41
(1,1)	1:A:56:THR:HG21	1:A:95:VAL:HG21	15	20.69	0.65	20.75
(1,1)	1:A:56:THR:HG21	1:A:95:VAL:HG22	15	20.69	0.65	20.75
(1,1)	1:A:56:THR:HG21	1:A:95:VAL:HG23	15	20.69	0.65	20.75
(1,1)	1:A:56:THR:HG22	1:A:95:VAL:HG21	15	20.69	0.65	20.75
(1,1)	1:A:56:THR:HG22	1:A:95:VAL:HG22	15	20.69	0.65	20.75
(1,1)	1:A:56:THR:HG22	1:A:95:VAL:HG23	15	20.69	0.65	20.75
(1,1)	1:A:56:THR:HG23	1:A:95:VAL:HG21	15	20.69	0.65	20.75
(1,1)	1:A:56:THR:HG23	1:A:95:VAL:HG22	15	20.69	0.65	20.75
(1,1)	1:A:56:THR:HG23	1:A:95:VAL:HG23	15	20.69	0.65	20.75
(1,2)	1:A:56:THR:HG21	1:A:95:VAL:HG21	15	20.69	0.65	20.75

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG21	4	30.32
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG22	4	30.32
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG23	4	30.32
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG21	4	30.32
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG22	4	30.32
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG23	4	30.32
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG21	9	29.97
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG22	9	29.97
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG23	9	29.97
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG21	9	29.97
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG22	9	29.97
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG23	9	29.97
(1,38)	1:A:95:VAL:H	1:A:56:THR:HG21	4	29.67
(1,38)	1:A:95:VAL:H	1:A:56:THR:HG22	4	29.67
(1,38)	1:A:95:VAL:H	1:A:56:THR:HG23	4	29.67
(1,37)	1:A:95:VAL:H	1:A:56:THR:HG21	4	29.67
(1,37)	1:A:95:VAL:H	1:A:56:THR:HG22	4	29.67

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,37)	1:A:95:VAL:H	1:A:56:THR:HG23	4	29.67
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG21	11	29.63
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG22	11	29.63
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG23	11	29.63
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG21	11	29.63
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG22	11	29.63
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG23	11	29.63
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG21	8	29.6
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG22	8	29.6
(1,34)	1:A:94:ARG:H	1:A:56:THR:HG23	8	29.6
(1,33)	1:A:94:ARG:H	1:A:56:THR:HG21	8	29.6

10 Dihedral-angle violation analysis

No dihedral-angle restraints found