

wwPDB X-ray Structure Validation Summary Report (i)

Sep 11, 2023 – 02:12 PM EDT

PDB ID	:	4L9Z
Title	:	Crystal Structure of Rhodobacter sphaeroides malyl-CoA lyase in complex
		with magnesium, oxalate, and CoA
Authors	:	Zarzycki, J.; Kerfeld, C.A.
Deposited on	:	2013-06-18
Resolution	:	2.01 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)		
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	$10434 \ (2.04-2.00)$
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	339	88%	5% 8%
1	В	339	% 	•• 7%
1	С	339	88%	• 7%
1	D	339	% 	5% 7%
1	Е	339	% 91%	• 8%



Mol	Chain	Length	Quality of chain		
			.%		
1	F	339	88%	•	8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OXL	А	402	-	Х	-	-
3	OXL	В	402	-	Х	-	-
3	OXL	С	402	-	Х	-	-
3	OXL	Е	402	-	Х	-	-
3	OXL	F	402	-	Х	-	-



$\mathbf{2}$ Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 17160 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace	
1	А	313	Total	С	Ν	Ο	\mathbf{S}	0	1	0	
	A	515	2381	1497	424	447	13	0	L	0	
1	В	314	Total	С	Ν	Ο	S	0	1	0	
	D	514	2387	1500	425	449	13	0	T	U	
1	С	314	Total	С	Ν	Ο	S	0	1	0	
	U	514	2387	1500	425	449	13	0		0	
1	D	314	Total	С	Ν	Ο	S	0	1	0	
	D	514	2387	1500	425	449	13	0	T		
1	Е	313	Total	С	Ν	Ο	S	0	1	0	
	Ľ	515	2381	1497	424	447	13	0	T	0	
1	F	313	Total	С	Ν	Ο	S	0	1	0	
	Г	515	2381	1497	424	447	13	0			

• Molecule 1 is a protein called Malyl-CoA lyase.

There are 126 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-20	MET	-	expression tag	UNP Q3J5L6
А	-19	GLY	-	expression tag	UNP Q3J5L6
А	-18	HIS	-	expression tag	UNP Q3J5L6
А	-17	HIS	-	expression tag	UNP Q3J5L6
А	-16	HIS	-	expression tag	UNP Q3J5L6
А	-15	HIS	-	expression tag	UNP Q3J5L6
А	-14	HIS	-	expression tag	UNP Q3J5L6
A	-13	HIS	-	expression tag	UNP Q3J5L6
А	-12	HIS	-	expression tag	UNP Q3J5L6
A	-11	HIS	-	expression tag	UNP Q3J5L6
А	-10	HIS	-	expression tag	UNP Q3J5L6
А	-9	HIS	-	expression tag	UNP Q3J5L6
А	-8	SER	-	expression tag	UNP Q3J5L6
А	-7	SER	-	expression tag	UNP Q3J5L6
А	-6	GLY	-	expression tag	UNP Q3J5L6
А	-5	HIS	-	expression tag	UNP Q3J5L6
А	-4	ILE	-	expression tag	UNP Q3J5L6



Chain	Residue	vious page Modelled	Actual	Comment	Reference
A	-3	GLU	-	expression tag	UNP Q3J5L6
A	-2	GLY	-	expression tag	UNP Q3J5L6
A	-1	ARG	_	expression tag	UNP Q3J5L6
A	0	HIS	-	expression tag	UNP Q3J5L6
В	-20	MET	_	expression tag	UNP Q3J5L6
В	-19	GLY	-	expression tag	UNP Q3J5L6
В	-18	HIS	-	expression tag	UNP Q3J5L6
В	-17	HIS	-	expression tag	UNP Q3J5L6
В	-16	HIS	-	expression tag	UNP Q3J5L6
В	-15	HIS	_	expression tag	UNP Q3J5L6
В	-14	HIS	_	expression tag	UNP Q3J5L6
В	-13	HIS	_	expression tag	UNP Q3J5L6
В	-12	HIS	_	expression tag	UNP Q3J5L6
В	-11	HIS	_	expression tag	UNP Q3J5L6
В	-10	HIS	_	expression tag	UNP Q3J5L6
В	-9	HIS	_	expression tag	UNP Q3J5L6
В	-8	SER	-	expression tag	UNP Q3J5L6
В	-7	SER	-	expression tag	UNP Q3J5L6
В	-6	GLY	-	expression tag	UNP Q3J5L6
В	-5	HIS	-	expression tag	UNP Q3J5L6
В	-4	ILE	-	expression tag	UNP Q3J5L6
В	-3	GLU	-	expression tag	UNP Q3J5L6
В	-2	GLY	-	expression tag	UNP Q3J5L6
В	-1	ARG	-	expression tag	UNP Q3J5L6
В	0	HIS	-	expression tag	UNP Q3J5L6
С	-20	MET	-	expression tag	UNP Q3J5L6
С	-19	GLY	-	expression tag	UNP Q3J5L6
С	-18	HIS	-	expression tag	UNP Q3J5L6
С	-17	HIS	-	expression tag	UNP Q3J5L6
С	-16	HIS	-	expression tag	UNP Q3J5L6
С	-15	HIS	-	expression tag	UNP Q3J5L6
С	-14	HIS	-	expression tag	UNP Q3J5L6
С	-13	HIS	-	expression tag	UNP Q3J5L6
С	-12	HIS	-	expression tag	UNP Q3J5L6
С	-11	HIS	-	expression tag	UNP Q3J5L6
С	-10	HIS	-	expression tag	UNP Q3J5L6
С	-9	HIS	-	expression tag	UNP Q3J5L6
С	-8	SER	-	expression tag	UNP Q3J5L6
С	-7	SER	-	expression tag	UNP Q3J5L6
С	-6	GLY	-	expression tag	UNP Q3J5L6
С	-5	HIS	-	expression tag	UNP Q3J5L6
С	-4	ILE	-	expression tag	UNP Q3J5L6




Chain	Residue	Modelled	Actual	Comment	Reference
С	-3	GLU	-	expression tag	UNP Q3J5L6
С	-2	GLY	-	expression tag	UNP Q3J5L6
С	-1	ARG	-	expression tag	UNP Q3J5L6
С	0	HIS	-	expression tag	UNP Q3J5L6
D	-20	MET	-	expression tag	UNP Q3J5L6
D	-19	GLY	-	expression tag	UNP Q3J5L6
D	-18	HIS	-	expression tag	UNP Q3J5L6
D	-17	HIS	-	expression tag	UNP Q3J5L6
D	-16	HIS	-	expression tag	UNP Q3J5L6
D	-15	HIS	-	expression tag	UNP Q3J5L6
D	-14	HIS	-	expression tag	UNP Q3J5L6
D	-13	HIS	-	expression tag	UNP Q3J5L6
D	-12	HIS	-	expression tag	UNP Q3J5L6
D	-11	HIS	-	expression tag	UNP Q3J5L6
D	-10	HIS	-	expression tag	UNP Q3J5L6
D	-9	HIS	-	expression tag	UNP Q3J5L6
D	-8	SER	-	expression tag	UNP Q3J5L6
D	-7	SER	-	expression tag	UNP Q3J5L6
D	-6	GLY	-	expression tag	UNP Q3J5L6
D	-5	HIS	-	expression tag	UNP Q3J5L6
D	-4	ILE	-	expression tag	UNP Q3J5L6
D	-3	GLU	-	expression tag	UNP Q3J5L6
D	-2	GLY	-	expression tag	UNP Q3J5L6
D	-1	ARG	-	expression tag	UNP Q3J5L6
D	0	HIS	-	expression tag	UNP Q3J5L6
Е	-20	MET	-	expression tag	UNP Q3J5L6
Е	-19	GLY	-	expression tag	UNP Q3J5L6
Е	-18	HIS	-	expression tag	UNP Q3J5L6
Е	-17	HIS	-	expression tag	UNP Q3J5L6
Е	-16	HIS	-	expression tag	UNP Q3J5L6
Е	-15	HIS	-	expression tag	UNP Q3J5L6
Е	-14	HIS	-	expression tag	UNP Q3J5L6
Е	-13	HIS	-	expression tag	UNP Q3J5L6
Е	-12	HIS	-	expression tag	UNP Q3J5L6
Е	-11	HIS	-	expression tag	UNP Q3J5L6
Е	-10	HIS	-	expression tag	UNP Q3J5L6
Е	-9	HIS	-	expression tag	UNP Q3J5L6
Е	-8	SER	-	expression tag	UNP Q3J5L6
Е	-7	SER	-	expression tag	UNP Q3J5L6
Е	-6	GLY	-	expression tag	UNP Q3J5L6
Е	-5	HIS	-	expression tag	UNP Q3J5L6
Ε	-4	ILE	-	expression tag	UNP Q3J5L6



Chain	Residue	Modelled	Actual	Comment	Reference
Е	-3	GLU	-	expression tag	UNP Q3J5L6
Е	-2	GLY	-	expression tag	UNP Q3J5L6
Е	-1	ARG	-	expression tag	UNP Q3J5L6
Е	0	HIS	-	expression tag	UNP Q3J5L6
F	-20	MET	-	expression tag	UNP Q3J5L6
F	-19	GLY	-	expression tag	UNP Q3J5L6
F	-18	HIS	-	expression tag	UNP Q3J5L6
F	-17	HIS	-	expression tag	UNP Q3J5L6
F	-16	HIS	-	expression tag	UNP Q3J5L6
F	-15	HIS	-	expression tag	UNP Q3J5L6
F	-14	HIS	-	expression tag	UNP Q3J5L6
F	-13	HIS	-	expression tag	UNP Q3J5L6
F	-12	HIS	-	expression tag	UNP Q3J5L6
F	-11	HIS	-	expression tag	UNP Q3J5L6
F	-10	HIS	-	expression tag	UNP Q3J5L6
F	-9	HIS	-	expression tag	UNP Q3J5L6
F	-8	SER	-	expression tag	UNP Q3J5L6
F	-7	SER	-	expression tag	UNP Q3J5L6
F	-6	GLY	-	expression tag	UNP Q3J5L6
F	-5	HIS	-	expression tag	UNP Q3J5L6
F	-4	ILE	-	expression tag	UNP Q3J5L6
F	-3	GLU	-	expression tag	UNP Q3J5L6
F	-2	GLY	-	expression tag	UNP Q3J5L6
F	-1	ARG	-	expression tag	UNP Q3J5L6
F	0	HIS	-	expression tag	UNP Q3J5L6


• Molecule 2 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
2	А	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
	Π	1	48	21	7	16	3	1	0	0	
2	В	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
2	D	1	48	21	7	16	3	1	0	0	
2	C	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
2	U	L	48	21	7	16	3	1	0	0	
2	О	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
2	D	1	48	21	7	16	3	1	0	0	
2	Е	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
2	Ľ	1	48	21	7	16	3	1	0	0	
2	F	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
	Ľ	1	48	21	7	16	3	1	0	0	

• Molecule 3 is OXALATE ION (three-letter code: OXL) (formula: C_2O_4).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 2 & 4 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 2 & 4 \end{array}$	0	0
3	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 2 4 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 2 & 4 \end{array}$	0	0
3	Е	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 2 & 4 \end{array}$	0	0
3	F	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 2 & 4 \end{array}$	0	0



- 4L9Z
- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total Mg 1 1	0	0
4	В	1	Total Mg 1 1	0	0
4	С	1	Total Mg 1 1	0	0
4	D	1	Total Mg 1 1	0	0
4	Ε	1	Total Mg 1 1	0	0
4	F	1	Total Mg 1 1	0	0


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	449	Total O 449 449	0	0
5	В	417	Total O 417 417	0	0
5	С	431	Total O 431 431	0	0
5	D	409	Total O 409 409	0	0
5	Е	410	Total O 410 410	0	0
5	F	410	Total O 410 410	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Malyl-CoA lyase

• Molecule 1: Malyl-CoA lyase







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61	Depositor
Cell constants	221.51Å 221.51 Å 96.26 Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.45 - 2.01	Depositor
Resolution (A)	38.45 - 2.01	EDS
% Data completeness	91.9 (38.45-2.01)	Depositor
(in resolution range)	85.8 (38.45-2.01)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	0.16	Depositor
$< I/\sigma(I) > 1$	$1.56 (at 2.01 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.8.2_1303	Depositor
D D.	0.170 , 0.194	Depositor
R, R_{free}	0.168 , 0.192	DCC
R_{free} test set	1740 reflections $(1.14%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	18.4	Xtriage
Anisotropy	0.099	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 47.4	EDS
L-test for twinning ²	$< L > = 0.51, < L^2 > = 0.35$	Xtriage
Estimated twinning fraction	0.009 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	17160	wwPDB-VP
Average B, all atoms $(Å^2)$	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.40% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OXL, MG, COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
	Ullalli	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.23	0/2433	0.42	0/3302
1	В	0.25	0/2439	0.42	0/3310
1	С	0.22	0/2439	0.41	0/3310
1	D	0.25	0/2439	0.42	0/3310
1	Ε	0.27	0/2433	0.42	0/3302
1	F	0.22	0/2433	0.41	0/3302
All	All	0.24	0/14616	0.41	0/19836

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2381	0	2354	9	0
1	В	2387	0	2359	11	0
1	С	2387	0	2359	9	0
1	D	2387	0	2359	10	0
1	Е	2381	0	2354	3	0
1	F	2381	0	2354	10	0
2	А	48	0	32	1	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	48	0	32	0	0
2	С	48	0	32	0	0
2	D	48	0	32	0	0
2	Е	48	0	32	0	0
2	F	48	0	32	0	0
3	А	6	0	0	0	0
3	В	6	0	0	0	0
3	С	6	0	0	0	0
3	D	6	0	0	0	0
3	Ε	6	0	0	0	0
3	F	6	0	0	0	0
4	А	1	0	0	0	0
4	В	1	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
4	Е	1	0	0	0	0
4	F	1	0	0	0	0
5	А	449	0	0	4	1
5	В	417	0	0	5	3
5	С	431	0	0	4	0
5	D	409	0	0	5	1
5	Е	410	0	0	1	1
5	F	410	0	0	6	0
All	All	17160	0	14331	50	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 50 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:5:LEU:HD23	5:B:858:HOH:O	1.68	0.92
1:C:81:ASP:OD2	5:C:928:HOH:O	2.00	0.79
1:C:315:MET:SD	5:C:712:HOH:O	2.41	0.79
1:C:311:ARG:O	5:C:893:HOH:O	2.04	0.74
1:D:2:SER:N	5:D:886:HOH:O	2.21	0.72

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:800:HOH:O	5:E:826:HOH:O[1_556]	2.06	0.14
5:B:618:HOH:O	5:D:727:HOH:O[1_556]	2.12	0.08
5:A:921:HOH:O	5:B:773:HOH:O[5_554]	2.14	0.06

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentile	s
1	А	312/339~(92%)	305~(98%)	7(2%)	0	100 100	
1	В	313/339~(92%)	307~(98%)	6~(2%)	0	100 100	
1	С	313/339~(92%)	307~(98%)	6 (2%)	0	100 100	
1	D	313/339~(92%)	307~(98%)	6(2%)	0	100 100	
1	Ε	312/339~(92%)	305~(98%)	7(2%)	0	100 100	
1	F	312/339~(92%)	306~(98%)	6(2%)	0	100 100	
All	All	1875/2034~(92%)	1837 (98%)	38 (2%)	0	100 100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	239/260~(92%)	235~(98%)	4 (2%)	60 63
1	В	240/260~(92%)	235~(98%)	5 (2%)	53 55
1	С	240/260~(92%)	237~(99%)	3 (1%)	69 72



Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	D	240/260~(92%)	238~(99%)	2(1%)	81 85		
1	Ε	239/260~(92%)	237~(99%)	2(1%)	81 85		
1	F	239/260~(92%)	238 (100%)	1 (0%)	91 93		
All	All	1437/1560~(92%)	1420 (99%)	17 (1%)	71 75		

Continued from previous page...

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	14	ASN
1	F	14	ASN
1	В	194	GLN
1	В	315	MET
1	С	5	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	205	GLN
1	F	312	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 6 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	gles
MOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OXL	F	402	4	$5,\!5,\!5$	1.54	0	$6,\!6,\!6$	1.42	2 (33%)
3	OXL	Е	402	4	$5,\!5,\!5$	1.54	0	$6,\!6,\!6$	1.46	2 (33%)
2	COA	С	401	-	41,50,50	1.47	6 (14%)	52,75,75	1.46	9 (17%)
2	COA	Е	401	-	41,50,50	1.43	6 (14%)	52,75,75	1.24	<mark>6 (11%)</mark>
2	COA	D	401	-	41,50,50	1.56	7 (17%)	52,75,75	1.16	5 (9%)
2	COA	F	401	-	41,50,50	1.27	4 (9%)	52,75,75	1.45	10 (19%)
3	OXL	В	402	4	$5,\!5,\!5$	1.45	0	$6,\!6,\!6$	1.66	2 (33%)
3	OXL	D	402	4	$5,\!5,\!5$	1.57	0	$6,\!6,\!6$	1.42	1 (16%)
3	OXL	А	402	4	$5,\!5,\!5$	1.45	0	$6,\!6,\!6$	1.70	2 (33%)
2	COA	А	401	-	41,50,50	1.28	3 (7%)	52,75,75	1.36	7 (13%)
2	COA	В	401	-	41,50,50	0.96	1 (2%)	52,75,75	1.05	2 (3%)
3	OXL	С	402	4	$5,\!5,\!5$	1.46	0	$6,\!6,\!6$	1.60	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OXL	F	402	4	-	4/4/4/4	-
3	OXL	Е	402	4	-	4/4/4/4	-
2	COA	С	401	-	-	3/44/64/64	0/3/3/3
2	COA	Е	401	-	-	3/44/64/64	0/3/3/3
2	COA	D	401	-	-	5/44/64/64	0/3/3/3
2	COA	F	401	-	-	6/44/64/64	0/3/3/3
3	OXL	В	402	4	-	4/4/4/4	-
3	OXL	D	402	4	-	4/4/4/4	-
3	OXL	А	402	4	-	4/4/4/4	-
2	COA	А	401	-	-	8/44/64/64	0/3/3/3
2	COA	В	401	-	-	2/44/64/64	0/3/3/3
3	OXL	С	402	4	-	4/4/4/4	-



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	С	401	COA	C4A-N3A	-4.09	1.30	1.35
2	Е	401	COA	P3B-O9A	-3.79	1.40	1.54
2	D	401	COA	P3B-O9A	-3.70	1.40	1.54
2	Е	401	COA	P1A-O1A	-3.49	1.38	1.50
2	В	401	COA	P3B-O7A	3.41	1.61	1.50

The worst 5 of 27 bond length outliers are listed below:

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	401	COA	N3A-C2A-N1A	-4.90	121.01	128.68
2	С	401	COA	N3A-C2A-N1A	-4.78	121.21	128.68
2	D	401	COA	N3A-C2A-N1A	-3.86	122.65	128.68
2	С	401	COA	C7P-N8P-C9P	3.85	129.45	122.59
2	А	401	COA	C6P-C5P-N4P	-3.67	110.24	116.42

There are no chirality outliers.

5 of 51 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	401	COA	C3B-O3B-P3B-O9A
2	А	401	COA	C5B-O5B-P1A-O1A
2	А	401	COA	N8P-C9P-CAP-OAP
2	А	401	COA	CAP-C9P-N8P-C7P
2	С	401	COA	C5B-O5B-P1A-O1A

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	COA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.




The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(A^2)$	Q<0.9
1	А	313/339~(92%)	-0.71	1 (0%) 94 93	4, 10, 27, 53	0
1	В	314/339~(92%)	-0.57	2 (0%) 89 89	5, 11, 31, 47	0
1	С	314/339~(92%)	-0.68	3 (0%) 82 82	4, 12, 25, 41	0
1	D	314/339~(92%)	-0.57	4 (1%) 77 76	4, 11, 28, 50	0
1	Ε	313/339~(92%)	-0.46	4 (1%) 77 76	5, 11, 33, 49	0
1	F	313/339~(92%)	-0.52	4 (1%) 77 76	4, 11, 28, 51	0
All	All	1881/2034~(92%)	-0.59	18 (0%) 82 82	4, 11, 29, 53	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	5	LEU	3.9
1	Ε	4	ARG	3.7
1	D	5	LEU	3.7
1	В	5	LEU	3.0
1	С	4	ARG	2.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	COA	А	401	48/48	0.89	0.14	$13,\!30,\!41,\!50$	48
2	COA	F	401	48/48	0.91	0.13	11,30,45,50	48
2	COA	С	401	48/48	0.92	0.12	16,27,40,44	48
3	OXL	В	402	6/6	0.93	0.17	7,9,19,22	6
2	COA	D	401	48/48	0.94	0.11	14,24,39,45	48
2	COA	Е	401	48/48	0.94	0.12	$16,\!25,\!38,\!44$	0
2	COA	В	401	48/48	0.95	0.10	12,21,31,41	48
3	OXL	С	402	6/6	0.96	0.14	$8,\!11,\!16,\!17$	6
3	OXL	D	402	6/6	0.96	0.12	7,8,16,16	0
3	OXL	F	402	6/6	0.96	0.11	$7,\!11,\!14,\!21$	0
3	OXL	Е	402	6/6	0.97	0.08	$10,\!10,\!16,\!17$	0
3	OXL	А	402	6/6	0.97	0.09	7,7,12,13	0
4	MG	D	403	1/1	0.97	0.10	9,9,9,9	0
4	MG	Е	403	1/1	0.97	0.07	8,8,8,8	0
4	MG	В	403	1/1	0.98	0.12	9,9,9,9	0
4	MG	F	403	1/1	0.98	0.08	9,9,9,9	0
4	MG	С	403	1/1	0.99	0.15	11,11,11,11	0
4	MG	А	403	1/1	0.99	0.06	$6,\!6,\!6,\!6$	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















6.5 Other polymers (i)

There are no such residues in this entry.

