

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 31, 2023 – 01:04 PM EDT

PDB ID : 3LBE

Title : The Crystal Structure of smu.793 from Streptococcus mutans UA159 bound

to acetyl CoA

Authors: Su, X.-D.; Hou, Q.M.; Fan, X.X.; Nan, J.; Liu, X.

Deposited on : 2010-01-08

Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	163	68%	5%	•	25%		
1	В	163	% <b>6</b> 7%	7%		24%		
1	С	163	% <b>6</b> 9%	5%		25%		
1	D	163	66%	8%	•	25%		

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard



residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	131	-	-	X	-



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Putative uncharacterized protein smu.793.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace			
1	Λ	122	Total	С	N	О	S	0	2	0
1	A	122	963	612	162	185	4	0		U
1	В	124	Total	С	N	О	S	0	7	0
1	Ъ	124	1004	638	169	193	4			U
1	С	122	Total	С	N	О	S	0	2	0
1		122	963	612	162	185	4	0	2	U
1	D	123	Total	С	N	О	S	0	4	0
1	ש	123	979	624	163	188	4			U

There are 136 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	EXPRESSION TAG	UNP Q8DUV0
A	-32	GLY	-	EXPRESSION TAG	UNP Q8DUV0
A	-31	SER	-	EXPRESSION TAG	UNP Q8DUV0
A	-30	SER	-	EXPRESSION TAG	UNP Q8DUV0
A	-29	HIS	-	EXPRESSION TAG	UNP Q8DUV0
A	-28	HIS	-	EXPRESSION TAG	UNP Q8DUV0
A	-27	HIS	-	EXPRESSION TAG	UNP Q8DUV0
A	-26	HIS	-	EXPRESSION TAG	UNP Q8DUV0
A	-25	HIS	-	EXPRESSION TAG	UNP Q8DUV0
A	-24	HIS	-	EXPRESSION TAG	UNP Q8DUV0
A	-23	SER	-	EXPRESSION TAG	UNP Q8DUV0
A	-22	SER	-	EXPRESSION TAG	UNP Q8DUV0
A	-21	GLY	-	EXPRESSION TAG	UNP Q8DUV0
A	-20	LEU	-	EXPRESSION TAG	UNP Q8DUV0
A	-19	VAL	-	EXPRESSION TAG	UNP Q8DUV0
A	-18	PRO	-	EXPRESSION TAG	UNP Q8DUV0
A	-17	ARG	=	EXPRESSION TAG	UNP Q8DUV0
A	-16	GLY	=	EXPRESSION TAG	UNP Q8DUV0
A	-15	SER	=	EXPRESSION TAG	UNP Q8DUV0
A	-14	HIS	-	EXPRESSION TAG	UNP Q8DUV0
A	-13	MET	-	EXPRESSION TAG	UNP Q8DUV0



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	ALA	-	EXPRESSION TAG	UNP Q8DUV0
A	-11	SER	-	EXPRESSION TAG	UNP Q8DUV0
A	-10	MET	-	EXPRESSION TAG	UNP Q8DUV0
A	-9	THR	-	EXPRESSION TAG	UNP Q8DUV0
A	-8	GLY	-	EXPRESSION TAG	UNP Q8DUV0
A	-7	GLY	-	EXPRESSION TAG	UNP Q8DUV0
A	-6	GLN	-	EXPRESSION TAG	UNP Q8DUV0
A	-5	GLN	-	EXPRESSION TAG	UNP Q8DUV0
A	-4	MET	-	EXPRESSION TAG	UNP Q8DUV0
A	-3	GLY	-	EXPRESSION TAG	UNP Q8DUV0
A	-2	ARG	-	EXPRESSION TAG	UNP Q8DUV0
A	-1	GLY	-	EXPRESSION TAG	UNP Q8DUV0
A	0	SER	-	EXPRESSION TAG	UNP Q8DUV0
В	-33	MET	-	EXPRESSION TAG	UNP Q8DUV0
В	-32	GLY	-	EXPRESSION TAG	UNP Q8DUV0
В	-31	SER	-	EXPRESSION TAG	UNP Q8DUV0
В	-30	SER	-	EXPRESSION TAG	UNP Q8DUV0
В	-29	HIS	-	EXPRESSION TAG	UNP Q8DUV0
В	-28	HIS	-	EXPRESSION TAG	UNP Q8DUV0
В	-27	HIS	-	EXPRESSION TAG	UNP Q8DUV0
В	-26	HIS	-	EXPRESSION TAG	UNP Q8DUV0
В	-25	HIS	-	EXPRESSION TAG	UNP Q8DUV0
В	-24	HIS	ı	EXPRESSION TAG	UNP Q8DUV0
В	-23	SER	-	EXPRESSION TAG	UNP Q8DUV0
В	-22	SER	-	EXPRESSION TAG	UNP Q8DUV0
В	-21	GLY	-	EXPRESSION TAG	UNP Q8DUV0
В	-20	LEU	-	EXPRESSION TAG	UNP Q8DUV0
В	-19	VAL	ı	EXPRESSION TAG	UNP Q8DUV0
В	-18	PRO	I	EXPRESSION TAG	UNP Q8DUV0
В	-17	ARG	-	EXPRESSION TAG	UNP Q8DUV0
В	-16	GLY	ı	EXPRESSION TAG	UNP Q8DUV0
В	-15	SER	ı	EXPRESSION TAG	UNP Q8DUV0
В	-14	HIS	-	EXPRESSION TAG	UNP Q8DUV0
В	-13	MET	ı	EXPRESSION TAG	UNP Q8DUV0
В	-12	ALA	-	EXPRESSION TAG	UNP Q8DUV0
В	-11	SER	-	EXPRESSION TAG	UNP Q8DUV0
В	-10	MET	-	EXPRESSION TAG	UNP Q8DUV0
В	-9	THR	-	EXPRESSION TAG	UNP Q8DUV0
В	-8	GLY	-	EXPRESSION TAG	UNP Q8DUV0
В	-7	GLY	-	EXPRESSION TAG	UNP Q8DUV0
В	-6	GLN	-	EXPRESSION TAG	UNP Q8DUV0
В	-5	GLN	-	EXPRESSION TAG	UNP Q8DUV0



Continued from previous page...

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
В	-4	MET	-	EXPRESSION TAG	UNP Q8DUV0
В	-3	GLY	-	EXPRESSION TAG	UNP Q8DUV0
В	-2	ARG	-	EXPRESSION TAG	UNP Q8DUV0
В	-1	GLY	-	EXPRESSION TAG	UNP Q8DUV0
В	0	SER	-	EXPRESSION TAG	UNP Q8DUV0
С	-33	MET	-	EXPRESSION TAG	UNP Q8DUV0
С	-32	GLY	-	EXPRESSION TAG	UNP Q8DUV0
С	-31	SER	-	EXPRESSION TAG	UNP Q8DUV0
С	-30	SER	-	EXPRESSION TAG	UNP Q8DUV0
С	-29	HIS	-	EXPRESSION TAG	UNP Q8DUV0
С	-28	HIS	-	EXPRESSION TAG	UNP Q8DUV0
С	-27	HIS	-	EXPRESSION TAG	UNP Q8DUV0
С	-26	HIS	-	EXPRESSION TAG	UNP Q8DUV0
С	-25	HIS	-	EXPRESSION TAG	UNP Q8DUV0
С	-24	HIS	-	EXPRESSION TAG	UNP Q8DUV0
С	-23	SER	-	EXPRESSION TAG	UNP Q8DUV0
С	-22	SER	-	EXPRESSION TAG	UNP Q8DUV0
С	-21	GLY	-	EXPRESSION TAG	UNP Q8DUV0
С	-20	LEU	-	EXPRESSION TAG	UNP Q8DUV0
С	-19	VAL	-	EXPRESSION TAG	UNP Q8DUV0
С	-18	PRO	-	EXPRESSION TAG	UNP Q8DUV0
С	-17	ARG	-	EXPRESSION TAG	UNP Q8DUV0
С	-16	GLY	-	EXPRESSION TAG	UNP Q8DUV0
С	-15	SER	-	EXPRESSION TAG	UNP Q8DUV0
С	-14	HIS	-	EXPRESSION TAG	UNP Q8DUV0
С	-13	MET	-	EXPRESSION TAG	UNP Q8DUV0
С	-12	ALA	-	EXPRESSION TAG	UNP Q8DUV0
С	-11	SER	-	EXPRESSION TAG	UNP Q8DUV0
С	-10	MET	-	EXPRESSION TAG	UNP Q8DUV0
С	-9	THR	-	EXPRESSION TAG	UNP Q8DUV0
С	-8	GLY	-	EXPRESSION TAG	UNP Q8DUV0
С	-7	GLY	-	EXPRESSION TAG	UNP Q8DUV0
С	-6	GLN	-	EXPRESSION TAG	UNP Q8DUV0
С	-5	GLN	-	EXPRESSION TAG	UNP Q8DUV0
С	-4	MET	-	EXPRESSION TAG	UNP Q8DUV0
С	-3	GLY	-	EXPRESSION TAG	UNP Q8DUV0
С	-2	ARG	-	EXPRESSION TAG	UNP Q8DUV0
С	-1	GLY	-	EXPRESSION TAG	UNP Q8DUV0
С	0	SER	-	EXPRESSION TAG	UNP Q8DUV0
D	-33	MET	-	EXPRESSION TAG	UNP Q8DUV0
D	-32	GLY	-	EXPRESSION TAG	UNP Q8DUV0
D	-31	SER	-	EXPRESSION TAG	UNP Q8DUV0




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-30	SER	-	EXPRESSION TAG	UNP Q8DUV0
D	-29	HIS	-	EXPRESSION TAG	UNP Q8DUV0
D	-28	HIS	-	EXPRESSION TAG	UNP Q8DUV0
D	-27	HIS	-	EXPRESSION TAG	UNP Q8DUV0
D	-26	HIS	-	EXPRESSION TAG	UNP Q8DUV0
D	-25	HIS	-	EXPRESSION TAG	UNP Q8DUV0
D	-24	HIS	-	EXPRESSION TAG	UNP Q8DUV0
D	-23	SER	-	EXPRESSION TAG	UNP Q8DUV0
D	-22	SER	-	EXPRESSION TAG	UNP Q8DUV0
D	-21	GLY	-	EXPRESSION TAG	UNP Q8DUV0
D	-20	LEU	-	EXPRESSION TAG	UNP Q8DUV0
D	-19	VAL	-	EXPRESSION TAG	UNP Q8DUV0
D	-18	PRO	-	EXPRESSION TAG	UNP Q8DUV0
D	-17	ARG	-	EXPRESSION TAG	UNP Q8DUV0
D	-16	GLY	-	EXPRESSION TAG	UNP Q8DUV0
D	-15	SER	-	EXPRESSION TAG	UNP Q8DUV0
D	-14	HIS	-	EXPRESSION TAG	UNP Q8DUV0
D	-13	MET	-	EXPRESSION TAG	UNP Q8DUV0
D	-12	ALA	-	EXPRESSION TAG	UNP Q8DUV0
D	-11	SER	-	EXPRESSION TAG	UNP Q8DUV0
D	-10	MET	-	EXPRESSION TAG	UNP Q8DUV0
D	-9	THR	-	EXPRESSION TAG	UNP Q8DUV0
D	-8	GLY	-	EXPRESSION TAG	UNP Q8DUV0
D	-7	GLY	-	EXPRESSION TAG	UNP Q8DUV0
D	-6	GLN	-	EXPRESSION TAG	UNP Q8DUV0
D	-5	GLN	-	EXPRESSION TAG	UNP Q8DUV0
D	-4	MET	-	EXPRESSION TAG	UNP Q8DUV0
D	-3	GLY	-	EXPRESSION TAG	UNP Q8DUV0
D	-2	ARG	-	EXPRESSION TAG	UNP Q8DUV0
D	-1	GLY	-	EXPRESSION TAG	UNP Q8DUV0
D	0	SER	-	EXPRESSION TAG	UNP Q8DUV0

 $\bullet$  Molecule 2 is COENZYME A (three-letter code: COA) (formula:  $\mathrm{C_{21}H_{36}N_7O_{16}P_3S}).$ 





Mol	Chain	Residues		A	ton	ıs			ZeroOcc	AltConf
2	A	1	Total	С	N	О	Р	S	0	0
	Λ	1	48	21	7	16	3	1	U	
2	В	1	Total	С	Ν	О	Р	S	0	0
	Ъ	1	48	21	7	16	3	1	0	U
2	C	1	Total	С	N	О	Р	S	0	0
		1	48	21	7	16	3	1	0	U
2	D	1	Total	С	N	О	Р	S	0	0
	ש	1	48	21	7	16	3	1	0	U

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	В	2	Total Cl 2 2	0	0
3	D	1	Total Cl 1 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	75	Total O 75 75	0	0
4	В	111	Total O 111 111	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	71	Total O 71 71	0	0
4	D	110	Total O 110 110	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative uncharacterized protein smu.793









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	68.54Å 68.54Å 118.82Å	Danagitan
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	34.29 - 1.70	Depositor
Resolution (A)	34.29 - 1.70	EDS
% Data completeness	96.5 (34.29-1.70)	Depositor
(in resolution range)	98.9 (34.29-1.70)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.27 (at 1.70Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
D.D.	0.129 , 0.188	Depositor
$R, R_{free}$	0.142 , $0.143$	DCC
$R_{free}$ test set	3108 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.2	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36, 38.8	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	0.479 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	4472	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths Bo		Bo	nd angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.39	0/979	0.61	2/1322~(0.2%)
1	В	0.47	0/1035	0.59	0/1396
1	С	0.37	0/979	0.50	0/1322
1	D	0.42	0/1004	0.54	0/1356
All	All	0.42	0/3997	0.56	$2/5396 \ (0.0\%)$

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
1	A	7	HIS	N-CA-C	6.03	127.28	111.00
1	A	6	LEU	CA-CB-CG	-5.75	102.08	115.30

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	963	0	958	7	0
1	В	1004	0	1013	13	0
1	С	963	0	958	6	0
1	D	979	0	982	14	0
2	A	48	0	32	0	0



$\alpha \cdots$	, r	•	
Continued	trom	mromonie	maaa
-	110116	DICULUUS	Duuc
	J	1	1

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
2	В	48	0	32	0	0
2	С	48	0	32	1	0
2	D	48	0	32	0	0
3	A	1	0	0	2	0
3	В	2	0	0	1	0
3	D	1	0	0	0	0
4	A	75	0	0	1	0
4	В	111	0	0	2	0
4	С	71	0	0	2	0
4	D	110	0	0	3	0
All	All	4472	0	4039	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 39 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:13:GLU:H	1:C:53:GLN:HE22	1.25	0.85
1:B:13:GLU:H	1:B:53:GLN:HE22	1.26	0.80
1:D:13:GLU:H	1:D:53:GLN:HE22	1.27	0.80
1:A:13:GLU:H	1:A:53:GLN:HE22	1.30	0.79
1:B:115:LYS:HE2	1:D:102:ASP:OD1	1.87	0.75

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
1	A	122/163 (75%)	120 (98%)	2 (2%)	0	100	100
1	В	129/163 (79%)	127 (98%)	2 (2%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	С	122/163~(75%)	120 (98%)	2 (2%)	0	100	100
1	D	125/163~(77%)	124 (99%)	1 (1%)	0	100	100
All	All	498/652 (76%)	491 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentil	es.
1	A	109/140 (78%)	105 (96%)	4 (4%)	34 15	
1	В	116/140 (83%)	112 (97%)	4 (3%)	37 18	
1	С	109/140 (78%)	106 (97%)	3 (3%)	43 25	
1	D	112/140 (80%)	109 (97%)	3 (3%)	44 26	
All	All	446/560 (80%)	432 (97%)	14 (3%)	39 21	

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	115	LYS
1	С	30	GLU
1	D	108	GLN
1	D	4	ASN
1	D	70	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	С	53	GLN
1	С	70	GLN
1	D	70	GLN
1	D	24	HIS
1	D	53	GLN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chai		Chain	Chain Res Link		Во	ond leng	$ ag{ths}$	Bond angles		
MIOI	$oxed{egin{array}{c c} Mol & Type & Chain \\ \end{array}}$	nes L	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
2	COA	D	130	-	41,50,50	1.64	3 (7%)	52,75,75	1.01	1 (1%)
2	COA	С	130	-	41,50,50	1.64	3 (7%)	52,75,75	1.08	1 (1%)
2	COA	A	130	-	41,50,50	1.65	3 (7%)	52,75,75	1.06	1 (1%)
2	COA	В	130	-	41,50,50	1.63	3 (7%)	52,75,75	1.05	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	D	130	-	-	5/44/64/64	0/3/3/3
2	COA	С	130	-	-	4/44/64/64	0/3/3/3
2	COA	A	130	-	-	5/44/64/64	0/3/3/3
2	COA	В	130	-	-	6/44/64/64	0/3/3/3



The worst 5 of	of 12 b	bond	length	outliers	are	listed	below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(A)
2	A	130	COA	O9P-C9P	8.50	1.40	1.23
2	С	130	COA	O9P-C9P	8.44	1.40	1.23
2	D	130	COA	O9P-C9P	8.40	1.40	1.23
2	В	130	COA	O9P-C9P	8.32	1.39	1.23
2	A	130	COA	C2A-N3A	4.21	1.38	1.32

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	С	130	COA	N3A-C2A-N1A	-5.64	119.86	128.68
2	A	130	COA	N3A-C2A-N1A	-5.59	119.93	128.68
2	D	130	COA	N3A-C2A-N1A	-5.41	120.22	128.68
2	В	130	COA	N3A-C2A-N1A	-5.29	120.40	128.68
2	В	130	COA	O3B-C3B-C2B	-2.31	103.31	111.68

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	130	COA	C3B-O3B-P3B-O7A
2	A	130	COA	CCP-O6A-P2A-O4A
2	A	130	COA	CCP-O6A-P2A-O5A
2	В	130	COA	P2A-O3A-P1A-O5B
2	В	130	COA	CCP-O6A-P2A-O4A

There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	130	COA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	122/163 (74%)	-0.45	2 (1%) 72 76	19, 32, 52, 70	0
1	В	124/163~(76%)	-0.53	2 (1%) 72 76	18, 25, 43, 78	0
1	С	122/163 (74%)	-0.49	2 (1%) 72 76	19, 32, 50, 62	0
1	D	123/163 (75%)	-0.55	1 (0%) 86 88	18, 25, 42, 78	0
All	All	491/652 (75%)	-0.51	7 (1%) 75 79	18, 28, 50, 78	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	6	LEU	4.6
1	A	6	LEU	4.3
1	D	6	LEU	3.6
1	В	6	LEU	3.2
1	A	108	GLN	2.9

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.





Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	COA	В	130	48/48	0.96	0.09	20,36,72,74	0
2	COA	С	130	48/48	0.96	0.10	21,40,77,80	0
2	COA	A	130	48/48	0.97	0.09	22,42,78,80	0
2	COA	D	130	48/48	0.97	0.09	19,39,73,75	0
3	CL	D	131	1/1	0.97	0.09	75,75,75,75	0
3	CL	В	132	1/1	0.98	0.06	78,78,78,78	0
3	CL	A	131	1/1	0.98	0.04	58,58,58,58	0
3	CL	В	131	1/1	1.00	0.04	48,48,48,48	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















## 6.5 Other polymers (i)

There are no such residues in this entry.

