



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2023 – 08:26 AM EDT

PDB ID : 3LEH  
Title : The Crystal Structure of smu.943c from Streptococcus mutans UA159  
Authors : Su, X.-D.; Liu, Y.H.; Liu, X.  
Deposited on : 2010-01-14  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	425	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2812 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative hydroxymethylglutaryl-CoA synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	356	2734	1760	451	515	8	0	0	0

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	expression tag	UNP Q8DUI5
A	-32	GLY	-	expression tag	UNP Q8DUI5
A	-31	SER	-	expression tag	UNP Q8DUI5
A	-30	SER	-	expression tag	UNP Q8DUI5
A	-29	HIS	-	expression tag	UNP Q8DUI5
A	-28	HIS	-	expression tag	UNP Q8DUI5
A	-27	HIS	-	expression tag	UNP Q8DUI5
A	-26	HIS	-	expression tag	UNP Q8DUI5
A	-25	HIS	-	expression tag	UNP Q8DUI5
A	-24	HIS	-	expression tag	UNP Q8DUI5
A	-23	SER	-	expression tag	UNP Q8DUI5
A	-22	SER	-	expression tag	UNP Q8DUI5
A	-21	GLY	-	expression tag	UNP Q8DUI5
A	-20	LEU	-	expression tag	UNP Q8DUI5
A	-19	VAL	-	expression tag	UNP Q8DUI5
A	-18	PRO	-	expression tag	UNP Q8DUI5
A	-17	ARG	-	expression tag	UNP Q8DUI5
A	-16	GLY	-	expression tag	UNP Q8DUI5
A	-15	SER	-	expression tag	UNP Q8DUI5
A	-14	HIS	-	expression tag	UNP Q8DUI5
A	-13	MET	-	expression tag	UNP Q8DUI5
A	-12	ALA	-	expression tag	UNP Q8DUI5
A	-11	SER	-	expression tag	UNP Q8DUI5
A	-10	MET	-	expression tag	UNP Q8DUI5
A	-9	THR	-	expression tag	UNP Q8DUI5
A	-8	GLY	-	expression tag	UNP Q8DUI5
A	-7	GLY	-	expression tag	UNP Q8DUI5

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLN	-	expression tag	UNP Q8DUI5
A	-5	GLN	-	expression tag	UNP Q8DUI5
A	-4	MET	-	expression tag	UNP Q8DUI5
A	-3	GLY	-	expression tag	UNP Q8DUI5
A	-2	ARG	-	expression tag	UNP Q8DUI5
A	-1	GLY	-	expression tag	UNP Q8DUI5
A	0	SER	-	expression tag	UNP Q8DUI5

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total N O 4 1 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	74	Total O 74 74	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative hydroxymethylglutaryl-CoA synthase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	131.64Å 56.08Å 51.55Å 90.00° 90.32° 90.00°	Depositor
Resolution (Å)	40.48 – 1.70 40.48 – 1.70	Depositor EDS
% Data completeness (in resolution range)	86.7 (40.48-1.70) 86.7 (40.48-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.59 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.5.0066	Depositor
R, $R_{free}$	0.217 , 0.254 0.215 , 0.256	Depositor DCC
$R_{free}$ test set	1815 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.0	Xtrriage
Anisotropy	0.945	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2812	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.10	1/2775 (0.0%)	1.05	5/3736 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	16	VAL	CB-CG1	5.75	1.65	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	136	LEU	CA-CB-CG	-5.86	101.83	115.30
1	A	39	LEU	CA-CB-CG	5.74	128.49	115.30
1	A	72	ASP	CB-CG-OD1	5.53	123.28	118.30
1	A	234	LEU	CA-CB-CG	5.33	127.56	115.30
1	A	181	ASP	CB-CG-OD1	5.30	123.07	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	379	LEU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2734	0	2744	76	0
2	A	4	0	0	0	0
3	A	74	0	0	4	0
All	All	2812	0	2744	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:ILE:CG2	1:A:204:GLN:HE22	1.49	1.26
1:A:246:LYS:O	1:A:247:ILE:CD1	1.88	1.20
1:A:246:LYS:O	1:A:247:ILE:HD12	0.97	1.14
1:A:182:ILE:HG21	1:A:204:GLN:NE2	1.69	1.08
1:A:182:ILE:HG23	1:A:204:GLN:HE22	1.19	1.07
1:A:32:LYS:HG2	1:A:32:LYS:O	1.53	1.04
1:A:224:LEU:HD11	1:A:247:ILE:CG2	1.94	0.97
1:A:224:LEU:HD11	1:A:247:ILE:HG23	1.49	0.95
1:A:182:ILE:CG2	1:A:204:GLN:NE2	2.24	0.93
1:A:36:GLY:O	1:A:238:LYS:HD2	1.68	0.93
1:A:182:ILE:HG21	1:A:204:GLN:CD	1.95	0.87
1:A:182:ILE:HG21	1:A:204:GLN:HE22	1.25	0.87
1:A:224:LEU:CD1	1:A:247:ILE:HG23	2.10	0.81
1:A:47:ILE:HD12	1:A:48:THR:N	2.00	0.75
1:A:19:MET:HE3	1:A:33:PHE:CB	2.17	0.74
1:A:51:VAL:HG11	1:A:80:SER:HB2	1.71	0.72
1:A:381:HIS:CD2	1:A:381:HIS:N	2.56	0.71
1:A:32:LYS:O	1:A:32:LYS:CG	2.37	0.71
1:A:47:ILE:HD12	1:A:48:THR:HG23	1.73	0.71
1:A:182:ILE:HG21	1:A:204:GLN:OE1	1.89	0.71
1:A:19:MET:HE3	1:A:33:PHE:HB3	1.71	0.70
1:A:261:LYS:NZ	1:A:265:GLU:OE2	2.21	0.70
1:A:180:ARG:NH1	1:A:207:ASP:OD2	2.21	0.68
1:A:47:ILE:CD1	1:A:48:THR:HG23	2.25	0.66

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:340:THR:OG1	1:A:369:GLN:NE2	2.29	0.65
1:A:19:MET:HE3	1:A:33:PHE:CG	2.34	0.62
1:A:172:HIS:HE1	1:A:319:THR:OG1	1.84	0.59
1:A:249:ASP:C	1:A:251:GLN:H	2.07	0.58
1:A:224:LEU:HD11	1:A:247:ILE:HG21	1.83	0.57
1:A:381:HIS:N	1:A:381:HIS:HD2	2.02	0.57
1:A:19:MET:CE	1:A:151:SER:HB2	2.36	0.56
1:A:234:LEU:HD21	1:A:268:ILE:HG22	1.89	0.55
1:A:291:GLU:HG2	1:A:328:LEU:HG	1.87	0.55
1:A:176:LEU:HD21	1:A:215:GLU:HB3	1.88	0.53
1:A:325:LYS:HA	1:A:328:LEU:HD22	1.92	0.52
1:A:47:ILE:HD12	1:A:47:ILE:C	2.30	0.52
1:A:253:PRO:O	1:A:257:GLN:NE2	2.43	0.51
1:A:47:ILE:HD13	1:A:365:PHE:CD1	2.45	0.51
1:A:37:LEU:O	1:A:237:PRO:HD2	2.12	0.50
1:A:19:MET:CE	1:A:33:PHE:CG	2.94	0.50
1:A:285:GLY:HA3	3:A:437:HOH:O	2.12	0.50
1:A:388:VAL:HG21	3:A:435:HOH:O	2.13	0.48
1:A:19:MET:CE	1:A:33:PHE:HB3	2.43	0.48
1:A:19:MET:HE1	1:A:151:SER:HB2	1.95	0.47
1:A:176:LEU:CD2	1:A:215:GLU:HB3	2.46	0.46
1:A:246:LYS:O	1:A:247:ILE:CG1	2.62	0.46
1:A:26:ARG:NH1	1:A:145:ILE:HG23	2.31	0.45
1:A:42:LEU:HD12	1:A:44:ILE:HD11	1.98	0.45
1:A:384:ILE:N	1:A:384:ILE:HD12	2.32	0.45
1:A:1:MET:CG	1:A:3:ILE:HD11	2.46	0.45
1:A:301:LYS:HG2	1:A:319:THR:OG1	2.17	0.45
1:A:245:ASN:O	1:A:247:ILE:N	2.50	0.44
1:A:51:VAL:CG1	1:A:80:SER:HB2	2.42	0.44
1:A:298:ALA:HB2	1:A:322:LYS:HA	2.00	0.44
1:A:289:LEU:HD11	1:A:296:LEU:HD11	2.01	0.43
1:A:384:ILE:N	1:A:384:ILE:CD1	2.82	0.43
1:A:286:LEU:O	1:A:290:LEU:HG	2.19	0.43
1:A:19:MET:HE2	1:A:151:SER:HB2	2.00	0.43
1:A:172:HIS:CE1	1:A:319:THR:OG1	2.69	0.43
1:A:19:MET:HE3	1:A:33:PHE:CD2	2.53	0.42
1:A:76:LEU:HD23	1:A:136:LEU:HB2	2.02	0.42
1:A:277:ILE:HG21	1:A:277:ILE:HD13	1.85	0.42
1:A:46:PRO:HA	1:A:350:ILE:HG22	2.02	0.42
1:A:182:ILE:HG23	1:A:204:GLN:NE2	2.05	0.42
1:A:1:MET:CE	1:A:129:PRO:HB3	2.50	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:380:GLU:HA	1:A:381:HIS:HA	1.55	0.41
1:A:123:LEU:HD23	1:A:123:LEU:HA	1.72	0.41
1:A:380:GLU:H	1:A:380:GLU:HG2	1.59	0.41
1:A:53:THR:HG21	1:A:374:ALA:HA	2.03	0.41
1:A:340:THR:HA	1:A:341:PRO:HD3	1.93	0.41
1:A:75:ILE:HG21	1:A:117:ALA:CB	2.51	0.41
1:A:291:GLU:CG	1:A:328:LEU:HG	2.49	0.41
1:A:178:GLN:NE2	3:A:462:HOH:O	2.33	0.40
1:A:15:TYR:OH	1:A:369:GLN:HG3	2.20	0.40
1:A:47:ILE:HG23	3:A:419:HOH:O	2.20	0.40
1:A:253:PRO:O	1:A:257:GLN:CG	2.70	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	339/425 (80%)	324 (96%)	12 (4%)	3 (1%)	<b>17</b>   <b>5</b>

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	246	LYS
1	A	250	LYS
1	A	109	GLU

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	286/358 (80%)	261 (91%)	25 (9%)	10 2

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	LEU
1	A	22	LEU
1	A	26	ARG
1	A	32	LYS
1	A	39	LEU
1	A	42	LEU
1	A	63	LEU
1	A	76	LEU
1	A	85	SER
1	A	94	SER
1	A	164	LYS
1	A	180	ARG
1	A	224	LEU
1	A	230	PHE
1	A	238	LYS
1	A	239	LEU
1	A	247	ILE
1	A	253	PRO
1	A	261	LYS
1	A	322	LYS
1	A	328	LEU
1	A	347	TYR
1	A	356	GLN
1	A	367	GLU
1	A	381	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	172	HIS
1	A	204	GLN
1	A	233	HIS
1	A	257	GLN
1	A	273	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	294	GLN
1	A	369	GLN
1	A	381	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NO3	A	392	-	1,3,3	3.52	1 (100%)	0,3,3	-	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	392	NO3	O1-N	3.52	1.40	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	356/425 (83%)	0.67	43 (12%) <b>4</b> <b>5</b>	15, 26, 51, 72	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	PHE	6.4
1	A	384	ILE	5.7
1	A	379	LEU	5.4
1	A	32	LYS	5.1
1	A	37	LEU	4.2
1	A	148	PRO	3.8
1	A	201	SER	3.7
1	A	202	THR	3.7
1	A	355	ALA	3.7
1	A	27	GLY	3.3
1	A	381	HIS	3.3
1	A	363	ALA	3.3
1	A	378	ILE	3.3
1	A	85	SER	3.2
1	A	143	TYR	3.2
1	A	256	LEU	3.0
1	A	254	SER	3.0
1	A	117	ALA	3.0
1	A	34	SER	3.0
1	A	356	GLN	3.0
1	A	135	VAL	2.9
1	A	38	LEU	2.9
1	A	136	LEU	2.8
1	A	364	SER	2.8
1	A	116	ALA	2.7
1	A	350	ILE	2.6
1	A	222	VAL	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	75	ILE	2.6
1	A	47	ILE	2.6
1	A	25	ALA	2.5
1	A	114	ALA	2.4
1	A	149	GLY	2.4
1	A	147	THR	2.3
1	A	247	ILE	2.3
1	A	255	ASP	2.2
1	A	28	GLU	2.2
1	A	283	PHE	2.2
1	A	377	GLU	2.2
1	A	239	LEU	2.2
1	A	76	LEU	2.1
1	A	380	GLU	2.1
1	A	244	PHE	2.1
1	A	221	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NO3	A	392	4/4	0.84	0.14	34,34,35,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.