



# wwPDB X-ray Structure Validation Summary Report

May 24, 2021 – 08:09 PM EDT

PDB ID : 7LJN  
Title : Structure of the Bradyrhizobium diazoefficiens CD-NTase CdnG in complex with GTP  
Authors : Govande, A.; Lowey, B.; Eaglesham, J.B.; Whiteley, A.T.; Kranzusch, P.J.  
Deposited on : 2021-01-29  
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.18  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.18


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	416	
1	B	416	
1	C	416	
1	D	416	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CD-NTase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	342	2696	1697	486	503	10	0	0	0
1	B	335	2640	1661	479	490	10	0	0	0
1	C	337	2651	1668	479	494	10	0	0	0
1	D	339	2673	1683	483	497	10	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q89Y83
B	1	SER	-	expression tag	UNP Q89Y83
C	1	SER	-	expression tag	UNP Q89Y83
D	1	SER	-	expression tag	UNP Q89Y83

- Molecule 2 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ) (labeled as "Ligand of Interest" by depositor).




*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	B	508	Total 508	O 508	0	0
4	C	512	Total 512	O 512	0	0
4	D	487	Total 487	O 487	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CD-NTase



- Molecule 1: CD-NTase



- Molecule 1: CD-NTase



ALA  
ALA  
PRO  
ALA  
ILE  
ILE  
GLY  
ALA  
ALA  
THR  
SER  
SER  
LEU  
ALA  
PRO  
VAL  
ALA  
ALA  
ARG  
ALA  
ALA  
HIS  
THR  
ASN  
MET  
GLY  
GLU  
ARG  
ARG

● Molecule 1: CD-NTase



SER  
SER  
ASN  
PHE  
PRO  
SER  
LEU  
ARG  
ARG  
ASP  
ASP  
ARG  
PRO  
ASP  
ASP  
P16  
L21  
E27  
L28  
E44  
E54  
R63  
T81  
Y88  
D89  
L107  
D108  
Y117  
P118  
R124  
D140  
E151  
K152  
A159  
LYS  
GLY  
THR  
ARG  
SER  
ASP  
P167  
M173  
Y176

K180  
R185  
V213  
L222  
V227  
R265  
P266  
G267  
M268  
E272  
R283  
E301  
R306  
R323  
R336  
THR  
GLY  
ASP  
VAL  
GLN  
LEU  
E343  
F353  
R356  
S361  
W362  
K363  
A364  
F365  
R366  
Q367  
ARG  
LEU  
GLY  
ARG  
GLN  
VAL  
GLN  
SER  
ARG  
GLN  
HIS  
GLY  
TYR  
THR  
ARG

SER  
GLY  
GLY  
LEU  
PHE  
VAL  
PRO  
ALA  
ALA  
PRO  
ALA  
ILE  
ILE  
GLY  
ALA  
ALA  
THR  
SER  
LEU  
ALA  
PRO  
VAL  
ALA  
ALA  
ARG  
ALA  
HIS  
THR  
MET  
MET  
GLY  
GLU  
ARG  
ARG

## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.00Å 68.69Å 112.43Å 90.00° 108.41° 90.00°	Depositor
Resolution (Å)	37.94 – 1.60 37.94 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.9 (37.94-1.60) 98.2 (37.94-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.99 (at 1.60Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.197 , 0.218 0.197 , 0.218	Depositor DCC
$R_{free}$ test set	2018 reflections (0.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.4	Xtrriage
Anisotropy	0.104	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 30.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.366 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	12830	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 52.78 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.6345e-05.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/2757	0.51	0/3750
1	B	0.35	0/2699	0.51	0/3670
1	C	0.34	0/2711	0.52	0/3689
1	D	0.33	0/2734	0.52	0/3717
All	All	0.34	0/10901	0.51	0/14826

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2696	0	2666	30	0
1	B	2640	0	2620	54	0
1	C	2651	0	2625	27	0
1	D	2673	0	2645	34	0
2	A	32	0	12	0	0
2	B	32	0	12	0	0
2	C	32	0	12	0	0
2	D	32	0	12	0	0
3	A	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	531	0	0	16	3
4	B	508	0	0	22	6
4	C	512	0	0	13	3
4	D	487	0	0	21	1
All	All	12830	0	10604	145	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 145 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:367:GLN:O	4:B:601:HOH:O	1.82	0.97
1:D:124:ARG:NH1	4:D:604:HOH:O	2.01	0.92
1:A:336:ARG:NH1	4:A:601:HOH:O	2.01	0.91
1:C:265:ARG:NH2	4:C:603:HOH:O	2.04	0.91
1:D:301:GLU:OE2	4:D:601:HOH:O	1.88	0.91

The worst 5 of 7 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1069:HOH:O	4:B:969:HOH:O[1_545]	1.94	0.26
4:B:1016:HOH:O	4:C:993:HOH:O[2_454]	2.00	0.20
4:A:1126:HOH:O	4:D:1014:HOH:O[2_545]	2.13	0.07
4:B:861:HOH:O	4:C:919:HOH:O[2_454]	2.13	0.07
4:B:1055:HOH:O	4:B:1085:HOH:O[2_454]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	336/416 (81%)	333 (99%)	3 (1%)	0	100	100
1	B	329/416 (79%)	324 (98%)	5 (2%)	0	100	100
1	C	331/416 (80%)	330 (100%)	1 (0%)	0	100	100
1	D	333/416 (80%)	330 (99%)	3 (1%)	0	100	100
All	All	1329/1664 (80%)	1317 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/338 (83%)	279 (99%)	2 (1%)	84	73
1	B	275/338 (81%)	274 (100%)	1 (0%)	91	84
1	C	276/338 (82%)	274 (99%)	2 (1%)	84	73
1	D	278/338 (82%)	277 (100%)	1 (0%)	91	84
All	All	1110/1352 (82%)	1104 (100%)	6 (0%)	88	80

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	51	ARG
1	C	173	MET
1	D	173	MET
1	A	173	MET
1	A	67	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GTP	B	501	3	26,34,34	0.92	1 (3%)	33,54,54	1.49	3 (9%)
2	GTP	A	501	3	26,34,34	1.06	2 (7%)	33,54,54	1.57	4 (12%)
2	GTP	D	501	3	26,34,34	0.90	1 (3%)	33,54,54	1.49	5 (15%)
2	GTP	C	501	3	26,34,34	0.86	1 (3%)	33,54,54	1.63	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	B	501	3	-	1/18/38/38	0/3/3/3
2	GTP	A	501	3	-	2/18/38/38	0/3/3/3
2	GTP	D	501	3	-	2/18/38/38	0/3/3/3
2	GTP	C	501	3	-	2/18/38/38	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	GTP	C6-N1	3.30	1.38	1.33
2	B	501	GTP	C6-N1	2.78	1.37	1.33
2	D	501	GTP	C6-N1	2.64	1.37	1.33
2	C	501	GTP	C6-N1	2.57	1.37	1.33
2	A	501	GTP	C2-N1	2.13	1.39	1.35

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	GTP	N3-C2-N1	-5.18	120.31	127.22
2	C	501	GTP	N3-C2-N1	-4.98	120.58	127.22
2	B	501	GTP	N3-C2-N1	-4.58	121.11	127.22
2	D	501	GTP	N3-C2-N1	-4.47	121.26	127.22
2	D	501	GTP	C5-C6-N1	-4.13	117.78	123.43

There are no chirality outliers.


5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	GTP	PG-O3B-PB-O2B
2	B	501	GTP	PG-O3B-PB-O1B
2	C	501	GTP	PG-O3B-PB-O2B
2	D	501	GTP	PG-O3B-PB-O2B
2	A	501	GTP	PG-O3B-PB-O1B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	342/416 (82%)	-0.47	3 (0%) 84 84	11, 19, 41, 63	0
1	B	335/416 (80%)	-0.49	4 (1%) 79 78	11, 20, 40, 58	0
1	C	337/416 (81%)	-0.51	5 (1%) 73 73	12, 20, 38, 50	0
1	D	339/416 (81%)	-0.50	6 (1%) 68 67	11, 20, 42, 65	0
All	All	1353/1664 (81%)	-0.49	18 (1%) 77 77	11, 20, 40, 65	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	365	PHE	10.0
1	A	342	LEU	4.9
1	D	267	GLY	4.4
1	B	367	GLN	3.6
1	D	363	LYS	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GTP	A	501	32/32	0.99	0.06	10,13,15,16	0
2	GTP	B	501	32/32	0.99	0.05	9,12,15,16	0
2	GTP	C	501	32/32	0.99	0.06	10,13,16,18	0
2	GTP	D	501	32/32	0.99	0.06	9,13,17,23	0
3	MG	B	502	1/1	0.99	0.04	12,12,12,12	0
3	MG	C	502	1/1	0.99	0.04	11,11,11,11	0
3	MG	D	502	1/1	0.99	0.04	12,12,12,12	0
3	MG	A	502	1/1	1.00	0.06	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.