

Full wwPDB X-ray Structure Validation Report (i)

Aug 7, 2020 – 05:13 PM BST

PDB ID	:	5LKR
Title	:	Human Butyrylcholinesterase complexed with N-Propargyliperidines
Authors	:	Coquelle, N.; Knez, D.; Colletier, J.P.; Gobec, S.
Deposited on	:	2016-07-23
Resolution	:	2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	÷	4.02b-467
Mogul		1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	$5743\ (2.54-2.50)$
Clashscore	141614	$6463 \ (2.54-2.50)$
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630(2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	574	4% 81%	10% • 8%
1	В	574	5%	11% • 9%
		011	1970	1170 • 970
2	С	2	100%	
2	D	2	50%	50%
2	F	2	100%	
2	G	2	100%	



Mol	Chain	Length	Quality	of chain
2	Н	2	10	00%
2	Ι	2	50%	50%
3	Е	2	50%	50%
4	J	2	10	00%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	FUL	А	604	-	-	-	Х
9	PEG	A	618	_	Х	-	-



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 8751 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Cholinesterase.

Mol	Chain	Residues		Atoms					AltConf	Trace
1	А	528	Total 4168	C 2694	N 695	O 763	S 16	0	3	1
1	В	525	Total 4130	C 2668	N 697	O 750	S 15	0	0	1

• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	С	2	Total C N O 28 16 2 10	0	0	0
2	D	2	Total C N O 28 16 2 10	0	0	0
2	F	2	Total C N O 28 16 2 10	0	0	0
2	G	2	Total C N O 28 16 2 10	0	0	0
2	Н	2	Total C N O 28 16 2 10	0	0	0
2	Ι	2	Total C N O 28 16 2 10	0	0	0

• Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-bet a-D-glucopyranose.





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	Е	2	Total 24	C 14	N 1	O 9	0	0	0

• Molecule 4 is an oligosaccharide called beta-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta -D-glucopyranose.



Mol	Chain	Residues	A	Aton	ıs		ZeroOcc	AltConf	Trace
4	J	2	Total 24	C 14	N 1	O 9	0	0	0

• Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf	
5	Λ	1	Total	С	Ν	Ο	0	0	
J	Л	, <u>1</u>	14	8	1	5	0	0	
5	В	1	Total	С	Ν	Ο	0	0	
	D	T	14	8	1	5	0	0	
5	В	1	Total	С	Ν	Ο	0	0	
	D	T	14	8	1	5	0	0	
5	В	1	Total	С	Ν	0	0	0	
	J D	L	14	8	1	5	0	U	



001100	.easa jisn		30			
Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
ц	D	1	Total C N O	0	0	
D B	L	14 8 1 5	0	0		
ĸ	D	D	1	Total C N O	0	0
5 В	D	1	14 8 1 5	0	0	
ĸ	р	1	Total C N O	0	0	
9 B	1	14 8 1 5	0	0		
5	р	1	Total C N O	0	0	
	D	В	14 8 1 5	0	U	

Continued from previous page...

• Molecule 6 is beta-L-fucopyranose (three-letter code: FUL) (formula: $C_6H_{12}O_5$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	А	1	$\begin{array}{cc} {\rm Total} & {\rm C} \\ 10 & 6 \end{array}$	O 4	0	0

• Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
7	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
7	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 8 is {N}-(2-methoxyethyl)- {N}-[[(3 {S})-1-prop-2-ynylpiperidin-3-yl]methyl]nap hthalene-2-carboxamide (three-letter code: 6YC) (formula: $C_{23}H_{28}N_2O_2$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	А	1	Total 27	С 23	N 2	O 2	9	0
8	В	1	Total 27	C 23	N 2	O 2	0	0

• Molecule 9 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
9	А	1	Total 7	$\begin{array}{c} \mathrm{C} \\ 4 \end{array}$	O 3	0	0

• Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	А	15	Total O 15 15	0	0
10	В	21	TotalO2121	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Cholinesterase

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:

100%



NAG1 NAG2

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:	50%	50%	
NAG1 NAG2			
• Molecule opyranose	2: 2-acetamido-2-deoxy-beta-D	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain F:	100	0%	
NAG1 NAG2			
• Molecule opyranose	2: 2-acetamido-2-deoxy-beta-D	9-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain G:	10	0%	
NAG1 NAG2			
• Molecule opyranose	2: 2-acetamido-2-deoxy-beta-D	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain H:	10	0%	
NAG1 NAG2			
• Molecule opyranose	2: 2-acetamido-2-deoxy-beta-D	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain I:	50%	50%	
NAG1 NAG2			
• Molecule	3: alpha-L-fucopyranose-(1-6)-	2-acetamido-2-deoxy-beta-D-gluc	opyranose
Chain E:	50%	50%	
NAG1 FUC2			

• Molecule 4: beta-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



Chain J:

100%

NAG 1 FUL2



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	73.68Å 79.46Å 228.87Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	48.86 - 2.52	Depositor
Resolution (A)	48.86 - 2.61	EDS
% Data completeness	99.1 (48.86-2.52)	Depositor
(in resolution range)	94.7 (48.86 - 2.61)	EDS
R_{merge}	0.06	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.46 (at 2.61 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
D D.	0.185 , 0.238	Depositor
Π, Π_{free}	0.199 , 0.202	DCC
R_{free} test set	1985 reflections (5.00%)	wwPDB-VP
Wilson B-factor $(Å^2)$	75.0	Xtriage
Anisotropy	0.314	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33 , 57.8	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8751	wwPDB-VP
Average B, all atoms $(Å^2)$	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, 6YC, FUC, FUL, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bond angles		
10101	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.35	0/4298	0.52	0/5849	
1	В	0.36	2/4247~(0.0%)	0.53	3/5773~(0.1%)	
All	All	0.35	2/8545~(0.0%)	0.52	3/11622~(0.0%)	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
1	В	364	PHE	CB-CG	-8.11	1.37	1.51
1	В	364	PHE	CD1-CE1	-6.00	1.27	1.39

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	В	364	PHE	N-CA-CB	-5.26	101.12	110.60
1	В	364	PHE	CB-CG-CD1	-5.16	117.19	120.80
1	В	95	LEU	CA-CB-CG	5.02	126.86	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4168	0	4015	35	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	4130	0	3994	37	0
2	С	28	0	25	0	0
2	D	28	0	25	0	0
2	F	28	0	25	1	0
2	G	28	0	25	0	0
2	Н	28	0	25	0	0
2	Ι	28	0	25	0	0
3	Е	24	0	22	0	0
4	J	24	0	22	0	0
5	А	14	0	13	1	0
5	В	98	0	91	1	0
6	А	10	0	10	0	0
7	А	12	0	16	3	0
7	В	6	0	8	0	0
8	А	27	0	0	0	0
8	В	27	0	0	0	0
9	А	7	0	10	1	0
10	A	15	0	0	1	0
10	В	21	0	0	2	0
All	All	8751	0	8351	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:323:LYS:HD2	1:A:422:GLU:HG2	1.67	0.75
1:B:240:ARG:NH1	10:B:703:HOH:O	2.17	0.74
5:B:607:NAG:O4	10:B:701:HOH:O	2.09	0.70
1:B:271:GLU:O	1:B:275:ASN:ND2	2.26	0.67
1:B:69:ILE:HD11	1:B:88:LEU:HD11	1.75	0.67
1:A:424:ARG:NH2	1:A:430:TRP:O	2.27	0.66
1:B:99:ILE:HD11	1:B:185:PHE:HB3	1.78	0.66
1:B:335:PRO:HD3	1:B:356:ILE:HD12	1.78	0.65
1:A:198:SER:OG	10:A:701:HOH:O	2.14	0.65
1:A:523:THR:HG22	7:A:615:GOL:H32	1.82	0.61
7:A:617:GOL:O3	9:A:618:PEG:O4	2.19	0.58
1:B:102:PRO:O	1:B:138:ARG:NH2	2.26	0.58
1:B:108:THR:HG21	1:B:476:LYS:HA	1.86	0.58
1:B:256:ASN:OD1	1:B:256:ASN:N	2.38	0.56



		Interatomic	Clash	
Atom-1	Atom-2	distance $(Å)$	overlap (Å)	
1:B:234:THR:HG22	1:B:238:GLU:HB2	1.87	0.56	
1:B:109:VAL:HB	1:B:192:VAL:HG22	1.88	0.56	
1:B:363:GLU:HA	1:B:366:LYS:HE3	1.88	0.55	
1:B:500:TYR:CZ	1:B:511:MET:HB2	2.42	0.55	
1:B:110:LEU:HD23	1:B:193:THR:HB	1.90	0.54	
1:B:195:PHE:CB	1:B:221:ILE:HB	2.39	0.53	
1:A:336:GLY:H	2:F:2:NAG:H61	1.73	0.53	
1:B:462:ILE:O	1:B:466:SER:OG	2.27	0.51	
1:A:419:TYR:HB3	1:A:490:TRP:CZ2	2.47	0.50	
1:A:500:TYR:CZ	1:A:511:MET:HB2	2.46	0.50	
1:A:99:ILE:HD11	1:A:185:PHE:HD2	1.77	0.50	
1:B:34:ALA:HB2	1:B:173:LEU:HD23	1.93	0.50	
1:B:239:ALA:HB2	1:B:293:THR:HG21	1.94	0.50	
1:A:341:ASN:OD1	1:A:341:ASN:N	2.44	0.49	
1:B:379:ASP:OD1	1:B:379:ASP:N	2.30	0.49	
1:A:154:LEU:HD11	1:A:243:THR:HG23	1.95	0.49	
1:A:328:ALA:HB2	1:A:437:MET:HE3	1.95	0.48	
1:B:450:LEU:HA	1:B:461:GLU:HG3	1.93	0.48	
1:B:195:PHE:HB2	1:B:221:ILE:HB	1.95	0.48	
1:A:523:THR:HA	7:A:615:GOL:H12	1.96	0.48	
1:A:526:PHE:O	1:A:529:VAL:HG12	2.13	0.47	
1:B:328:ALA:HB2	1:B:437:MET:HE3	1.97	0.47	
1:A:17:ASN:HD22	5:A:601:NAG:H83	1.79	0.47	
1:B:236:LEU:HD11	1:B:294:VAL:O	2.14	0.47	
1:A:368:SER:HG	1:A:525:PHE:HE2	1.60	0.47	
1:A:218:THR:HG22	1:A:219:ARG:HG3	1.98	0.46	
1:A:168:LEU:HD11	1:A:292:PRO:HB3	1.97	0.46	
1:A:450:LEU:HA	1:A:461:GLU:HG3	1.98	0.46	
1:B:320:GLY:HA3	1:B:419:TYR:CE2	2.51	0.46	
1:B:224:SER:OG	1:B:325:GLU:OE2	2.25	0.46	
1:A:39:GLY:O	1:A:265:ARG:HD3	2.15	0.46	
1:B:219:ARG:HD3	1:B:316:GLN:OE1	2.16	0.46	
1:B:299:LEU:HD23	1:B:299:LEU:HA	1.82	0.45	
1:B:321:VAL:HG21	1:B:399:ILE:HG12	1.98	0.45	
1:A:16:MET:HB3	1:A:59:THR:HG22	1.99	0.45	
$1:A:424:AR\overline{G:HG3}$	1:A:428:LEU:HD23	1.99	0.45	
1:B:155:ALA:HB3	1:B:239:ALA:HB1	1.99	0.45	
1:A:369:ILE:HD11	1:A:526:PHE:CE1	2.52	0.44	
1:A:69:ILE:HD11	1:A:88:LEU:HD11	1.99	0.44	
1:B:362:SER:C	1:B:364:PHE:H	2.21	0.44	
1:A:159:ASN:OD1	1:A:161:GLU:HB2	2.17	0.44	



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:310:GLY:HA2	1:A:312:PHE:CE2	2.53	0.44
1:A:319:VAL:O	1:A:418:PHE:HA	2.18	0.43
1:A:500:TYR:CE1	1:A:514:LEU:HB2	2.53	0.43
1:B:362:SER:C	1:B:364:PHE:N	2.72	0.43
1:A:208:LEU:HD23	1:A:217:PHE:HZ	1.84	0.43
1:A:195:PHE:CB	1:A:221:ILE:HB	2.49	0.43
1:B:234:THR:HB	1:B:293:THR:HG22	2.01	0.42
1:A:427:LYS:HD3	1:A:456:TYR:CZ	2.53	0.42
1:B:489:SER:O	1:B:510:ILE:HD11	2.19	0.42
1:A:66[B]:CYS:SG	1:A:273:LEU:HD11	2.60	0.42
1:B:66:CYS:HB3	1:B:273:LEU:HD11	2.02	0.41
1:B:96:ASN:O	1:B:142:VAL:HA	2.21	0.41
1:B:197:GLU:HA	1:B:223:GLN:O	2.20	0.41
1:A:41:LEU:HD23	1:A:41:LEU:HA	1.85	0.41
1:B:427:LYS:HD3	1:B:456:TYR:CZ	2.56	0.40
1:A:379:ASP:O	1:A:380:GLN:HG2	2.21	0.40
1:A:341:ASN:HB2	1:A:343:SER:H	1.86	0.40
1:A:99:ILE:HD11	1:A:185:PHE:CD2	2.55	0.40
1:B:197:GLU:HG3	1:B:441:GLU:OE2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	529/574~(92%)	503~(95%)	23 (4%)	3 (1%)	25	41
1	В	521/574~(91%)	488 (94%)	32 (6%)	1 (0%)	47	67
All	All	1050/1148~(92%)	991 (94%)	55~(5%)	4 (0%)	34	53

All (4) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	В	506	GLU
1	А	282	TYR
1	А	54	ASP
1	А	281	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	444/494~(90%)	435~(98%)	9 (2%)	55 78		
1	В	436/494~(88%)	417~(96%)	19 (4%)	28 49	1	
All	All	880/988~(89%)	852~(97%)	28~(3%)	39 63		

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	48	SER
1	А	54	ASP
1	А	195	PHE
1	А	341	ASN
1	А	368	SER
1	А	383	GLU
1	А	425	SER
1	А	428	LEU
1	А	471	TRP
1	В	72	SER
1	В	79	SER
1	В	105	LYS
1	В	144	MET
1	В	156	LEU
1	В	176	GLN
1	В	195	PHE
1	В	234	THR
1	В	240	ARG
1	В	256	ASN
1	В	286	LEU



continuea front problout page								
\mathbf{Mol}	Chain	\mathbf{Res}	Type					
1	В	344	ILE					
1	В	364	PHE					
1	В	379	ASP					
1	В	408	LYS					
1	В	428	LEU					
1	В	466	SER					
1	В	506	GLU					
1	В	511	MET					

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Tune		Chain	Chain	Dog	Tink	Bo	ond leng	\mathbf{ths}	B	ond ang	les		
		Unaim		Ullalli	Unain	Ullalli	nes	Res LINK	Counts	RMSZ	# Z > 2	Counts	RMSZ
2	NAG	C	1	1,2	14,14,15	0.20	0	17,19,21	0.49	0			
2	NAG	C	2	2	14, 14, 15	0.30	0	17,19,21	0.49	0			
2	NAG	D	1	1,2	14,14,15	0.41	0	17,19,21	0.82	1 (5%)			
2	NAG	D	2	2	14,14,15	0.70	0	17,19,21	0.48	0			
3	NAG	E	1	1,3	14, 14, 15	0.39	0	17,19,21	0.45	0			
3	FUC	E	2	3	10, 10, 11	1.07	0	14,14,16	1.22	2 (14%)			
2	NAG	F	1	1,2	14,14,15	0.49	0	17,19,21	1.02	2 (11%)			



Mal	Tune	Chain	Dec	Tink	Bo	ond leng	$_{\rm ths}$	B	ond ang	les
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	F	2	2	$14,\!14,\!15$	0.37	0	17,19,21	0.45	0
2	NAG	G	1	1,2	14, 14, 15	0.35	0	17,19,21	0.79	1(5%)
2	NAG	G	2	2	14,14,15	0.56	0	17,19,21	0.79	1 (5%)
2	NAG	Н	1	1,2	14,14,15	0.59	0	17,19,21	0.67	0
2	NAG	Н	2	2	14, 14, 15	0.33	0	17,19,21	0.55	0
2	NAG	Ι	1	1,2	14,14,15	0.77	1(7%)	17,19,21	0.83	1(5%)
2	NAG	Ι	2	2	14, 14, 15	0.21	0	17,19,21	0.57	0
4	NAG	J	1	1,4	14, 14, 15	0.60	0	$17,\!19,\!21$	0.80	1 (5%)
4	FUL	J	2	4	10, 10, 11	1.99	3 (30%)	14,14,16	2.12	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	С	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	С	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
3	NAG	Е	1	1,3	-	0/6/23/26	0/1/1/1
3	FUC	Е	2	3	-	-	0/1/1/1
2	NAG	F	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	F	2	2	-	2/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1
2	NAG	Н	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	Н	2	2	-	4/6/23/26	0/1/1/1
2	NAG	Ι	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	Ι	2	2	-	2/6/23/26	0/1/1/1
4	NAG	J	1	1,4	-	0/6/23/26	0/1/1/1
4	FUL	J	2	4	-	_	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
4	J	2	FUL	O5-C1	4.96	1.51	1.43
2	Ι	1	NAG	O5-C1	-2.57	1.39	1.43
4	J	2	FUL	C2-C3	-2.53	1.48	1.52



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	J	2	FUL	O5-C5	2.23	1.48	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
4	J	2	FUL	C1-C2-C3	4.43	115.11	109.67
4	J	2	FUL	C6-C5-C4	-3.11	107.32	113.07
4	J	2	FUL	O5-C5-C4	2.96	114.84	109.52
4	J	2	FUL	O5-C1-C2	2.92	115.28	110.77
3	Ε	2	FUC	O2-C2-C1	2.65	114.58	109.15
2	G	2	NAG	C1-O5-C5	2.65	115.78	112.19
4	J	2	FUL	C1-O5-C5	2.62	118.72	112.78
2	D	1	NAG	O4-C4-C3	-2.37	104.87	110.35
2	F	1	NAG	C1-O5-C5	2.36	115.38	112.19
3	Ε	2	FUC	O5-C5-C4	2.34	113.72	109.52
2	G	1	NAG	C1-O5-C5	2.28	115.28	112.19
4	J	1	NAG	C3-C4-C5	2.27	114.30	110.24
2	F	1	NAG	C4-C3-C2	-2.17	107.84	111.02
2	Ι	1	NAG	O4-C4-C5	-2.15	103.96	109.30

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
2	С	2	NAG	C4-C5-C6-O6
2	F	1	NAG	C4-C5-C6-O6
2	С	2	NAG	O5-C5-C6-O6
2	F	1	NAG	O5-C5-C6-O6
2	F	2	NAG	C4-C5-C6-O6
2	F	2	NAG	O5-C5-C6-O6
2	Н	1	NAG	C4-C5-C6-O6
2	С	1	NAG	O5-C5-C6-O6
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	Н	2	NAG	C8-C7-N2-C2
2	Н	2	NAG	O7-C7-N2-C2
2	Н	1	NAG	O5-C5-C6-O6
2	С	1	NAG	C4-C5-C6-O6
2	Ι	2	NAG	C4-C5-C6-O6
2	G	1	NAG	C4-C5-C6-O6
2	Ι	2	NAG	O5-C5-C6-O6
2	Н	2	NAG	C4-C5-C6-O6

All (21) torsion outliers are listed below:



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	G	1	NAG	O5-C5-C6-O6
2	Н	2	NAG	O5-C5-C6-O6
2	Ι	1	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



























5.6 Ligand geometry (i)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type	Chain	Bos	Link	Bo	ond leng	ths	Bond angles		
WIOI	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	В	601	1	$14,\!14,\!15$	0.74	1 (7%)	$17,\!19,\!21$	0.61	0
5	NAG	А	601	1	14,14,15	0.45	0	17,19,21	0.73	1 (5%)
7	GOL	А	615	-	$5,\!5,\!5$	0.39	0	5, 5, 5	0.28	0
5	NAG	В	609	1	14,14,15	1.07	1(7%)	17,19,21	1.81	1(5%)
5	NAG	В	612	-	14,14,15	0.77	1 (7%)	17,19,21	0.68	0
9	PEG	А	618	-	6,6,6	2.35	4 (66%)	5, 5, 5	1.60	2 (40%)
8	6YC	А	616	-	29,29,29	1.43	1 (3%)	37,38,38	1.17	3 (8%)
5	NAG	В	613	1	14,14,15	0.34	0	17,19,21	0.52	0
8	6YC	В	611	-	29,29,29	1.45	1 (3%)	37,38,38	1.23	<mark>3 (8%)</mark>
7	GOL	А	617	-	$5,\!5,\!5$	0.82	0	5, 5, 5	0.88	0
7	GOL	В	610	-	$5,\!5,\!5$	0.37	0	5, 5, 5	0.30	0
6	FUL	A	604	-	10, 10, 11	1.72	<mark>3 (30%)</mark>	14,14,16	1.37	3 (21%)
5	NAG	В	607	1	14,14,15	0.37	0	17,19,21	0.67	0



Mol	Tune	Chain	Dog	Link	Bo	Bond lengths			ond ang	les
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	В	608	1	14,14,15	0.56	0	17,19,21	0.67	1 (5%)
5	NAG	В	606	-	14,14,15	0.68	1 (7%)	17,19,21	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	В	601	1	-	2/6/23/26	0/1/1/1
5	NAG	А	601	1	-	4/6/23/26	0/1/1/1
7	GOL	А	615	-	-	4/4/4/4	-
5	NAG	В	609	1	-	3/6/23/26	0/1/1/1
5	NAG	В	612	-	-	3/6/23/26	0/1/1/1
9	PEG	А	618	-	-	2/4/4/4	-
8	6YC	А	616	-	-	5/18/29/29	0/3/3/3
5	NAG	В	613	1	-	0/6/23/26	0/1/1/1
8	6YC	В	611	-	-	4/18/29/29	0/3/3/3
7	GOL	А	617	-	-	4/4/4/4	-
7	GOL	В	610	-	-	2/4/4/4	-
6	FUL	А	604	-	-	-	0/1/1/1
5	NAG	В	607	1	-	3/6/23/26	0/1/1/1
5	NAG	В	608	1	-	2/6/23/26	0/1/1/1
5	NAG	В	606	-	-	0/6/23/26	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
8	В	611	6YC	CAR-CAB	-6.12	1.39	1.50
8	А	616	6YC	CAR-CAB	-6.10	1.39	1.50
6	А	604	FUL	O5-C1	3.97	1.50	1.43
5	В	609	NAG	O5-C1	3.71	1.49	1.43
9	А	618	PEG	O2-C3	3.17	1.55	1.42
9	А	618	PEG	O4-C4	2.81	1.56	1.42
6	А	604	FUL	C2-C3	-2.49	1.48	1.52
5	В	606	NAG	O5-C1	-2.45	1.39	1.43
5	В	601	NAG	O5-C1	2.28	1.47	1.43
6	A	604	FUL	O5-C5	2.13	1.48	1.43
9	А	618	PEG	O2-C2	2.11	1.51	1.42



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
9	А	618	PEG	C3-C4	2.08	1.60	1.49
5	В	612	NAG	O5-C1	-2.08	1.40	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
5	В	609	NAG	C1-O5-C5	7.13	121.85	112.19
8	А	616	6YC	CAI-CAH-NAC	-3.96	107.51	113.80
8	В	611	6YC	CAI-CAH-NAC	-3.93	107.56	113.80
6	А	604	FUL	C3-C4-C5	3.18	114.72	109.77
6	А	604	FUL	O5-C5-C4	3.00	114.91	109.52
9	А	618	PEG	C3-O2-C2	2.68	124.90	113.29
5	А	601	NAG	C1-O5-C5	2.43	115.48	112.19
8	В	611	6YC	CAR-CAB-NAC	-2.40	115.67	118.72
8	В	611	6YC	CAN-NAM-CAL	-2.33	109.00	111.71
5	В	608	NAG	C1-O5-C5	2.29	115.30	112.19
6	А	604	FUL	C6-C5-C4	-2.19	109.02	113.07
8	A	616	6YC	CAN-NAM-CAL	-2.19	109.17	111.71
9	А	618	PEG	O1-C1-C2	-2.10	99.62	111.81
8	A	616	6YC	CAR-CAB-NAC	-2.08	116.07	118.72

There are no chirality outliers.

Mol	Chain	\mathbf{Res}	Type	Atoms
7	А	615	GOL	O1-C1-C2-O2
7	А	615	GOL	O1-C1-C2-C3
7	А	615	GOL	C1-C2-C3-O3
8	В	611	6YC	CAO-CAN-NAM-CAQ
7	А	617	GOL	C1-C2-C3-O3
7	А	617	GOL	O2-C2-C3-O3
8	А	616	6YC	CAO-CAN-NAM-CAQ
5	В	608	NAG	O5-C5-C6-O6
5	В	601	NAG	O5-C5-C6-O6
5	А	601	NAG	O5-C5-C6-O6
5	В	601	NAG	C4-C5-C6-O6
5	В	609	NAG	O5-C5-C6-O6
5	А	601	NAG	C4-C5-C6-O6
5	В	608	NAG	C4-C5-C6-O6
5	A	601	NAG	C8-C7-N2-C2
5	A	601	NAG	O7-C7-N2-C2
5	В	$\overline{607}$	NAG	C8-C7-N2-C2

All (38) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	В	607	NAG	O7-C7-N2-C2
5	В	612	NAG	C4-C5-C6-O6
8	В	611	6YC	CAE-CAD-NAC-CAH
8	А	616	6YC	CAE-CAD-NAC-CAH
7	А	617	GOL	O1-C1-C2-O2
8	В	611	6YC	NAC-CAD-CAE-OAF
8	А	616	6YC	NAC-CAD-CAE-OAF
5	В	609	NAG	C4-C5-C6-O6
5	В	612	NAG	O5-C5-C6-O6
8	В	611	6YC	CAE-CAD-NAC-CAB
8	А	616	6YC	CAE-CAD-NAC-CAB
7	А	617	GOL	O1-C1-C2-C3
7	В	610	GOL	O1-C1-C2-C3
7	А	615	GOL	O2-C2-C3-O3
5	В	607	NAG	O5-C5-C6-O6
9	А	618	PEG	O1-C1-C2-O2
5	В	612	NAG	C3-C2-N2-C7
9	A	618	PEG	C4-C3-O2-C2
5	В	609	NAG	C3-C2-N2-C7
7	В	610	GOL	O1-C1-C2-O2
8	A	616	6YC	CAD-CAE-OAF-CAG

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	А	601	NAG	1	0
7	А	615	GOL	2	0
9	А	618	PEG	1	0
7	А	617	GOL	1	0
5	В	607	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



Ligand 6YC A 616 Bond lengths Bond angles Torsions Rings

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$<$ RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	528/574~(91%)	0.51	24 (4%) 33 36	45, 74, 101, 118	1 (0%)
1	В	525/574~(91%)	0.57	26 (4%) 28 31	47, 73, 104, 129	0
All	All	1053/1148~(91%)	0.54	50 (4%) 31 34	45, 73, 102, 129	1 (0%)

All (50) RSRZ outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	RSRZ
1	В	530	LEU	17.4
1	В	52	TRP	6.8
1	А	530	LEU	6.3
1	В	53	SER	6.2
1	А	53	SER	6.0
1	В	49	LEU	4.3
1	В	54	ASP	4.3
1	В	11	GLY	4.1
1	А	4	ILE	4.0
1	А	529	VAL	3.9
1	А	156	LEU	3.3
1	А	3	ASP	3.1
1	В	55	ILE	3.1
1	В	10	ASN	3.0
1	А	22	GLY	3.0
1	В	529	VAL	3.0
1	В	8	THR	2.9
1	В	454	ASP	2.9
1	A	352	GLU	2.9
1	В	266	ASN	2.8
1	A	278	PHE	2.8
1	В	185	PHE	2.8
1	В	50	THR	2.7
1	A	264	LEU	2.7



Mol	Chain	Res	Type	RSRZ
1	А	263	CYS	2.5
1	В	371	PHE	2.5
1	А	262	LYS	2.5
1	В	161	GLU	2.4
1	В	9	LYS	2.3
1	А	419	TYR	2.3
1	В	412	TRP	2.3
1	А	259	GLU	2.3
1	А	380	GLN	2.2
1	А	364[A]	PHE	2.2
1	А	159	ASN	2.2
1	А	55	ILE	2.2
1	А	274	LEU	2.2
1	В	291	GLY	2.2
1	В	378	ASP	2.2
1	В	453	ARG	2.2
1	А	266	ASN	2.2
1	В	490	TRP	2.2
1	В	7	ALA	2.1
1	А	237	TYR	2.1
1	В	56	TRP	2.1
1	В	357	PHE	2.1
1	В	383	GLU	2.1
1	А	275	ASN	2.1
1	А	282	TYR	2.0
1	A	23	GLY	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{\AA}^2)$	Q<0.9
2	NAG	Η	2	14/15	0.72	0.33	$145,\!151,\!158,\!159$	0
2	NAG	Н	1	14/15	0.78	0.14	$98,\!127,\!141,\!146$	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
3	NAG	Е	1	14/15	0.81	0.29	$120,\!131,\!139,\!139$	0
2	NAG	G	2	14/15	0.82	0.21	$75,\!94,\!100,\!101$	14
2	NAG	D	2	14/15	0.82	0.40	114,133,142,145	0
2	NAG	С	1	14/15	0.83	0.19	$87,\!111,\!121,\!132$	0
2	NAG	Ι	2	14/15	0.84	0.48	$107,\!135,\!138,\!142$	0
2	NAG	C	2	14/15	0.87	0.27	$90,\!124,\!132,\!135$	0
2	NAG	D	1	14/15	0.88	0.27	$74,\!93,\!117,\!125$	0
2	NAG	F	1	14/15	0.88	0.15	$95,\!105,\!125,\!132$	0
2	NAG	G	1	14/15	0.89	0.21	70,81,93,95	14
2	NAG	F	2	14/15	0.90	0.33	113,122,131,132	0
4	NAG	J	1	14/15	0.90	0.17	$83,\!99,\!109,\!116$	0
3	FUC	Ē	2	10/11	0.91	0.43	105,111,118,121	0
2	NAG	I	1	14/15	0.91	0.23	$100,\!113,\!129,\!133$	0
4	FUL	J	2	10/11	0.93	0.32	$91,\!101,\!107,\!124$	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.































6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	\mathbf{Res}	Atoms	RSCC	RSR	${f B}$ -factors(${ m \AA}^2)$	Q<0.9
7	GOL	А	617	6/6	0.72	0.18	$89,\!97,\!103,\!109$	0
6	FUL	А	604	10/11	0.74	0.48	$109,\!116,\!125,\!133$	0
7	GOL	А	615	6/6	0.78	0.31	$74,\!83,\!96,\!96$	0
5	NAG	В	613	14/15	0.79	0.27	$109,\!129,\!137,\!137$	0
5	NAG	В	601	14/15	0.81	0.20	$104,\!127,\!134,\!137$	0
5	NAG	В	606	14/15	0.81	0.43	$93,\!112,\!118,\!119$	0
5	NAG	В	609	14/15	0.82	0.20	$106,\!119,\!133,\!134$	0
9	PEG	А	618	7/7	0.84	0.14	$93,\!96,\!100,\!103$	0
5	NAG	В	612	14/15	0.87	0.14	$85,\!105,\!120,\!127$	0
8	6YC	В	611	27/27	0.88	0.29	$58,\!79,\!105,\!110$	27
8	6YC	А	616	27/27	0.90	0.23	$67,\!96,\!117,\!128$	9
5	NAG	А	601	14/15	0.91	0.25	$86,\!91,\!107,\!108$	0
5	NAG	В	607	14/15	0.91	0.12	$95,\!106,\!113,\!131$	0
7	GOL	В	610	6/6	0.92	0.20	77,78,93,94	0
5	NAG	B	608	14/15	0.94	0.12	$81,\!91,\!101,\!105$	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

