



Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 11:20 pm BST

PDB ID : 2LRP
Title : Solution structure, dynamics and binding studies of CtCBM11
Authors : Viegas, A.; Cabrita, E.J.
Deposited on : 2012-04-11

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 81%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	172	83% 12% • 5%

2 Ensemble composition and analysis i

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:6-A:169 (164)	0.81	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 6, 7, 8, 9, 11, 13, 16, 20
2	3, 5
3	18, 19
4	12, 17
Single-model clusters	10; 14; 15

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2613 atoms, of which 1276 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Endoglucanase H.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	172	2611	846	1276	216	267	6	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP P16218
A	2	ALA	-	EXPRESSION TAG	UNP P16218
A	3	SER	-	EXPRESSION TAG	UNP P16218
A	171	LEU	-	EXPRESSION TAG	UNP P16218
A	172	GLU	-	EXPRESSION TAG	UNP P16218

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	
			Total	Ca
2	A	2	2	2

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Endoglucanase H




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Endoglucanase H




4.2.2 Score per residue for model 2 (medoid)

- Molecule 1: Endoglucanase H




4.2.3 Score per residue for model 3

- Molecule 1: Endoglucanase H




4.2.4 Score per residue for model 4

- Molecule 1: Endoglucanase H




4.2.5 Score per residue for model 5

- Molecule 1: Endoglucanase H




4.2.6 Score per residue for model 6

- Molecule 1: Endoglucanase H



4.2.7 Score per residue for model 7

- Molecule 1: Endoglucanase H





4.2.8 Score per residue for model 8

- Molecule 1: Endoglucanase H

Chain A: 80% 13% 5%



4.2.9 Score per residue for model 9

- Molecule 1: Endoglucanase H



Chain A: 80% 12% 5%



4.2.10 Score per residue for model 10

- Molecule 1: Endoglucanase H

Chain A: 75% 17% 5%



4.2.11 Score per residue for model 11


- Molecule 1: Endoglucanase H

Chain A: 81% 12% 5%




4.2.12 Score per residue for model 12

- Molecule 1: Endoglucanase H




4.2.13 Score per residue for model 13

- Molecule 1: Endoglucanase H




4.2.14 Score per residue for model 14

- Molecule 1: Endoglucanase H




4.2.15 Score per residue for model 15

- Molecule 1: Endoglucanase H



4.2.16 Score per residue for model 16

- Molecule 1: Endoglucanase H






4.2.17 Score per residue for model 17

- Molecule 1: Endoglucanase H


Chain A: 77% 15% • • 5%



4.2.18 Score per residue for model 18

- Molecule 1: Endoglucanase H

Chain A: 81% 11% • • 5%



4.2.19 Score per residue for model 19

- Molecule 1: Endoglucanase H


Chain A: 81% 10% • • 5%



4.2.20 Score per residue for model 20

- Molecule 1: Endoglucanase H

Chain A: 77% 12% 5% • 5%



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *molecular dynamics*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
AMBER	refinement	9.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	1790
Number of shifts mapped to atoms	1790
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	81%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1281	1219	1219	9±6
All	All	25660	24380	24360	189

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:76:LYS:HE3	1:A:113:TRP:CE3	0.77	2.13	20	6
1:A:19:TRP:CZ2	1:A:58:TYR:CE2	0.76	2.73	4	10
1:A:105:TYR:CD2	1:A:107:ILE:HD12	0.75	2.17	16	2
1:A:19:TRP:CZ3	1:A:58:TYR:CG	0.64	2.85	9	10
1:A:19:TRP:CH2	1:A:58:TYR:CE2	0.62	2.88	17	10
1:A:76:LYS:HE3	1:A:113:TRP:CD2	0.62	2.30	20	2
1:A:76:LYS:CE	1:A:113:TRP:CE3	0.62	2.83	20	1
1:A:18:ASN:ND2	1:A:19:TRP:CZ2	0.61	2.69	5	10
1:A:19:TRP:CH2	1:A:58:TYR:CZ	0.61	2.89	2	7
1:A:19:TRP:CH2	1:A:58:TYR:CD2	0.59	2.91	5	7
1:A:22:TYR:CE1	1:A:57:VAL:HG23	0.58	2.32	6	1
1:A:19:TRP:CZ2	1:A:58:TYR:CD2	0.58	2.92	5	5
1:A:19:TRP:CZ3	1:A:58:TYR:CD2	0.58	2.91	5	10
1:A:11:ASP:HA	1:A:13:PHE:CE2	0.55	2.36	18	1
1:A:76:LYS:HE2	1:A:113:TRP:CE3	0.54	2.36	9	1
1:A:71:ILE:HD11	1:A:118:ILE:HD12	0.54	1.80	16	1
1:A:19:TRP:CH2	1:A:58:TYR:CE1	0.53	2.96	2	2
1:A:18:ASN:ND2	1:A:19:TRP:CE2	0.52	2.78	4	9
1:A:105:TYR:CE2	1:A:118:ILE:HD13	0.52	2.39	5	3
1:A:76:LYS:HD3	1:A:113:TRP:CE3	0.52	2.39	17	3
1:A:19:TRP:CE3	1:A:58:TYR:CD2	0.52	2.97	3	7
1:A:18:ASN:ND2	1:A:19:TRP:CH2	0.51	2.78	2	4
1:A:46:TYR:CE2	1:A:54:TRP:CD1	0.51	2.98	14	7
1:A:46:TYR:CE1	1:A:54:TRP:HB2	0.51	2.40	19	2
1:A:76:LYS:HD2	1:A:113:TRP:HA	0.50	1.82	10	2
1:A:58:TYR:CG	1:A:148:ILE:HD11	0.50	2.41	10	7
1:A:54:TRP:CZ3	1:A:152:TYR:CG	0.50	2.99	8	1
1:A:58:TYR:CG	1:A:148:ILE:HD12	0.49	2.43	5	2
1:A:46:TYR:CE2	1:A:54:TRP:NE1	0.48	2.81	11	1
1:A:63:GLY:HA2	1:A:65:TRP:CZ3	0.48	2.43	4	1
1:A:57:VAL:HG13	1:A:149:HIS:CE1	0.48	2.44	2	6
1:A:118:ILE:HG22	1:A:119:PRO:HD2	0.47	1.86	20	1
1:A:12:ASP:H	1:A:13:PHE:HB3	0.47	1.70	19	1
1:A:76:LYS:HD3	1:A:113:TRP:CD2	0.47	2.45	8	3
1:A:76:LYS:HD2	1:A:113:TRP:CZ3	0.47	2.45	17	1
1:A:58:TYR:CG	1:A:148:ILE:CD1	0.46	2.97	5	2
1:A:46:TYR:CD2	1:A:54:TRP:CD1	0.46	3.03	11	1
1:A:19:TRP:CZ3	1:A:58:TYR:CD1	0.46	3.04	2	2
1:A:54:TRP:CZ2	1:A:152:TYR:HB3	0.45	2.46	14	3
1:A:58:TYR:CD1	1:A:148:ILE:HD11	0.45	2.45	3	6
1:A:12:ASP:H	1:A:13:PHE:CB	0.45	2.25	19	1
1:A:58:TYR:CD1	1:A:148:ILE:CD1	0.45	2.99	3	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:102:HIS:CE1	1:A:129:TYR:CE1	0.45	3.05	4	1
1:A:76:LYS:HE3	1:A:113:TRP:CZ3	0.45	2.47	11	3
1:A:46:TYR:CE1	1:A:54:TRP:CB	0.44	3.00	9	2
1:A:11:ASP:HA	1:A:13:PHE:CD1	0.44	2.48	19	1
1:A:76:LYS:CE	1:A:161:VAL:HG13	0.44	2.42	7	1
1:A:70:LYS:HA	1:A:120:PHE:H	0.44	1.72	20	1
1:A:11:ASP:HA	1:A:13:PHE:CZ	0.44	2.48	18	1
1:A:76:LYS:CE	1:A:113:TRP:CD2	0.44	3.01	9	1
1:A:46:TYR:CE1	1:A:54:TRP:HB3	0.43	2.48	9	1
1:A:76:LYS:HE3	1:A:161:VAL:HG13	0.43	1.90	7	1
1:A:105:TYR:CE2	1:A:107:ILE:HD12	0.43	2.48	16	1
1:A:76:LYS:HE3	1:A:113:TRP:CE2	0.43	2.49	6	1
1:A:76:LYS:HE2	1:A:113:TRP:CE2	0.43	2.49	3	1
1:A:105:TYR:CE2	1:A:107:ILE:CD1	0.42	3.02	2	1
1:A:37:LYS:HE2	1:A:113:TRP:CE3	0.42	2.49	13	1
1:A:105:TYR:HD2	1:A:107:ILE:HD12	0.42	1.68	20	1
1:A:54:TRP:CE3	1:A:152:TYR:CD1	0.42	3.08	8	1
1:A:58:TYR:CD2	1:A:148:ILE:HD12	0.42	2.49	5	1
1:A:76:LYS:HE2	1:A:161:VAL:CG2	0.41	2.45	20	1
1:A:76:LYS:HE3	1:A:113:TRP:HA	0.40	1.93	9	1
1:A:58:TYR:CD2	1:A:148:ILE:HD11	0.40	2.51	10	1
1:A:19:TRP:CZ2	1:A:58:TYR:CZ	0.40	3.09	2	1

5.2 Torsion angles [i](#)

5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/172 (95%)	149±4 (91±3%)	13±4 (8±2%)	2±1 (1±1%)	20	68
All	All	3280/3440 (95%)	2982 (91%)	266 (8%)	32 (1%)	20	68

All 22 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	79	ASP	3
1	A	66	SER	3
1	A	164	ASN	3
1	A	13	PHE	2
1	A	113	TRP	2
1	A	120	PHE	2
1	A	14	GLU	2
1	A	16	VAL	1
1	A	100	GLY	1
1	A	123	PHE	1
1	A	63	GLY	1
1	A	26	GLY	1
1	A	64	ASP	1
1	A	55	GLY	1
1	A	80	GLY	1
1	A	12	ASP	1
1	A	24	GLY	1
1	A	121	SER	1
1	A	6	GLY	1
1	A	81	SER	1
1	A	109	PRO	1
1	A	119	PRO	1

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/145 (97%)	126±3 (90±2%)	14±3 (10±2%)	12	57
All	All	2800/2900 (97%)	2529 (90%)	271 (10%)	12	57

All 76 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	134	GLN	16
1	A	32	LYS	16
1	A	54	TRP	15
1	A	76	LYS	13

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	135	ASP	12
1	A	73	PHE	12
1	A	107	ILE	11
1	A	161	VAL	10
1	A	65	TRP	10
1	A	58	TYR	9
1	A	124	ARG	8
1	A	142	LEU	6
1	A	18	ASN	6
1	A	21	SER	6
1	A	126	ARG	5
1	A	148	ILE	5
1	A	40	ASN	5
1	A	118	ILE	5
1	A	127	LEU	5
1	A	60	LEU	4
1	A	87	PHE	4
1	A	113	TRP	4
1	A	56	THR	3
1	A	22	TYR	3
1	A	129	TYR	3
1	A	166	LYS	3
1	A	12	ASP	3
1	A	130	GLN	3
1	A	114	LYS	3
1	A	155	ASN	3
1	A	86	ARG	3
1	A	140	LEU	3
1	A	42	MET	2
1	A	13	PHE	2
1	A	84	GLU	2
1	A	67	LYS	2
1	A	164	ASN	2
1	A	17	LEU	2
1	A	108	THR	2
1	A	143	ASP	2
1	A	152	TYR	2
1	A	46	TYR	2
1	A	144	ASN	1
1	A	160	PHE	1
1	A	69	LEU	1
1	A	83	ASN	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	123	PHE	1
1	A	44	VAL	1
1	A	112	SER	1
1	A	70	LYS	1
1	A	88	MET	1
1	A	125	ARG	1
1	A	31	THR	1
1	A	30	SER	1
1	A	51	ASP	1
1	A	62	ASP	1
1	A	79	ASP	1
1	A	28	LYS	1
1	A	150	PHE	1
1	A	154	ASN	1
1	A	72	SER	1
1	A	64	ASP	1
1	A	25	GLU	1
1	A	11	ASP	1
1	A	151	MET	1
1	A	38	THR	1
1	A	168	ILE	1
1	A	29	VAL	1
1	A	23	SER	1
1	A	74	ASP	1
1	A	77	SER	1
1	A	14	GLU	1
1	A	157	SER	1
1	A	105	TYR	1
1	A	89	ILE	1
1	A	156	LYS	1

5.2.3 RNA

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.5 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

5.6 Other polymers [i](#)

There are no such molecules in this entry.

5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 81% for the well-defined parts and 80% for the entire structure.

6.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

6.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1790
Number of shifts mapped to atoms	1790
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	11

6.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	161	2.62 ± 0.12	Should be applied
$^{13}\text{C}_\beta$	142	2.20 ± 0.14	Should be applied
$^{13}\text{C}'$	157	2.89 ± 0.14	Should be applied
^{15}N	158	0.48 ± 0.32	None needed (< 0.5 ppm)

6.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 81%, i.e. 1569 atoms were assigned a chemical shift out of a possible 1936. 0 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	772/810 (95%)	310/323 (96%)	309/328 (94%)	153/159 (96%)
Sidechain	777/921 (84%)	477/540 (88%)	290/347 (84%)	10/34 (29%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	20/205 (10%)	15/107 (14%)	0/90 (0%)	5/8 (62%)
Overall	1569/1936 (81%)	802/970 (83%)	599/765 (78%)	168/201 (84%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 80%, i.e. 1611 atoms were assigned a chemical shift out of a possible 2015. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	795/850 (94%)	319/339 (94%)	318/344 (92%)	158/167 (95%)
Sidechain	796/960 (83%)	488/562 (87%)	298/364 (82%)	10/34 (29%)
Aromatic	20/205 (10%)	15/107 (14%)	0/90 (0%)	5/8 (62%)
Overall	1611/2015 (80%)	822/1008 (82%)	616/798 (77%)	173/209 (83%)

6.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	37	LYS	HB2	-0.36	3.03 – 0.53	-8.5
1	A	75	ILE	CG1	13.16	36.54 – 18.94	-8.3
1	A	58	TYR	HB2	0.15	4.76 – 1.06	-7.5
1	A	140	LEU	HB2	-0.88	3.32 – -0.08	-7.3
1	A	75	ILE	CG2	27.02	24.63 – 10.43	6.7
1	A	86	ARG	HD2	1.64	4.27 – 1.97	-6.4
1	A	76	LYS	HD2	0.18	2.76 – 0.46	-6.2
1	A	86	ARG	HD3	1.64	4.36 – 1.86	-5.9
1	A	101	GLU	HB3	3.28	3.10 – 0.90	5.8
1	A	44	VAL	HB	0.30	3.59 – 0.39	-5.3
1	A	74	ASP	HA	6.15	6.15 – 3.05	5.0

6.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

