

# wwPDB NMR Structure Validation Summary Report (i)

Jun 5, 2023 – 01:49 AM EDT

PDB ID : 2LSG BMRB ID : 18431

Title : Solution structure of the mouse Rev1 C-terminal domain

Authors : Liu, J.; Wojtaszek, J.; Zhou, P.

Deposited on : 2012-04-30

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

wwPDB-ShiftChecker : v1.2 BMRB Restraints Analysis : v1.2

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment is 86%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	NMR archive
Metric	$(\#  ext{Entries})$	$(\#  ext{Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	A	104	66%	15%	12%	7%	



# 2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid mo					
1	A:18-A:102 (85)	0.37	1		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 6, 8, 11, 12, 15, 16, 18, 20
2	3, 5, 7, 9
Single-model clusters	10; 13; 14; 17; 19



# 3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1575 atoms, of which 794 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called DNA repair protein REV1.

Mol	Chain	Residues	Atoms				Trace		
1	Λ	07	Total	С	Н	N	О	S	0
	A	97	1575	502	794	121	153	5	U

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP Q920Q2
A	2	SER	-	expression tag	UNP Q920Q2
A	3	GLY	-	expression tag	UNP Q920Q2
A	4	GLY	-	expression tag	UNP Q920Q2




# 4 Residue-property plots (i)

#### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: DNA repair protein REV1



# 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

• Molecule 1: DNA repair protein REV1





#### Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: simulated annealing.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: target function.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
CYANA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1227
Number of shifts mapped to atoms	1180
Number of unparsed shifts	0
Number of shifts with mapping errors	47
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	86%



# 6 Model quality (i)

### 6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	704	716	716	10±3
All	All	14080	14320	14320	208

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

5 of 56 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:48:THR:HG23	1:A:92:VAL:HG21	0.87	1.44	13	13
1:A:29:ILE:HD12	1:A:69:LEU:HD13	0.79	1.53	16	13
1:A:25:LEU:HD11	1:A:62:VAL:HG13	0.69	1.62	4	3
1:A:23:THR:HG22	1:A:27:GLU:OE1	0.64	1.92	20	1
1:A:28:TRP:CZ3	1:A:43:VAL:HG11	0.63	2.28	9	13

# 6.3 Torsion angles (i)

#### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	85/104 (82%)	83±1 (98±1%)	2±1 (2±1%)	0±0 (0±0%)	100	100
All	All	1700/2080 (82%)	1669 (98%)	31 (2%)	0 (0%)	100	100

There are no Ramachandran outliers.

#### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Chain Analysed Rotameric Outl		Outliers	Percentiles
1	A	83/94 (88%)	77±2 (92±2%)	6±2 (8±2%)	16 64
All	All	1660/1880 (88%)	1533 (92%)	127 (8%)	16 64

5 of 25 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	56	LEU	20
1	A	93	LEU	20
1	A	70	MET	19
1	A	59	LEU	8
1	A	60	ASP	8

### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

# 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.



# 6.6 Ligand geometry (i)

There are no ligands in this entry.

# 6.7 Other polymers (i)

There are no such molecules in this entry.

# 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 86% for the well-defined parts and 86% for the entire structure.

#### 7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned\_chem\_shift\_list\_1

#### 7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1227
Number of shifts mapped to atoms	1180
Number of unparsed shifts	0
Number of shifts with mapping errors	47
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

• No matching atom found in the structure. First 5 (of 47) occurrences are reported below.

T:at ID	Clasica	Dag	Т	A 4 a		Shift Data				
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity			
1	A	3	GLY	HA2	3.977	0.000				
1	A	3	GLY	HA3	3.977	0.000	•			
1	A	3	GLY	CA	45.388	0.000				
1	A	4	GLY	Н	8.251	0.001	•			
1	A	4	GLY	HA2	3.895	0.012				
1	A	4	GLY	HA3	3.895	0.012				
1	A	4	GLY	CA	44.934	0.054				
1	A	4	GLY	N	108.668	0.040	•			
1	A	5	PHE	Н	8.161	0.003				
1	A	5	PHE	HA	4.561	0.010				
1	A	5	PHE	HB2	3.016	0.014	•			
1	A	5	PHE	HB3	3.016	0.014				
1	A	5	PHE	HD1	7.19	0.009	•			
1	A	5	PHE	HD2	7.19	0.009	•			

Continued on next page...



 $Continued\ from\ previous\ page...$ 

	a from pr			<b>A</b> .		l	
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	5	PHE	HE1	7.269 0.011		
1	A	5	PHE	HE2	7.269	0.011	
1	A	5	PHE	HZ	7.235	0.006	
1	A	5	PHE	CA	57.634	0.032	
1	A	5	PHE	СВ	39.615	0.172	•
1	A	5	PHE	CD1	131.17	0.068	
1	A	5	PHE	CE1	130.867	0.010	•
1	A	5	PHE	CZ	129.159	0.098	
1	A	5	PHE	N	120.371	0.132	
1	A	6	ARG	Н	8.161	0.001	•
1	A	6	ARG	HA	4.544	0.010	•
1	A	6	ARG	HB2	1.633	0.009	•
1	A	6	ARG	HB3	1.753	0.010	•
1	A	6	ARG	HG2	1.55	0.012	•
1	A	6	ARG	HG3	1.55	0.012	•
1	A	6	ARG	HD2	3.155	0.012	
1	A	6	ARG	HD3	3.155	0.012	•
1	A	6	ARG	CA	53.393	0.218	
1	A	6	ARG	СВ	30.495	0.086	•
1	A	6	ARG	CG	26.505	0.171	•
1	A	6	ARG	CD	43.162	0.118	
1	A	6	ARG	N	125.018	0.134	•
1	A	7	PRO	HA	4.298	0.012	
1	A	7	PRO	HB2	1.917	0.004	•
1	A	7	PRO	HB3	2.274	0.010	•
1	A	7	PRO	HG2	1.957	0.009	·
1	A	7	PRO	HG3	1.957	0.009	•
1	A	7	PRO	HD2	3.562	0.009	•
1	A	7	PRO	HD3	3.562	0.009	•
1	A	7	PRO	CA	62.909	0.112	•
1	A	7	PRO	СВ	31.862	0.154	•
1	A	7	PRO	CG	27.039	0.100	•
1	A	7	PRO	CD	50.376	0.043	

### 7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, $ppm$	Suggested action
$^{13}\mathrm{C}_{\alpha}$	102	$-0.41 \pm 0.28$	None needed ( $< 0.5 \text{ ppm}$ )
$^{13}\mathrm{C}_{\beta}$	98	$0.38 \pm 0.17$	None needed (< 0.5 ppm)

Continued on next page...



Continued from previous page...

Nucleus	# values	Correction $\pm$ precision, $ppm$	Suggested action
<sup>13</sup> C′	0		None (insufficient data)
$^{15}N$	91	$0.33 \pm 0.26$	None needed ( $< 0.5 \text{ ppm}$ )

#### 7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 86%, i.e. 1060 atoms were assigned a chemical shift out of a possible 1229. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	328/424~(77%)	166/170 (98%)	85/170~(50%)	77/84 (92%)
Sidechain	667/724~(92%)	455/471 (97%)	205/232~(88%)	7/21 (33%)
Aromatic	65/81 (80%)	39/39 (100%)	24/40 (60%)	2/2 (100%)
Overall	1060/1229~(86%)	660/680 (97%)	314/442 (71%)	86/107 (80%)

#### 7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.


List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	35	PRO	HG2	-0.20	0.41 - 3.45	-7.0
1	A	22	LYS	HG2	-0.09	0.13 - 2.61	-5.9
1	A	78	TRP	HH2	5.17	5.24 - 8.73	-5.2

### 7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:







# 8 NMR restraints analysis (i)

# 8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1969
Intra-residue ( $ i-j =0$ )	497
Sequential ( $ i-j =1$ )	481
Medium range ( $ i-j >1$ and $ i-j <5$ )	622
Long range ( i-j ≥5)	369
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	35
Number of restraints per residue	18.9
Number of long range restraints per residue <sup>1</sup>	3.5

<sup>&</sup>lt;sup>1</sup>Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

# 8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

# 8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	1.4	0.16
0.2-0.5 (Medium)	0.3	0.35
>0.5 (Large)	None	None



# 8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than  $1^{\circ}$  are not included in the calculation. There are no dihedral-angle violations



# 9 Distance violation analysis (i)

# 9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destroints tune	Count	<b>%</b> <sup>1</sup>	Vio	lated	3	Consis	tentl	${ m y~Violated^4}$
Restraints type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue ( i-j =0)	497	25.2	3	0.6	0.2	0	0.0	0.0
Backbone-Backbone	4	0.2	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	386	19.6	3	0.8	0.2	0	0.0	0.0
Sidechain-Sidechain	107	5.4	0	0.0	0.0	0	0.0	0.0
Sequential ( i-j =1)	481	24.4	4	0.8	0.2	0	0.0	0.0
Backbone-Backbone	114	5.8	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	316	16.0	3	0.9	0.2	0	0.0	0.0
Sidechain-Sidechain	51	2.6	1	2.0	0.1	0	0.0	0.0
Medium range ( $ i-j >1 \&  i-j <5$ )	622	31.6	8	1.3	0.4	0	0.0	0.0
Backbone-Backbone	178	9.0	2	1.1	0.1	0	0.0	0.0
Backbone-Sidechain	307	15.6	5	1.6	0.3	0	0.0	0.0
Sidechain-Sidechain	137	7.0	1	0.7	0.1	0	0.0	0.0
Long range ( i-j ≥5)	369	18.7	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	5	0.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	116	5.9	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	248	12.6	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1969	100.0	15	0.8	0.8	0	0.0	0.0
Backbone-Backbone	301	15.3	2	0.7	0.1	0	0.0	0.0
Backbone-Sidechain	1125	57.1	11	1.0	0.6	0	0.0	0.0
Sidechain-Sidechain	543	27.6	2	0.4	0.1	0	0.0	0.0

 $<sup>^1</sup>$  percentage calculated with respect to the total number of distance restraints,  $^2$  percentage calculated with respect to the number of restraints in a particular restraint category,  $^3$  violated in at least one model,  $^4$  violated in all the models



#### 9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

### 9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID		Nun	nber o	f viola	ations	3	Maan (Å)	Max (Å)	${ m SD}^6$ (Å)	Modian (Å)
Model ID	$IR^1$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Mean (Å)	Max (A)	$SD^*(A)$	Median (Å)
1	0	1	0	0	0	1	0.12	0.12	0.0	0.12
2	0	1	0	0	0	1	0.13	0.13	0.0	0.13
3	0	1	0	0	0	1	0.12	0.12	0.0	0.12
4	0	0	0	0	0	0	0.0	0.0	0.0	0.0
5	0	0	0	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	1	1	0	0	0	2	0.12	0.13	0.01	0.12
8	0	1	0	0	0	1	0.12	0.12	0.0	0.12
9	0	1	0	0	0	1	0.13	0.13	0.0	0.13
10	0	0	0	0	0	0	0.0	0.0	0.0	0.0
11	0	1	1	0	0	2	0.18	0.22	0.04	0.18

Continued on next page...




Continued from previous page...

Madal ID	Number of violations					3	Mean (Å)	Max (Å)	${ m SD}^6$ (Å)	Median (Å)
Model ID	$IR^1$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	0	2	0	0	0	2	0.15	0.16	0.02	0.15
13	1	1	0	0	0	2	0.12	0.13	0.01	0.12
14	0	1	2	0	0	3	0.13	0.15	0.01	0.13
15	1	1	1	0	0	3	0.18	0.31	0.09	0.12
16	0	1	2	0	0	3	0.19	0.35	0.11	0.12
17	0	0	0	0	0	0	0.0	0.0	0.0	0.0
18	0	2	1	0	0	3	0.16	0.21	0.04	0.15
19	1	2	2	0	0	5	0.15	0.26	0.06	0.13
20	1	2	1	0	0	4	0.17	0.26	0.05	0.15

 $<sup>^1</sup>$ Intra-residue restraints,  $^2$ Sequential restraints,  $^3$ Medium range restraints,  $^4$ Long range restraints,  $^5$ Inter-chain restraints,  $^6$ Standard deviation

### 9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



#### 9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1954(IR:494, SQ:477, MR:614, LR:369, IC:0) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	Fraction of the ensemble			
$IR^1$	$SQ^2$	$MR^3$	$LR^4$	$IC^5$	Total	Count <sup>6</sup>	%
1	2	6	0	0	9	1	5.0
2	1	2	0	0	5	2	10.0
0	0	0	0	0	0	3	15.0
0	0	0	0	0	0	4	20.0
0	0	0	0	0	0	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	1	0	0	0	1	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

 $<sup>^1</sup>$ Intra-residue restraints,  $^2$ Sequential restraints,  $^3$ Medium range restraints,  $^4$ Long range restraints,  $^5$ Inter-chain restraints,  $^6$  Number of models with violations



#### 9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




# 9.4 Most violated distance restraints in the ensemble (i)

# 9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





#### 9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	$Models^1$	Mean (Å)	$SD^1$ (Å)	Median (Å)
(1,1355)	1:A:73:SER:HA	1:A:74:VAL:HB	15	0.13	0.0	0.13
(1,389)	1:A:31:THR:HA	1:A:31:THR:HG21	2	0.26	0.0	0.26
(1,389)	1:A:31:THR:HA	1:A:31:THR:HG22	2	0.26	0.0	0.26
(1,389)	1:A:31:THR:HA	1:A:31:THR:HG23	2	0.26	0.0	0.26
(1,1917)	1:A:102:LYS:HG2	1:A:104:THR:H	2	0.26	0.05	0.26
(1,1917)	1:A:102:LYS:HG3	1:A:104:THR:H	2	0.26	0.05	0.26
(1,397)	1:A:31:THR:HB	1:A:32:ILE:H	2	0.15	0.02	0.15
(1,291)	1:A:27:GLU:HA	1:A:31:THR:HG21	2	0.13	0.01	0.13
(1,291)	1:A:27:GLU:HA	1:A:31:THR:HG22	2	0.13	0.01	0.13
(1,291)	1:A:27:GLU:HA	1:A:31:THR:HG23	2	0.13	0.01	0.13
(1,962)	1:A:58:LYS:HA	1:A:58:LYS:HD2	2	0.11	0.0	0.11
(1,962)	1:A:58:LYS:HA	1:A:58:LYS:HD3	2	0.11	0.0	0.11


<sup>&</sup>lt;sup>1</sup>Number of violated models, <sup>2</sup>Standard deviation



### 9.5 All violated distance restraints (i)

#### 9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



### 9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,66)	1:A:13:ALA:HB1	1:A:15:ALA:HA	16	0.35
(1,66)	1:A:13:ALA:HB2	1:A:15:ALA:HA	16	0.35
(1,66)	1:A:13:ALA:HB3	1:A:15:ALA:HA	16	0.35
(1,1917)	1:A:102:LYS:HG2	1:A:104:THR:H	15	0.31
(1,1917)	1:A:102:LYS:HG3	1:A:104:THR:H	15	0.31
(1,389)	1:A:31:THR:HA	1:A:31:THR:HG21	19	0.26
(1,389)	1:A:31:THR:HA	1:A:31:THR:HG22	19	0.26
(1,389)	1:A:31:THR:HA	1:A:31:THR:HG23	19	0.26
(1,389)	1:A:31:THR:HA	1:A:31:THR:HG21	20	0.26
(1,389)	1:A:31:THR:HA	1:A:31:THR:HG22	20	0.26
(1,389)	1:A:31:THR:HA	1:A:31:THR:HG23	20	0.26

Continued on next page...



### Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,465)	1:A:37:GLU:HB3	1:A:40:ILE:HD11	11	0.22
(1,465)	1:A:37:GLU:HB3	1:A:40:ILE:HD12	11	0.22
(1,465)	1:A:37:GLU:HB3	1:A:40:ILE:HD13	11	0.22
(1,1917)	1:A:102:LYS:HG2	1:A:104:THR:H	18	0.21
(1,1917)	1:A:102:LYS:HG3	1:A:104:THR:H	18	0.21
(1,948)	1:A:57:GLU:H	1:A:58:LYS:HE2	12	0.16
(1,948)	1:A:57:GLU:H	1:A:58:LYS:HE3	12	0.16
(1,397)	1:A:31:THR:HB	1:A:32:ILE:H	20	0.16
(1,1915)	1:A:102:LYS:HE2	1:A:104:THR:H	14	0.15
(1,1915)	1:A:102:LYS:HE3	1:A:104:THR:H	14	0.15
(1,1913)	1:A:102:LYS:HE2	1:A:103:VAL:HG11	18	0.15



# 10 Dihedral-angle violation analysis (i)

Dihedral angle analysis failed due to data error in the dihedral angle restraints, possibly missing target value

