

# wwPDB X-ray Structure Validation Summary Report (i)

#### Sep 11, 2023 – 10:27 PM EDT

PDB ID	:	4LS3
Title	:	THE crystal STRUCTURE OF HELICOBACTER PYLORI
		CEUE(HP1561)/NI-HIS COMPL
Authors	:	Salamina, M.; Shaik, M.M.; Cendron, L.; Zanotti, G.
Deposited on		
Resolution	:	1.70  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	А	333	87%	•• 9%			
1	В	333	% <b>8</b> 4%	6% 9%			



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5838 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Δ	302	Total C N O S O	C N O S O	N O S		0	0		
1 A	302	2399	1569	390	435	5	0	0	U	
1	D	302	Total	С	Ν	0	S	0	0	0
	502	2399	1569	390	435	5	0	0	U	

• Molecule 1 is a protein called Nickel (III) ABC transporter, periplasmic iron-bindin gprotein.

Chain	Residue	Modelled	Actual	Comment	Reference
А	335	LYS	-	expression tag	UNP B5Z9J2
А	336	GLY	-	expression tag	UNP B5Z9J2
А	337	GLU	-	expression tag	UNP B5Z9J2
А	338	LEU	-	expression tag	UNP B5Z9J2
А	339	ASN	-	expression tag	UNP B5Z9J2
А	340	SER	-	expression tag	UNP B5Z9J2
А	341	LYS	-	expression tag	UNP B5Z9J2
А	342	LEU	-	expression tag	UNP B5Z9J2
А	343	GLU	-	expression tag	UNP B5Z9J2
A	344	GLY	-	expression tag	UNP B5Z9J2
A	345	LYS	-	expression tag	UNP B5Z9J2
A	346	PRO	-	expression tag	UNP B5Z9J2
A	347	ILE	-	expression tag	UNP B5Z9J2
А	348	PRO	-	expression tag	UNP B5Z9J2
A	349	ASN	-	expression tag	UNP B5Z9J2
А	350	PRO	-	expression tag	UNP B5Z9J2
А	351	LEU	-	expression tag	UNP B5Z9J2
А	352	LEU	-	expression tag	UNP B5Z9J2
А	353	GLY	-	expression tag	UNP B5Z9J2
A	354	LEU	-	expression tag	UNP B5Z9J2
А	355	ASP	-	expression tag	UNP B5Z9J2
А	356	SER	-	expression tag	UNP B5Z9J2
А	357	THR	-	expression tag	UNP B5Z9J2
А	358	ARG	-	expression tag	UNP B5Z9J2
А	359	THR	-	expression tag	UNP B5Z9J2

There are 64 discrepancies between the modelled and reference sequences:




Continued from previous page						
Chain	Residue	Modelled	Actual	Comment	Reference	
A	360	GLY	-	expression tag	UNP B5Z9J2	
A	361	HIS	-	expression tag	UNP B5Z9J2	
A	362	HIS	-	expression tag	UNP B5Z9J2	
A	363	HIS	-	expression tag	UNP B5Z9J2	
А	364	HIS	-	expression tag	UNP B5Z9J2	
A	365	HIS	-	expression tag	UNP B5Z9J2	
A	366	HIS	-	expression tag	UNP B5Z9J2	
В	335	LYS	-	expression tag	UNP B5Z9J2	
В	336	GLY	-	expression tag	UNP B5Z9J2	
В	337	GLU	-	expression tag	UNP B5Z9J2	
В	338	LEU	-	expression tag	UNP B5Z9J2	
В	339	ASN	-	expression tag	UNP B5Z9J2	
В	340	SER	-	expression tag	UNP B5Z9J2	
В	341	LYS	-	expression tag	UNP B5Z9J2	
В	342	LEU	-	expression tag	UNP B5Z9J2	
В	343	GLU	-	expression tag	UNP B5Z9J2	
В	344	GLY	-	expression tag	UNP B5Z9J2	
В	345	LYS	-	expression tag	UNP B5Z9J2	
В	346	PRO	-	expression tag	UNP B5Z9J2	
В	347	ILE	-	expression tag	UNP B5Z9J2	
В	348	PRO	-	expression tag	UNP B5Z9J2	
В	349	ASN	-	expression tag	UNP B5Z9J2	
В	350	PRO	-	expression tag	UNP B5Z9J2	
В	351	LEU	-	expression tag	UNP B5Z9J2	
В	352	LEU	-	expression tag	UNP B5Z9J2	
В	353	GLY	-	expression tag	UNP B5Z9J2	
В	354	LEU	-	expression tag	UNP B5Z9J2	
В	355	ASP	-	expression tag	UNP B5Z9J2	
В	356	SER	-	expression tag	UNP B5Z9J2	
В	357	THR	-	expression tag	UNP B5Z9J2	
В	358	ARG	-	expression tag	UNP B5Z9J2	
В	359	THR	-	expression tag	UNP B5Z9J2	
В	360	GLY	-	expression tag	UNP B5Z9J2	
В	361	HIS	-	expression tag	UNP B5Z9J2	
В	362	HIS	-	expression tag	UNP B5Z9J2	
В	363	HIS	-	expression tag	UNP B5Z9J2	
В	364	HIS	-	expression tag	UNP B5Z9J2	
В	365	HIS	-	expression tag	UNP B5Z9J2	
В	366	HIS	-	expression tag	UNP B5Z9J2	

Continued from previous page...

• Molecule 2 is HISTIDINE (three-letter code: HIS) (formula:  $C_6H_{10}N_3O_2$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total         C         N         O           11         6         3         2	0	0
2	А	1	Total         C         N         O           11         6         3         2	0	0
2	В	1	Total         C         N         O           11         6         3         2	0	0
2	В	1	Total         C         N         O           11         6         3         2	0	0

• Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Ni 1 1	0	0
3	В	1	Total Ni 1 1	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	510	Total O 510 510	0	0
4	В	484	Total         O           484         484	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nickel (III) ABC transporter, periplasmic iron-bindin gprotein





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants	71.42Å $60.65$ Å $77.13$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $91.02^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	30.32 - 1.70	Depositor
Resolution (A)	30.33 - 1.70	EDS
% Data completeness	$96.5(30.32  ext{-} 1.70)$	Depositor
(in resolution range)	91.6 (30.33 - 1.70)	EDS
R <sub>merge</sub>	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.37 (at 1.70 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.168 , $0.204$	Depositor
$n, n_{free}$	0.175 , $0.206$	DCC
$R_{free}$ test set	3554 reflections $(5.07%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	15.3	Xtriage
Anisotropy	0.381	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.33, 29.8	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.387 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5838	wwPDB-VP
Average B, all atoms $(Å^2)$	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.13% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.37	0/2452	0.56	1/3309~(0.0%)	
1	В	0.37	0/2452	0.57	0/3309	
All	All	0.37	0/4904	0.57	1/6618~(0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^{o})$	$Ideal(^{o})$
1	А	$\overline{72}$	ARG	NE-CZ-NH2	-5.24	117.68	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2399	0	2484	9	0
1	В	2399	0	2484	13	0
2	А	22	0	12	0	0
2	В	22	0	12	1	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
4	А	510	0	0	5	1
4	В	484	0	0	4	1

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5838	0	4992	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:LYS:NZ	4:A:1152:HOH:O	2.19	0.76
1:A:71:ASP:HB3	4:A:1080:HOH:O	1.90	0.72
1:B:335:LYS:NZ	4:B:1149:HOH:O	2.25	0.69
1:B:37:LYS:HG2	1:B:46:THR:HG22	1.74	0.68
1:B:105:ALA:HB1	1:B:106:ALA:HA	1.80	0.63

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
4:A:1103:HOH:O	4:B:1069:HOH:O[2_656]	2.04	0.16	

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Favoured Allowed		Perce	ntiles
1	А	300/333~(90%)	292~(97%)	8(3%)	0	100	100
1	В	300/333~(90%)	292~(97%)	8(3%)	0	100	100
All	All	600/666~(90%)	584 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	262/289~(91%)	259~(99%)	3(1%)	73 63		
1	В	262/289~(91%)	259~(99%)	3 (1%)	73 63		
All	All	524/578~(91%)	518~(99%)	6 (1%)	65 63		

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	112	LEU
1	В	138	ILE
1	В	185	LYS
1	А	112	LEU
1	А	72	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Dec	Res Link	Bond lengths			Bond angles		
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
2	HIS	А	602	3	6,11,11	0.89	0	7,14,14	1.44	2 (28%)
2	HIS	А	601	3	6,11,11	1.00	1 (16%)	7,14,14	1.35	2 (28%)
2	HIS	В	601	3	6,11,11	0.97	1 (16%)	7,14,14	1.59	2 (28%)
2	HIS	В	602	3	6,11,11	0.94	1 (16%)	7,14,14	1.53	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HIS	А	602	3	-	1/8/8/8	0/1/1/1
2	HIS	А	601	3	-	3/8/8/8	0/1/1/1
2	HIS	В	601	3	-	3/8/8/8	0/1/1/1
2	HIS	В	602	3	-	0/8/8/8	0/1/1/1

All $(3)$ bond	length	outliers	$\operatorname{are}$	listed	below:
----------------	--------	----------	----------------------	--------	--------

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	А	601	HIS	OXT-C	-2.20	1.23	1.30
2	В	601	HIS	OXT-C	-2.18	1.23	1.30
2	В	602	HIS	OXT-C	-2.04	1.23	1.30

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	601	HIS	OXT-C-O	-2.97	117.35	124.09
2	В	602	HIS	OXT-C-O	-2.48	118.46	124.09
2	А	602	HIS	OXT-C-O	-2.29	118.88	124.09
2	В	601	HIS	OXT-C-CA	2.27	121.13	113.38
2	В	602	HIS	OXT-C-CA	2.16	120.75	113.38

There are no chirality outliers.



Mol	Chain	Res	Type	Atoms
2	А	601	HIS	CA-CB-CG-ND1
2	А	601	HIS	CA-CB-CG-CD2
2	В	601	HIS	CA-CB-CG-ND1
2	В	601	HIS	CA-CB-CG-CD2
2	А	602	HIS	C-CA-CB-CG

5 of 7 torsion outliers are listed below:

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	601	HIS	1	0

### 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\begin{tabular}{lllllllllllllllllllllllllllllllllll$		$OWAB(Å^2)$	Q<0.9
1	А	302/333~(90%)	-0.49	2 (0%) 87 90	10, 16, 30, 43	0
1	В	302/333~(90%)	-0.44	2 (0%) 87 90	9, 16, 30, 41	0
All	All	604/666~(90%)	-0.46	4 (0%) 87 90	9, 16, 31, 43	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	105	ALA	4.9
1	В	138	ILE	2.8
1	А	138	ILE	2.6
1	А	133	ALA	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
2	HIS	А	602	11/11	0.93	0.09	$10,\!14,\!17,\!18$	0

Continued on next page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	HIS	В	602	11/11	0.94	0.08	$10,\!14,\!18,\!19$	0
2	HIS	В	601	11/11	0.97	0.07	11,12,14,15	0
2	HIS	А	601	11/11	0.98	0.06	$10,\!13,\!15,\!15$	0
3	NI	А	603	1/1	1.00	0.02	13,13,13,13	0
3	NI	В	603	1/1	1.00	0.02	14,14,14,14	0

Continued from previous page...

## 6.5 Other polymers (i)

There are no such residues in this entry.

