

# wwPDB X-ray Structure Validation Summary Report (i)

#### Jan 6, 2024 – 12:06 pm GMT

PDB ID	:	5LS6
Title	:	Structure of Human Polycomb Repressive Complex 2 (PRC2) with inhibitor
Authors	:	Zhang, Y.; Justin, N.; Chen, S.; Wilson, J.; Gamblin, S.
Deposited on	:	2016-08-22
Resolution	:	3.47  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 3.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	130704	1379 (3.56-3.40)
Clashscore	141614	1461 (3.56-3.40)
Ramachandran outliers	138981	1424 (3.56-3.40)
Sidechain outliers	138945	1425 (3.56-3.40)
RSRZ outliers	127900	1289 (3.56-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain							
1	А	695	69%	12%	·	18%				
1	D	695	66% 1	.4%		18%	_			
1	G	695	68%	13%	•	18%	_			
1	J	695	70%	11%	·	18%				
2	В	367	4%			19%	••			



Mol	Chain	Length		Quality of chain						
2	Е	367	.%	8%	20% ••					
2	Н	367	2% 	86%	13% •					
2	K	367	% •	84%	14% ••					
3	С	129	6% 71%		21% 5% •					
3	F	129	72%	,	25% •					
3	Ι	129	4% 75	%	22% •					
3	L	129	7	8%	18% ••					
4	Q	11	36%	45%	9% 9%					
4	R	11	45%	36%	9% 9%					
4	S	11	45%	45%	9%					
4	Т	11	45%	27%	18% 9%					

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ZN	А	801	-	-	Х	-
5	ZN	G	806	-	-	Х	-
5	ZN	J	806	-	-	Х	-
6	74D	А	809	Х	-	-	-
6	74D	D	809	Х	-	-	-
6	74D	G	809	Х	-	-	-
6	74D	J	809	Х	-	-	-



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 34891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Histone-lysine N-methyltransferase EZH2, Histone-lysine N-methyltransferase EZH2, Histone-lysine N-methyltransferase EZH2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	570	Total	С	Ν	0	$\mathbf{S}$	0	0	0
1	A	570	4588	2878	813	855	42		0	0
1	Л	570	Total	С	Ν	0	S	0	0	0
	D	570	4588	2878	813	855	42			0
1	С	572	Total	С	Ν	0	S	0	0	0
	I G		4603	2886	816	859	42			0
1	1 J	570	Total	С	Ν	0	S	0	0	0
			4588	2878	813	855	42	0		U

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-9	GLU	-	expression tag	UNP Q15910
А	-8	THR	-	expression tag	UNP Q15910
А	-7	SER	-	expression tag	UNP Q15910
А	-6	LEU	-	expression tag	UNP Q15910
А	-5	ALA	-	expression tag	UNP Q15910
А	-4	GLU	-	expression tag	UNP Q15910
А	-3	GLU	-	expression tag	UNP Q15910
А	-2	LYS	-	expression tag	UNP Q15910
А	-1	LEU	-	expression tag	UNP Q15910
А	0	THR	-	expression tag	UNP Q15910
А	249	GLN	GLU	$\operatorname{conflict}$	UNP Q15910
А	419	PRO	-	linker	UNP Q15910
А	420	GLY	-	linker	UNP Q15910
D	-9	GLU	-	expression tag	UNP Q15910
D	-8	THR	-	expression tag	UNP Q15910
D	-7	SER	-	expression tag	UNP Q15910
D	-6	LEU	_	expression tag	UNP Q15910
D	-5	ALA	-	expression tag	UNP Q15910
D	-4	GLU	-	expression tag	UNP Q15910
D	-3	GLU	-	expression tag	UNP Q15910



Chain	<b>Besidue</b>	Modelled	Actual	Comment	Reference
D	_2	LVS	-	expression tag	UNP 015010
	1	LEU	_	expression tag	$\frac{000}{100}$
	-1	THR	-	expression tag	$\frac{000}{1000}$
	240		CLU	expression tag	UNP Q15910
	410	BBO	GLU	linkor	$\frac{000}{1000}$
	419		-	linker	$\frac{\text{UND}}{\text{UND}} \frac{015910}{015010}$
	420	GLI	-		$\frac{\text{UNF Q15910}}{\text{UND Q15910}}$
G	-9	GLU	-	expression tag	UNP Q15910
G	-8		-	expression tag	UNP Q15910
G	-7	SER	-	expression tag	UNP Q15910
G	-6	LEU	-	expression tag	UNP Q15910
G	-5	ALA	-	expression tag	UNP Q15910
G	-4	GLU	-	expression tag	UNP Q15910
G	-3	GLU	-	expression tag	UNP Q15910
G	-2	LYS	-	expression tag	UNP Q15910
G	-1	LEU	-	expression tag	UNP Q15910
G	0	THR	-	expression tag	UNP Q15910
G	249	GLN	GLU	conflict	UNP Q15910
G	419	PRO	-	linker	UNP Q15910
G	420	GLY	-	linker	UNP Q15910
J	-9	GLU	-	expression tag	UNP Q15910
J	-8	THR	-	expression tag	UNP Q15910
J	-7	SER	-	expression tag	UNP Q15910
J	-6	LEU	-	expression tag	UNP Q15910
J	-5	ALA	-	expression tag	UNP Q15910
J	-4	GLU	-	expression tag	UNP Q15910
J	-3	GLU	-	expression tag	UNP Q15910
J	-2	LYS	-	expression tag	UNP Q15910
J	-1	LEU	-	expression tag	UNP Q15910
J	0	THR	-	expression tag	UNP Q15910
J	249	GLN	GLU	conflict	UNP Q15910
J	419	PRO	-	linker	UNP Q15910
J	420	GLY	-	linker	UNP Q15910

• Molecule 2 is a protein called Polycomb protein EED.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
0	D	265	Total	С	Ν	0	S	0	0	0
	D		2959	1873	521	543	22	0	0	0
0	Б	265	Total	С	Ν	0	$\mathbf{S}$	0	0	0
	Ľ	305	2959	1873	521	543	22	0	0	
0	п	265	Total	С	Ν	0	$\mathbf{S}$	0	0	0
	п	909	2959	1873	521	543	22			0



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	K	365	Total 2959	C 1873	N 521	0 543	S 22	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	75	GLY	-	expression tag	UNP 075530
В	76	SER	-	expression tag	UNP 075530
Е	75	GLY	-	expression tag	UNP 075530
Е	76	SER	-	expression tag	UNP 075530
Н	75	GLY	-	expression tag	UNP 075530
Н	76	SER	-	expression tag	UNP 075530
K	75	GLY	-	expression tag	UNP 075530
K	76	SER	-	expression tag	UNP 075530

• Molecule 3 is a protein called Polycomb protein SUZ12.

Mol	Chain	Residues		Atoms					AltConf	Trace
3 C	125	Total	С	Ν	0	$\mathbf{S}$	0	0	0	
		1042	657	180	193	12	0	0	0	
2	Б	195	Total	С	Ν	0	S	0	0	0
J	ЭГ	120	1042	657	180	193	12	0		
2	т	195	Total	С	Ν	0	S	0	0	0
0 1	120	1042	657	180	193	12	0	0	0	
9	9 I	195	Total	С	Ν	0	S	0	0	0
3 L	125	1042	657	180	193	12	0	0	U	

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	557	GLY	-	expression tag	UNP Q15022
F	557	GLY	-	expression tag	UNP Q15022
Ι	557	GLY	-	expression tag	UNP Q15022
L	557	GLY	-	expression tag	UNP Q15022

• Molecule 4 is a protein called Jarid2 K116me3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4 Q	10	Total	С	Ν	0	0	0	0	
	Q	10	85	54	17	14	0	0	0
4	4 D	10	Total	С	Ν	0	0	0	0
4	n	10	85	54	17	14			0



001000	iraca jien	v precioac pa	.ge						
Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4 S	10	Total	С	Ν	0	0	0	0	
	G	10	85	54	17	14	0		0
4 T	10	Total	С	Ν	0	0	0	0	
		10	85	54	17	14	0	0	U

Continued from previous page...

• Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	8	Total Zn 8 8	0	0
5	D	8	Total Zn 8 8	0	0
5	G	8	Total Zn 8 8	0	0
5	J	8	Total Zn 8 8	0	0

• Molecule 6 is 1-[(1 {R})-1-[1-[2,2-bis(fluoranyl)propyl]piperidin-4-yl]ethyl]- {N}-[(4-metho xy-6-methyl-2-oxidanylidene-3 {H}-pyridin-3-yl)methyl]-2-methyl-indole-3-carboxamide (three-letter code: 74D) (formula:  $C_{28}H_{36}F_2N_4O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
6	Λ	1	Total	С	F	Ν	Ο	0	0
0 A	I	37	28	2	4	3	0	0	
6	6 D	1	Total	С	F	Ν	Ο	0	0
0		L	37	28	2	4	3	0	



001000	e onomaca from process as pagem										
Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
6	С	1	Total	С	F	Ν	Ο	0	0		
	G	1	37	28	2	4	3	0	0		
6	J	1	Total	С	F	Ν	Ο	0	0		
			37	28	2	4	3	0	0		

Continued from previous page...



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

 $\bullet$  Molecule 1: Histone-lysine N-methyltransferase EZH2,<br/>Histone-lysine N-methyltransferase EZH2, Histone-lysine N-methyltransferase EZH2



 $\bullet$  Molecule 1: Histone-lysine N-methyltransferase EZH2,<br/>Histone-lysine N-methyltransferase EZH2, Histone-lysine N-methyltransferase EZH2







• Molecule 1: Histone-lysine N-methyltransferase EZH2, Histone-lysine N-methyltransferase EZH2, Histone-lysine N-methyltransferase EZH2



 $\bullet$  Molecule 1: Histone-lysine N-methyltransferase EZH2,<br/>Histone-lysine N-methyltransferase EZH2, Histone-lysine N-methyltransferase EZH2





# 1403 1224 1403 1224 1404 1224 1405 1226 1404 1226 141 1226 1434 1226 1434 1226 1434 1226 1434 1226 1434 1226 1434 1226 1434 1257 1434 1256 1434 1256 1431 1257 1431 1256 1431 1256 1431 1254 1254 1256 1319 <tr

• Molecule 2: Polycomb protein EED



- Molecule 2: Polycomb protein EED



# N307 1318 1318 1318 1318 1326 1331 1331 1332 1333 1333 1333 1333 1333 1333 1333 1333 1333 1333 1335

 $\bullet$  Molecule 3: Polycomb protein SUZ12



#### 1671 1674 1681

• Molecule 3: Polycomb protein SUZ12



• Molecule 3: Polycomb protein SUZ12









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	131.31Å 170.27Å 275.54Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution(A)	68.84 - 3.47	Depositor
Resolution (A)	68.84 - 3.47	EDS
% Data completeness	99.9 (68.84-3.47)	Depositor
(in resolution range)	99.9(68.84 - 3.47)	EDS
R <sub>merge</sub>	0.30	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.92 (at 3.49 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
P. P.	0.229 , $0.299$	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.231 , $0.297$	DCC
$R_{free}$ test set	3952 reflections $(4.91%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	78.0	Xtriage
Anisotropy	0.722	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.31, 56.0	EDS
L-test for $twinning^2$	$ < L >=0.42, < L^2>=0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	34891	wwPDB-VP
Average B, all atoms $(Å^2)$	105.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 34.62 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.6281e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 74D, ZN, M3L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bo	ond angles
	Ullaili	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.50	0/4691	0.74	0/6327
1	D	0.51	0/4691	0.74	0/6327
1	G	0.51	0/4706	0.76	2/6347~(0.0%)
1	J	0.53	1/4691~(0.0%)	0.75	0/6327
2	В	0.52	0/3034	0.76	0/4107
2	Е	0.50	0/3034	0.76	0/4107
2	Н	0.52	0/3034	0.78	0/4107
2	Κ	0.50	0/3034	0.77	0/4107
3	С	0.51	0/1063	0.79	2/1427~(0.1%)
3	F	0.49	0/1063	0.77	0/1427
3	Ι	0.46	0/1063	0.77	0/1427
3	L	0.50	0/1063	0.77	0/1427
4	Q	0.65	0/73	1.05	0/96
4	R	0.70	0/73	0.73	0/96
4	S	0.66	0/73	1.02	0/96
4	Т	0.75	0/73	1.10	0/96
All	All	0.51	1/35459~(0.0%)	0.76	4/47848~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	3
1	D	0	3
1	G	0	2
1	J	0	2
3	С	0	3
3	F	0	1
3	L	0	2



Mol	Chain	#Chirality outliers	#Planarity outliers
All	All	0	16

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
1	J	182	ASN	C-O	5.38	1.33	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	G	685	ARG	NE-CZ-NH2	5.97	123.29	120.30
1	G	315	LEU	CA-CB-CG	5.97	129.02	115.30
3	С	563	ARG	NE-CZ-NH1	5.20	122.90	120.30
3	С	608	GLU	N-CA-C	5.11	124.80	111.00

There are no chirality outliers.

5 of 16 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	134	MET	Peptide
1	А	311	THR	Peptide
1	А	326	GLN	Peptide
3	С	561	HIS	Peptide
3	С	579	MET	Peptide

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4588	0	4445	74	0
1	D	4588	0	4442	55	0
1	G	4603	0	4457	51	0
1	J	4588	0	4447	37	0
2	В	2959	0	2881	49	0
2	Е	2959	0	2881	44	0
2	Н	2959	0	2881	30	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	K	2959	0	2881	34	0
3	С	1042	0	1021	22	0
3	F	1042	0	1021	20	0
3	Ι	1042	0	1021	17	0
3	L	1042	0	1021	17	0
4	Q	85	0	90	6	0
4	R	85	0	90	3	0
4	S	85	0	90	5	0
4	Т	85	0	90	5	0
5	А	8	0	0	2	0
5	D	8	0	0	0	0
5	G	8	0	0	3	0
5	J	8	0	0	2	0
6	А	37	0	0	4	0
6	D	37	0	0	0	0
6	G	37	0	0	0	0
6	J	37	0	0	0	0
All	All	34891	0	33759	401	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 401 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:667:PHE:CD2	1:A:708:ILE:HD12	1.39	1.55
1:A:667:PHE:CD2	1:A:708:ILE:CD1	2.00	1.41
1:A:667:PHE:CE2	1:A:708:ILE:CD1	2.07	1.36
1:A:667:PHE:CE2	1:A:708:ILE:HD13	1.72	1.22
1:A:667:PHE:CE2	1:A:708:ILE:HD12	1.76	1.12

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



5LS6
------

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	558/695~(80%)	501 (90%)	51 (9%)	6 (1%)	14 50
1	D	558/695~(80%)	504 (90%)	44 (8%)	10~(2%)	8 38
1	G	560/695~(81%)	513 (92%)	41 (7%)	6(1%)	14 50
1	J	558/695~(80%)	501 (90%)	50 (9%)	7~(1%)	12 45
2	В	363/367~(99%)	330 (91%)	31 (8%)	2(1%)	25 63
2	Ε	363/367~(99%)	331 (91%)	27 (7%)	5(1%)	11 43
2	Η	363/367~(99%)	346~(95%)	16 (4%)	1 (0%)	41 75
2	Κ	363/367~(99%)	334 (92%)	27 (7%)	2(1%)	25 63
3	С	123/129~(95%)	113~(92%)	8~(6%)	2(2%)	9 41
3	F	123/129~(95%)	112 (91%)	10 (8%)	1 (1%)	19 57
3	Ι	123/129~(95%)	118 (96%)	4(3%)	1 (1%)	19 57
3	L	123/129~(95%)	113~(92%)	10 (8%)	0	100 100
4	Q	7/11~(64%)	6 (86%)	0	1 (14%)	0 2
4	R	7/11~(64%)	7~(100%)	0	0	100 100
4	S	7/11~(64%)	7 (100%)	0	0	100 100
4	Т	7/11~(64%)	4 (57%)	2(29%)	1 (14%)	0 2
All	All	4206/4808 (88%)	3840 (91%)	321 (8%)	45 (1%)	14 50

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	68	VAL
1	А	103	ALA
2	В	257	ASP
1	D	103	ALA
1	J	103	ALA

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	А	511/618~(83%)	486 (95%)	25~(5%)	25	57
1	D	511/618~(83%)	482 (94%)	29~(6%)	20	52
1	G	513/618~(83%)	482 (94%)	31 (6%)	19	51
1	J	511/618 (83%)	482 (94%)	29 (6%)	20	52
2	В	328/329~(100%)	316 (96%)	12 (4%)	34	64
2	Е	328/329~(100%)	316 (96%)	12 (4%)	34	64
2	Н	328/329~(100%)	318 (97%)	10 (3%)	41	70
2	Κ	328/329~(100%)	321 (98%)	7 (2%)	53	78
3	С	119/121 (98%)	113 (95%)	6 (5%)	24	56
3	F	119/121 (98%)	116 (98%)	3 (2%)	47	74
3	Ι	119/121 (98%)	115 (97%)	4 (3%)	37	67
3	L	119/121~(98%)	118 (99%)	1 (1%)	81	92
4	Q	7/8~(88%)	5 (71%)	2 (29%)	0	2
4	R	7/8~(88%)	5 (71%)	2(29%)	0	2
4	S	7/8~(88%)	6 (86%)	1 (14%)	3	17
4	Т	7/8~(88%)	5 (71%)	2 (29%)	0	2
All	All	3862/4304~(90%)	3686 (95%)	176 (5%)	27	59

5 of 176 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	G	727	ARG
1	J	134	MET
2	Н	94	GLN
3	Ι	642	ASN
1	J	311	THR

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such side chains are listed below:

Mol	Chain	Res	Type
3	F	655	ASN
3	L	642	ASN
2	Н	176	ASN
3	L	633	ASN
4	R	3	GLN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain		Pog Link	Bo	Bond lengths			Bond angles		
	туре	Unain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
4	M3L	S	7	4	10,11,12	0.63	0	9,14,16	0.61	0	
4	M3L	Q	7	4	10,11,12	0.65	0	9,14,16	0.55	0	
4	M3L	Т	7	4	10,11,12	0.55	0	9,14,16	0.48	0	
4	M3L	R	7	4	10,11,12	0.83	0	9,14,16	0.56	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	M3L	S	7	4	-	1/9/10/12	-
4	M3L	Q	7	4	-	0/9/10/12	-
4	M3L	Т	7	4	-	0/9/10/12	-
4	M3L	R	7	4	-	1/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	S	7	M3L	O-C-CA-CB
4	R	7	M3L	N-CA-CB-CG



There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	S	7	M3L	1	0
4	Q	7	M3L	1	0
4	Т	7	M3L	2	0
4	R	7	M3L	1	0

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 36 ligands modelled in this entry, 32 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Timle	B	Bond lengths			Bond angles		
IVIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2	
6	74D	D	809	-	32,40,40	3.11	7 (21%)	41,59,59	2.64	11 (26%)	
6	74D	G	809	-	32,40,40	2.82	7 (21%)	41,59,59	2.83	16 (39%)	
6	74D	J	809	-	32,40,40	<mark>3.13</mark>	9 (28%)	41,59,59	2.74	14 (34%)	
6	74D	А	809	-	32,40,40	<mark>3.34</mark>	10 (31%)	41,59,59	2.18	8 (19%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	74D	D	809	-	1/1/9/11	6/17/50/50	0/4/4/4
6	74D	G	809	-	1/1/9/11	11/17/50/50	0/4/4/4
6	74D	J	809	-	1/1/9/11	9/17/50/50	0/4/4/4
6	74D	А	809	-	1/1/9/11	9/17/50/50	0/4/4/4



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	А	809	74D	CAN-CAM	-12.72	1.33	1.51
6	J	809	74D	CAN-CAM	-11.44	1.34	1.51
6	D	809	74D	CAN-CAM	-10.37	1.36	1.51
6	G	809	74D	CAN-CAM	-9.59	1.37	1.51
6	А	809	74D	CAC-CAB	-9.45	1.39	1.50

The worst 5 of 33 bond length outliers are listed below:

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
6	D	809	74D	CAK-CAC-CAD	12.90	123.11	108.89
6	G	809	74D	CAK-CAC-CAD	11.94	122.05	108.89
6	J	809	74D	CAK-CAC-CAD	11.94	122.05	108.89
6	А	809	74D	CAK-CAC-CAD	9.95	119.85	108.89
6	J	809	74D	OAI-CAB-CAA	-6.71	119.94	126.00

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	А	809	74D	CAC
6	D	809	74D	CAC
6	G	809	74D	CAC
6	J	809	74D	CAC

5 of 35 torsion outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	Atoms
6	А	809	74D	NBD-CBG-CBH-FBI
6	А	809	74D	CAZ-CAY-NAQ-CAP
6	А	809	74D	CAZ-CAY-NAQ-CAR
6	А	809	74D	CAC-CAB-OAI-CAJ
6	А	809	74D	CAA-CAB-OAI-CAJ

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	А	809	74D	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will



also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and similar rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







#### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	А	1
1	G	1
1	D	1



Mol	Chain	Number of breaks
1	J	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	182:ASN	С	211:GLU	N	6.31
1	G	182:ASN	С	211:GLU	N	5.74
1	D	182:ASN	С	211:GLU	N	5.71
1	J	182:ASN	С	211:GLU	N	5.36



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	570/695~(82%)	0.86	74 (12%) 3 5	51, 108, 183, 239	0
1	D	570/695~(82%)	0.55	49 (8%) 10 12	56, 100, 177, 259	0
1	G	572/695~(82%)	0.81	83 (14%) 2 3	49, 107, 192, 252	0
1	J	570/695~(82%)	0.95	93 (16%) 1 2	52, 108, 203, 254	0
2	В	365/367~(99%)	0.33	16 (4%) 34 32	54, 94, 139, 213	0
2	E	365/367~(99%)	0.24	3 (0%) 86 82	57, 93, 137, 198	0
2	Н	365/367~(99%)	0.17	7 (1%) 66 63	49, 80, 120, 216	0
2	K	365/367~(99%)	0.13	3 (0%) 86 82	46, 74, 118, 208	0
3	С	125/129~(96%)	0.55	8 (6%) 19 19	67, 100, 168, 233	0
3	F	125/129~(96%)	0.36	6 (4%) 30 29	58, 92, 161, 219	0
3	Ι	125/129~(96%)	0.43	5 (4%) 38 35	73, 104, 163, 202	0
3	L	125/129~(96%)	0.67	14 (11%) 5 7	69, 113, 163, 229	0
4	Q	9/11~(81%)	0.27	0 100 100	77, 101, 126, 131	0
4	R	9/11~(81%)	0.61	1 (11%) 5 7	77, 95, 135, 138	0
4	S	9/11 (81%)	0.81	1 (11%) 5 7	93, 119, 137, 139	0
4	Т	9/11~(81%)	1.20	2(22%) 0 1	101, 117, 159, 165	0
All	All	4278/4808 (88%)	0.56	365 (8%) 10 13	46, 95, 181, 259	0

The worst 5 of 365 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	479	PRO	10.4
1	J	480	ALA	9.7
1	J	429	ASN	9.5
1	G	480	ALA	8.8
1	А	75	SER	8.8



#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
4	M3L	R	7	12/13	0.96	0.25	$50,\!57,\!84,\!84$	0
4	M3L	S	7	12/13	0.96	0.27	60,71,87,89	0
4	M3L	Т	7	12/13	0.96	0.25	67,74,83,87	0
4	M3L	Q	7	12/13	0.97	0.23	55,63,83,84	0

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
5	ZN	J	802	1/1	0.94	0.10	147,147,147,147	0
6	74D	А	809	37/37	0.94	0.29	62,78,86,91	0
6	74D	D	809	37/37	0.95	0.30	61,70,89,93	0
6	74D	J	809	37/37	0.95	0.27	65,77,87,89	0
6	74D	G	809	37/37	0.96	0.31	61,69,76,77	0
5	ZN	А	802	1/1	0.96	0.11	132,132,132,132	0
5	ZN	G	802	1/1	0.98	0.12	146,146,146,146	0
5	ZN	J	807	1/1	0.98	0.18	71,71,71,71	0
5	ZN	G	803	1/1	0.98	0.13	73,73,73,73	0
5	ZN	D	806	1/1	0.99	0.15	58, 58, 58, 58	0
5	ZN	D	807	1/1	0.99	0.17	66,66,66,66	0
5	ZN	D	808	1/1	0.99	0.17	56, 56, 56, 56	0
5	ZN	G	801	1/1	0.99	0.14	71,71,71,71	0
5	ZN	А	801	1/1	0.99	0.13	70,70,70,70	0
5	ZN	А	804	1/1	0.99	0.14	56, 56, 56, 56	0
5	ZN	G	804	1/1	0.99	0.14	70,70,70,70	0
5	ZN	G	805	1/1	0.99	0.13	72,72,72,72	0
5	ZN	G	806	1/1	0.99	0.16	61,61,61,61	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
5	ZN	G	807	1/1	0.99	0.17	59, 59, 59, 59, 59	0
5	ZN	J	801	1/1	0.99	0.15	68,68,68,68	0
5	ZN	А	805	1/1	0.99	0.15	66,66,66,66	0
5	ZN	J	803	1/1	0.99	0.12	89,89,89,89	0
5	ZN	J	804	1/1	0.99	0.13	94,94,94,94	0
5	ZN	J	805	1/1	0.99	0.11	80,80,80,80	0
5	ZN	J	806	1/1	0.99	0.17	$75,\!75,\!75,\!75$	0
5	ZN	А	806	1/1	0.99	0.13	$53,\!53,\!53,\!53$	0
5	ZN	А	807	1/1	0.99	0.15	$55,\!55,\!55,\!55$	0
5	ZN	D	801	1/1	0.99	0.15	$68,\!68,\!68,\!68$	0
5	ZN	D	802	1/1	0.99	0.11	84,84,84,84	0
5	ZN	D	805	1/1	0.99	0.13	$65,\!65,\!65,\!65$	0
5	ZN	А	803	1/1	1.00	0.13	$65,\!65,\!65,\!65$	0
5	ZN	J	808	1/1	1.00	0.16	$49,\!49,\!49,\!49$	0
5	ZN	А	808	1/1	1.00	0.16	$55,\!55,\!55,\!55$	0
5	ZN	D	803	1/1	1.00	0.13	59, 59, 59, 59	0
5	ZN	G	808	1/1	1.00	0.17	49,49,49,49	0
5	ZN	D	804	1/1	1.00	0.13	61,61,61,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











## 6.5 Other polymers (i)

There are no such residues in this entry.

