



Full wwPDB NMR Structure Validation Report ⓘ

May 29, 2020 – 05:35 am BST

PDB ID : 2M61
Title : NMR and Mass Spectrometric Studies of M-2 Branch Mini-M Conotoxins from Indian Cone Snails
Authors : Sarma, S.P.; Rajesh, R.P.; Kumar, G.S.; Sudarslal, S.; Sabareesh, V.; Gowd, K.H.; Gupta, K.; Krishnan, K.S.; Balaram, P.
Deposited on : 2013-03-18

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 68%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	15	 73% 7% 7% 13%

2 Ensemble composition and analysis

This entry contains 30 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:11, A:13-A:14 (13)	0.00	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: No clusters in NmrClust output

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 181 atoms, of which 86 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Conotoxin Ar1446.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	15	181	54	86	20	15	6	1

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	15	NH2	-	AMIDATION	UNP P0CI24

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Conotoxin Ar1446




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Conotoxin Ar1446




4.2.2 Score per residue for model 2

- Molecule 1: Conotoxin Ar1446




4.2.3 Score per residue for model 3

- Molecule 1: Conotoxin Ar1446




4.2.4 Score per residue for model 4

- Molecule 1: Conotoxin Ar1446




4.2.5 Score per residue for model 5 (medoid)

- Molecule 1: Conotoxin Ar1446




4.2.6 Score per residue for model 6

- Molecule 1: Conotoxin Ar1446




4.2.7 Score per residue for model 7

- Molecule 1: Conotoxin Ar1446




4.2.8 Score per residue for model 8

- Molecule 1: Conotoxin Ar1446




4.2.9 Score per residue for model 9

- Molecule 1: Conotoxin Ar1446




4.2.10 Score per residue for model 10

- Molecule 1: Conotoxin Ar1446




4.2.11 Score per residue for model 11

- Molecule 1: Conotoxin Ar1446




4.2.12 Score per residue for model 12

- Molecule 1: Conotoxin Ar1446




4.2.13 Score per residue for model 13

- Molecule 1: Conotoxin Ar1446




4.2.14 Score per residue for model 14

- Molecule 1: Conotoxin Ar1446




4.2.15 Score per residue for model 15

- Molecule 1: Conotoxin Ar1446




4.2.16 Score per residue for model 16

- Molecule 1: Conotoxin Ar1446




4.2.17 Score per residue for model 17

- Molecule 1: Conotoxin Ar1446




4.2.18 Score per residue for model 18

- Molecule 1: Conotoxin Ar1446




4.2.19 Score per residue for model 19

- Molecule 1: Conotoxin Ar1446




4.2.20 Score per residue for model 20

- Molecule 1: Conotoxin Ar1446




4.2.21 Score per residue for model 21

- Molecule 1: Conotoxin Ar1446




4.2.22 Score per residue for model 22

- Molecule 1: Conotoxin Ar1446




4.2.23 Score per residue for model 23

- Molecule 1: Conotoxin Ar1446




4.2.24 Score per residue for model 24

- Molecule 1: Conotoxin Ar1446




4.2.25 Score per residue for model 25

- Molecule 1: Conotoxin Ar1446




4.2.26 Score per residue for model 26

- Molecule 1: Conotoxin Ar1446




4.2.27 Score per residue for model 27

- Molecule 1: Conotoxin Ar1446




4.2.28 Score per residue for model 28

- Molecule 1: Conotoxin Ar1446




4.2.29 Score per residue for model 29

- Molecule 1: Conotoxin Ar1446



4.2.30 Score per residue for model 30

- Molecule 1: Conotoxin Ar1446



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 75 calculated structures, 30 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	113
Number of shifts mapped to atoms	113
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	68%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HYP, NH2

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	86	77	79	0±0
All	All	2580	2310	2370	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	12/15 (80%)	11±0 (92±0%)	1±0 (8±0%)	0±0 (0±0%)	100	100
All	All	360/450 (80%)	330 (92%)	30 (8%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	10/10 (100%)	8±0 (77±5%)	2±0 (23±5%)	3	28
All	All	300/300 (100%)	230 (77%)	70 (23%)	3	28

All 4 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	6	CYS	30
1	A	11	HIS	30
1	A	3	ARG	6
1	A	8	LEU	4

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

MODRES-GEOMETRY INFOmissingINFO

6.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.5 Ligand geometry [i](#)

There are no ligands in this entry.

6.6 Other polymers [i](#)

There are no such molecules in this entry.

6.7 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 68% for the well-defined parts and 68% for the entire structure.

7.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *ar1446_assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	113
Number of shifts mapped to atoms	113
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 68%, i.e. 89 atoms were assigned a chemical shift out of a possible 131. 1 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	37/65 (57%)	24/26 (92%)	13/26 (50%)	0/13 (0%)
Sidechain	50/59 (85%)	31/36 (86%)	19/20 (95%)	0/3 (0%)
Aromatic	2/7 (29%)	2/4 (50%)	0/2 (0%)	0/1 (0%)
Overall	89/131 (68%)	57/66 (86%)	32/48 (67%)	0/17 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 68%, i.e. 89 atoms were assigned a chemical shift out of a possible 131. 1 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	37/65 (57%)	24/26 (92%)	13/26 (50%)	0/13 (0%)
Sidechain	50/59 (85%)	31/36 (86%)	19/20 (95%)	0/3 (0%)
Aromatic	2/7 (29%)	2/4 (50%)	0/2 (0%)	0/1 (0%)
Overall	89/131 (68%)	57/66 (86%)	32/48 (67%)	0/17 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

