

Full wwPDB X-ray Structure Validation Report (i)

May 14, 2020 – 01:57 pm BST

PDB ID : 3M8Y

Title : Phosphopentomutase from Bacillus cereus after glucose-1,6-bisphosphate

activation

Authors: Panosian, T.D.; Nannemann, D.P.; Watkins, G.; Wadzinski, B.; Bachmann,

B.O.; Iverson, T.M.

Deposited on : 2010-03-19

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	399	89%	9%	-
1	В	399	86%	11%	-
1	С	399	2%	9%	



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 10042 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Phosphopentomutase.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace	
1	Λ	391	Total	С	N	О	Р	S	4	2	0
1	Λ	391	3079	1945	508	608	1	17	4		
1	В	390	Total	С	N	О	Р	S	0	1	0
1	Б	390	3066	1939	503	606	1	17	0	1	
1	С	390	Total	С	N	О	Р	S	0	0	0
1		390	3060	1935	503	604	1	17	8	0	

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP Q818Z9
A	-3	SER	-	EXPRESSION TAG	UNP Q818Z9
A	-2	HIS	-	EXPRESSION TAG	UNP Q818Z9
A	0	ALA	-	EXPRESSION TAG	UNP Q818Z9
A	1	SER	-	EXPRESSION TAG	UNP Q818Z9
В	-4	GLY	-	EXPRESSION TAG	UNP Q818Z9
В	-3	SER	-	EXPRESSION TAG	UNP Q818Z9
В	-2	HIS	-	EXPRESSION TAG	UNP Q818Z9
В	0	ALA	-	EXPRESSION TAG	UNP Q818Z9
В	1	SER	-	EXPRESSION TAG	UNP Q818Z9
С	-4	GLY	-	EXPRESSION TAG	UNP Q818Z9
С	-3	SER	-	EXPRESSION TAG	UNP Q818Z9
С	-2	HIS	-	EXPRESSION TAG	UNP Q818Z9
С	0	ALA	-	EXPRESSION TAG	UNP Q818Z9
С	1	SER	-	EXPRESSION TAG	UNP Q818Z9

• Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	4	Total Mn 4 4	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Mn 3 3	0	0
2	С	3	Total Mn 3 3	0	0


 \bullet Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $\mathrm{C_2H_3O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0

 \bullet Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $\mathrm{C_3H_8O_3}).$






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	В	1	Total C O 6 3 3	0	0
4	В	1	Total C O 6 3 3	0	0
4	В	1	Total C O 6 3 3	0	0

 \bullet Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3).$





Mol	Chain	Residues	${f Atoms}$				ZeroOcc	AltConf
5	A	1	Total 8		N 1		0	0
5	С	1	Total 8	C 4	N 1	O 3	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	269	Total O 269 269	0	0
6	В	322	Total O 322 322	0	0
6	С	164	Total O 164 164	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphopentomutase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	91.15Å 76.66Å 107.76Å	Depositor
a, b, c, α , β , γ	90.00° 108.92° 90.00°	Depositor
Resolution (Å)	20.00 - 2.10	Depositor
Resolution (A)	45.90 - 2.10	EDS
% Data completeness	91.8 (20.00-2.10)	Depositor
(in resolution range)	91.8 (45.90-2.10)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$< I/\sigma(I) > 1$	3.97 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.0, CNS	Depositor
P. P.	0.153 , 0.198	Depositor
R, R_{free}	0.157 , 0.209	DCC
R_{free} test set	5614 reflections (6.83%)	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.553	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38 , 46.9	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10042	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, GOL, TRS, MN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	77 1		RMSZ	# Z > 5	
1	A	0.45	0/3136	0.59	0/4233	
1	В	0.47	0/3118	0.61	$1/4209 \ (0.0\%)$	
1	С	0.42	0/3109	0.57	0/4197	
All	All	0.45	0/9363	0.59	$1/12639 \ (0.0\%)$	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mo	l Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	362	LEU	CA-CB-CG	5.06	126.93	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3079	0	3016	23	0
1	В	3066	0	3007	26	0
1	С	3060	0	3001	22	0
2	A	3	0	0	0	0
2	В	4	0	0	0	0
2	С	3	0	0	0	0
3	A	8	0	6	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	8	0	6	1	0
3	С	4	0	3	0	0
4	A	18	0	24	0	0
4	В	18	0	24	2	0
5	A	8	0	12	0	0
5	С	8	0	12	2	0
6	A	269	0	0	0	0
6	В	322	0	0	0	0
6	С	164	0	0	0	0
All	All	10042	0	9111	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4:TYR:H	1:B:376:ASN:HD21	1.18	0.89
1:A:164:HIS:HD2	1:A:166:GLU:H	1.22	0.87
1:C:4:TYR:H	1:C:376:ASN:HD21	1.22	0.87
1:A:4:TYR:H	1:A:376:ASN:HD21	1.20	0.86
1:C:164:HIS:HD2	1:C:166:GLU:H	1.28	0.79
1:B:164:HIS:HD2	1:B:166:GLU:H	1.31	0.78
1:B:240:LYS:HE3	1:B:244:ILE:HD11	1.69	0.75
1:C:45:GLN:H	1:C:303:GLN:HE21	1.34	0.74
1:A:45:GLN:H	1:A:303:GLN:HE21	1.41	0.68
1:B:45:GLN:H	1:B:303:GLN:HE21	1.45	0.64
1:B:213:HIS:HD2	1:C:234:ASP:OD1	1.80	0.64
1:B:234:ASP:OD1	1:C:213:HIS:HD2	1.80	0.63
1:A:164:HIS:HE1	1:A:201:GLY:O	1.83	0.62
1:B:227:GLU:HG3	1:B:379:VAL:HB	1.81	0.61
1:A:56:ASN:HD21	1:A:66:VAL:H	1.47	0.61
1:B:56:ASN:HD21	1:B:66:VAL:H	1.47	0.61
1:B:186:ASP:OD2	1:B:188:LYS:HE3	2.02	0.60
1:C:214:ASP:OD2	5:C:399:TRS:H21	2.01	0.60
1:B:232:ASP:OD1	1:C:213:HIS:HE1	1.86	0.59
1:A:32:ASP:OD1	1:A:58:ARG:NH2	2.34	0.57
1:A:117:GLU:HG3	1:A:124:ILE:HD12	1.87	0.57
1:C:164:HIS:HE1	1:C:201:GLY:O	1.87	0.56
1:B:164:HIS:HE1	1:B:201:GLY:O	1.89	0.55
1:C:256:THR:HG21	1:C:262:GLY:HA2	1.89	0.55



Continued from previous page...

Continuea from previo		Interatomic	Clash		
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)		
1:A:210:PRO:HG2	1:B:167:VAL:HG13	1.88	0.54		
1:B:45:GLN:H	1:B:303:GLN:NE2	2.04	0.54		
1:A:211:ASN:HD22	4:B:402:GOL:H32	1.73	0.53		
1:A:14:SER:HB2	1:A:330:ASN:HB2	1.89	0.53		
1:B:24:GLU:HG3	1:B:29:LEU:HD11	1.90	0.53		
1:A:4:TYR:N	1:A:376:ASN:HD21	1.99	0.53		
1:B:164:HIS:CD2	1:B:166:GLU:H	2.20	0.51		
1:C:14:SER:HB2	1:C:330:ASN:HB2	1.93	0.50		
1:A:164:HIS:CD2	1:A:166:GLU:H	2.14	0.49		
1:A:32:ASP:CG	1:A:58:ARG:HH22	2.16	0.49		
1:A:211:ASN:ND2	4:B:402:GOL:H32	2.28	0.49		
1:C:45:GLN:H	1:C:303:GLN:NE2	2.08	0.48		
1:B:172:GLU:HA	1:B:175:LYS:HE3	1.96	0.47		
1:A:45:GLN:H	1:A:303:GLN:NE2	2.11	0.47		
1:C:56:ASN:HD21	1:C:66:VAL:H	1.63	0.47		
1:B:41:MET:HG3	1:B:44:LEU:HD13	1.97	0.46		
1:C:56:ASN:HD21	1:C:66:VAL:HG22	1.80	0.46		
1:A:240:LYS:HE3	1:A:244:ILE:HD11	1.97	0.46		
1:A:56:ASN:HB3	1:A:73:TYR:CE2	2.51	0.46		
1:A:222[B]:ARG:NH1	1:A:230:ASP:OD2	2.47	0.46		
1:A:56:ASN:ND2	1:A:66:VAL:H	2.14	0.46		
1:B:259:ASN:HD21	1:B:284:ASP:H	1.63	0.46		
1:C:267:VAL:HA	1:C:270:LEU:HD12	1.98	0.46		
1:C:240:LYS:NZ	1:C:283:VAL:HG23	2.33	0.44		
1:C:4:TYR:N	1:C:376:ASN:HD21	2.01	0.44		
1:B:56:ASN:ND2	1:B:66:VAL:H	2.14	0.44		
1:B:256:THR:HG21	1:B:262:GLY:HA2	2.00	0.44		
1:B:14:SER:HB2	1:B:330:ASN:HB2	1.99	0.44		
1:A:21:PRO:HD3	1:A:343:TYR:CZ	2.53	0.44		
1:C:101:PHE:HE2	1:C:217:LEU:HD12	1.82	0.44		
1:A:14:SER:HB2	1:A:330:ASN:CB	2.49	0.43		
1:C:164:HIS:CD2	1:C:166:GLU:H	2.20	0.43		
1:A:193:ARG:HD3	1:A:214:ASP:OD2	2.19	0.43		
1:B:232:ASP:OD1	1:C:213:HIS:CE1	2.71	0.42		
1:B:170:LEU:HD22	1:B:174:TYR:CE2	2.54	0.42		
1:B:273:ASP:OD2	1:C:165:GLU:OE2	2.38	0.42		
1:B:273:ASP:HB2	3:B:399:ACT:H3	2.01	0.42		
1:A:202:GLU:HB2	1:A:203:PRO:HD2	2.02	0.42		
1:C:181:ARG:O	1:C:185:LEU:HD13	2.20	0.41		
1:C:214:ASP:H	5:C:399:TRS:H11	1.85	0.41		
1:B:143:GLN:HG2	1:B:148:SER:O	2.20	0.41		



There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Perce	\mathbf{ntiles}
1	A	390/399~(98%)	376 (96%)	14 (4%)	0	100	100
1	В	388/399 (97%)	373 (96%)	15 (4%)	0	100	100
1	С	387/399 (97%)	376 (97%)	11 (3%)	0	100	100
All	All	1165/1197 (97%)	1125 (97%)	40 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	334/338 (99%)	325 (97%)	9 (3%)	44 48		
1	В	332/338 (98%)	323 (97%)	9 (3%)	44 48		
1	С	331/338 (98%)	320 (97%)	11 (3%)	38 40		
All	All	997/1014 (98%)	968 (97%)	29 (3%)	42 46		

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	LEU
1	A	191	VAL



Continued from previous page...

Mol	Chain	Res	$oxed{\mathbf{Type}}$
1	A	207	THR
1	A	279	PHE
1	A	281	ASN
1	A	298	TYR
1	A	355	GLU
1	A	360	LEU
1	A	391	GLU
1	В	118	GLU
1	В	123	LYS
1	В	170	LEU
1	В	189	TYR
1	В	254	LEU
1	В	279	PHE
1	В	281	ASN
1	В	298	TYR
1	В	360	LEU
1	С	61	LYS
1	С	67	GLU
1	С	185	LEU
1	С	187	GLU
1	С	190	MET
1	С	191	VAL
1	C C C C C C C	254	LEU
1	С	279	PHE
1	С	298	TYR
1	С	316	LEU
1	\overline{C}	360	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (37) such sidechains are listed below:

Mol	Chain	${f Res}$	\mathbf{Type}
1	A	45	GLN
1	A	48	ASN
1	A	56	ASN
1	A	77	GLN
1	A	89	HIS
1	A	164	HIS
1	A	205	ASN
1	A	211	ASN
1	A	259	ASN
1	A	281	ASN
1	A	303	GLN



Continued from previous page...

Mol	Chain	Res	$egin{array}{c} ext{Type} \end{array}$
1	A	376	ASN
1	A	390	ASN
1	В	45	GLN
1	В	48	ASN
1	В	56	ASN
1	В	77	GLN
1	В	89	HIS
1	В	164	HIS
1	В	205	ASN
1	В	211	ASN
1	В	213	HIS
1	В	259	ASN
1	В	281	ASN
1	В	303	GLN
1	В	358	GLN
1	В	376	ASN
1	В	390	ASN
1	С	56	ASN
1	C C C	89	HIS
1	С	164	HIS
1	\overline{C}	205	ASN
1	С	213	HIS
1	C C C	259	ASN
1	С	281	ASN
1	С	303	GLN
1	C	376	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol Type		Chain	chain Res I	Dog	Pos	Link	B	ond leng	$_{ m gths}$	В	ond ang	les
MIOI	Mol Type Chain	LIIIK		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2			
1	TPO	С	85	1,2	8,10,11	0.75	0	10,14,16	1.18	1 (10%)		
1	TPO	A	85	1,2	8,10,11	0.69	0	10,14,16	1.27	1 (10%)		
1	TPO	В	85	1,2	8,10,11	0.75	0	10,14,16	1.24	1 (10%)		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	С	85	1,2	-	1/9/11/13	-
1	TPO	A	85	1,2	-	3/9/11/13	-
1	TPO	В	85	1,2	-	4/9/11/13	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
1	A	85	TPO	O3P-P-O2P	2.37	116.71	107.64
1	В	85	TPO	O-C-CA	-2.08	119.33	124.78
1	С	85	TPO	O3P-P-O2P	2.03	115.40	107.64

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	85	TPO	CG2-CB-OG1-P
1	A	85	TPO	CB-OG1-P-O1P
1	В	85	TPO	O-C-CA-CB
1	В	85	TPO	CG2-CB-OG1-P
1	В	85	TPO	CB-OG1-P-O1P
1	В	85	TPO	CB-OG1-P-O2P
1	С	85	TPO	O-C-CA-CB
1	A	85	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.



5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 23 ligands modelled in this entry, 10 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	TD	Chair	Dag	T : 1-	В	ond leng	gths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	401	_	5,5,5	0.37	0	5, 5, 5	0.43	0
5	TRS	A	403	_	7,7,7	0.29	0	9,9,9	0.53	0
3	ACT	A	399	_	1,3,3	1.75	0	0,3,3	0.00	-
4	GOL	В	401	-	5,5,5	0.41	0	5,5,5	0.22	0
4	GOL	В	402	_	5,5,5	0.37	0	5,5,5	0.35	0
4	GOL	A	400	_	5,5,5	0.40	0	5,5,5	0.55	0
5	TRS	С	399	-	7,7,7	0.33	0	9,9,9	0.69	0
3	ACT	С	398	_	1,3,3	1.81	0	0,3,3	0.00	-
4	GOL	В	403	-	5,5,5	0.45	0	5,5,5	0.37	0
3	ACT	В	399	-	1,3,3	1.73	0	0,3,3	0.00	-
4	GOL	A	402	-	5,5,5	0.41	0	5,5,5	0.16	0
3	ACT	В	400	-	1,3,3	1.42	0	0,3,3	0.00	-
3	ACT	A	398	_	1,3,3	1.28	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	401	_	-	0/4/4/4	-
4	GOL	В	401	_	-	0/4/4/4	-
5	TRS	A	403	_	-	3/9/9/9	_
4	GOL	В	402	-	-	0/4/4/4	-
4	GOL	A	400	_	-	0/4/4/4	-
5	TRS	С	399	-	-	2/9/9/9	-



Continued from previous page...

	Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
	4	GOL	В	403	-	-	0/4/4/4	ı
ſ	4	GOL	A	402	_	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	402	GOL	C1-C2-C3-O3
5	С	399	TRS	N-C-C1-O1
5	A	403	TRS	C1-C-C3-O3
5	A	403	TRS	C2-C-C3-O3
5	A	403	TRS	N-C-C3-O3
5	С	399	TRS	C3-C-C1-O1
4	A	402	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	402	GOL	2	0
5	С	399	TRS	2	0
3	В	399	ACT	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	390/399~(97%)	-0.35	3 (0%) 86 88	17, 26, 44, 60	1 (0%)
1	В	389/399 (97%)	-0.47	0 100 100	16, 25, 40, 47	2 (0%)
1	С	389/399 (97%)	-0.12	8 (2%) 63 68	20, 37, 63, 80	2 (0%)
All	All	1168/1197 (97%)	-0.31	11 (0%) 84 86	16, 29, 54, 80	5 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	189	TYR	4.4
1	С	110	LYS	3.7
1	С	188	LYS	3.6
1	С	355	GLU	3.2
1	С	186	ASP	3.1
1	A	203	PRO	2.7
1	С	106	GLU	2.5
1	С	105	PRO	2.2
1	A	2	ASN	2.1
1	С	390	ASN	2.1
1	A	202	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

\mathbf{N}	Iol	\mathbf{Type}	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f A}^2)$	Q<0.9
	1	TPO	С	85	11/12	0.96	0.11	30,32,39,39	0
	1	TPO	A	85	11/12	0.97	0.11	22,25,31,33	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q<0.9
1	TPO	В	85	11/12	0.97	0.10	19,21,29,30	0

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$oxed{ \mathbf{B\text{-}factors}(\mathbf{\mathring{A}}^2) }$	Q < 0.9
4	GOL	A	400	6/6	0.67	0.24	44,47,48,49	0
5	TRS	С	399	8/8	0.74	0.28	51,51,51,52	0
3	ACT	A	399	4/4	0.78	0.19	53,53,53,53	0
4	GOL	A	402	6/6	0.79	0.24	55,55,55,57	0
3	ACT	С	398	4/4	0.83	0.15	56,56,56,56	0
4	GOL	A	401	6/6	0.84	0.23	38,46,48,49	0
3	ACT	В	400	4/4	0.85	0.16	56,56,56,57	0
4	GOL	В	403	6/6	0.85	0.19	41,43,44,45	0
3	ACT	A	398	4/4	0.87	0.13	48,49,49,49	0
5	TRS	A	403	8/8	0.88	0.15	44,44,45,45	0
3	ACT	В	399	4/4	0.90	0.11	50,50,50,50	0
4	GOL	В	402	6/6	0.94	0.10	49,49,50,50	0
4	GOL	В	401	6/6	0.96	0.10	32,32,33,34	0
2	MN	В	398	1/1	0.98	0.21	40,40,40,40	1
2	MN	В	397	1/1	0.98	0.05	40,40,40,40	1
2	MN	С	397	1/1	0.98	0.03	44,44,44,44	0
2	MN	С	395	1/1	0.99	0.08	32,32,32,32	0
2	MN	A	396	1/1	1.00	0.12	20,20,20,20	0
2	MN	A	397	1/1	1.00	0.06	26,26,26,26	1
2	MN	A	395	1/1	1.00	0.11	22,22,22,22	0
2	MN	В	395	1/1	1.00	0.09	22,22,22,22	0
2	MN	С	396	1/1	1.00	0.09	27,27,27,27	0
2	MN	В	396	1/1	1.00	0.09	21,21,21,21	0



6.5 Other polymers (i)

There are no such residues in this entry.

