



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 11, 2023 – 01:03 PM EDT

PDB ID : 4M9V  
Title : Zfp57 mutant (E182Q) in complex with 5-carboxylcytosine DNA  
Authors : Liu, Y.; Olanrewaju, Y.O.; Zhang, X.; Cheng, X.  
Deposited on : 2013-08-15  
Resolution : 0.97 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 0.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1243 (1.06-0.86)
Clashscore	141614	1321 (1.06-0.86)
Ramachandran outliers	138981	1233 (1.06-0.86)
Sidechain outliers	138945	1235 (1.06-0.86)
RSRZ outliers	127900	1209 (1.06-0.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	11	
1	D	11	
2	B	11	
2	E	11	
3	C	64	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	64	 6% 91% 5% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MPD	E	101	-	-	X	-
5	ACT	B	102	-	-	X	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4151 atoms, of which 1597 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*TP\*AP\*TP\*TP\*GP\*CP\*(5CM)P\*GP\*CP\*AP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				P
1	A	11	Total 415	C 122	H 156	N 40	O 83	P 14	0	6	0
1	D	11	Total 408	C 121	H 155	N 40	O 79	P 13	0	5	0

- Molecule 2 is a DNA chain called DNA (5'-D(\*AP\*CP\*TP\*GP\*(1CC)P\*GP\*GP\*CP\*AP\*AP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				P
2	B	11	Total 350	C 108	H 124	N 43	O 65	P 10	0	0	0
2	E	11	Total 350	C 108	H 124	N 43	O 65	P 10	0	0	0

- Molecule 3 is a protein called Zinc finger protein 57.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
3	C	61	Total 989	C 315	H 477	N 107	O 84	S 6	0	4	2
3	F	61	Total 1016	C 322	H 493	N 109	O 86	S 6	0	5	1

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	132	GLY	-	expression tag	UNP Q8C6P8
C	133	PRO	-	expression tag	UNP Q8C6P8
C	134	LEU	-	expression tag	UNP Q8C6P8
C	135	GLY	-	expression tag	UNP Q8C6P8
C	136	SER	-	expression tag	UNP Q8C6P8
C	182	GLN	GLU	engineered mutation	UNP Q8C6P8

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	132	GLY	-	expression tag	UNP Q8C6P8
F	133	PRO	-	expression tag	UNP Q8C6P8
F	134	LEU	-	expression tag	UNP Q8C6P8
F	135	GLY	-	expression tag	UNP Q8C6P8
F	136	SER	-	expression tag	UNP Q8C6P8
F	182	GLN	GLU	engineered mutation	UNP Q8C6P8

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	B	1	22	6	14	2	0	0
4	C	1	22	6	14	2	0	0
4	E	1	22	6	14	2	0	0
4	F	1	22	6	14	2	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			7	2	3	2		
5	C	1	Total	C	H	O	0	0
			7	2	3	2		
5	C	1	Total	C	H	O	0	0
			7	2	3	2		
5	E	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Ca	0	0
			1	1		
6	E	1	Total	Ca	0	0
			1	1		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	2	Total	Zn	0	0
			2	2		
7	F	2	Total	Zn	0	0
			2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	66	Total O 70 70	0	5
8	B	76	Total O 82 82	0	6
8	C	93	Total O 98 98	0	9
8	D	56	Total O 59 59	0	4
8	E	83	Total O 87 87	0	4
8	F	98	Total O 105 105	0	11







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	36.39Å 96.09Å 36.41Å 90.00° 113.14° 90.00°	Depositor
Resolution (Å)	33.46 – 0.97 33.46 – 0.97	Depositor EDS
% Data completeness (in resolution range)	95.5 (33.46-0.97) 95.5 (33.46-0.97)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.22 (at 0.97Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, $R_{free}$	0.130 , 0.143 0.131 , 0.143	Depositor DCC
$R_{free}$ test set	6543 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	8.6	Xtrriage
Anisotropy	0.310	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 44.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.448 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	4151	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MPD, ACT, 1CC, 5CM, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.86	2/364 (0.5%)	1.39	2/556 (0.4%)
1	D	0.80	0/340	1.37	4/521 (0.8%)
2	B	0.64	0/228	1.22	1/348 (0.3%)
2	E	0.63	0/228	1.24	1/348 (0.3%)
3	C	0.47	0/528	0.71	0/705
3	F	0.39	0/539	0.67	0/718
All	All	0.62	2/2227 (0.1%)	1.09	8/3196 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	3[A]	DT	C3'-O3'	-5.41	1.36	1.44
1	A	3[B]	DT	C3'-O3'	-5.41	1.36	1.44

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	4[A]	DT	C3'-C2'-C1'	-6.00	95.30	102.50
1	D	4[B]	DT	C3'-C2'-C1'	-6.00	95.30	102.50
2	B	7	DG	O4'-C1'-N9	-5.86	103.90	108.00
1	A	3[A]	DT	O4'-C1'-C2'	5.81	110.55	105.90
1	A	3[B]	DT	O4'-C1'-C2'	5.81	110.55	105.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	259	156	117	2	0
1	D	253	155	125	1	0
2	B	226	124	124	1	0
2	E	226	124	124	3	0
3	C	512	477	474	4	1
3	F	523	493	490	2	0
4	B	8	14	14	0	0
4	C	8	14	14	0	0
4	E	8	14	14	7	0
4	F	8	14	14	1	0
5	B	4	3	3	3	0
5	C	8	6	6	0	0
5	E	4	3	3	1	0
6	B	1	0	0	0	0
6	E	1	0	0	0	0
7	C	2	0	0	0	0
7	F	2	0	0	0	0
8	A	70	0	0	3	0
8	B	82	0	0	2	0
8	C	98	0	0	3	0
8	D	59	0	0	1	0
8	E	87	0	0	1	0
8	F	105	0	0	1	1
All	All	2554	1597	1522	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:101:MPD:H12	4:E:101:MPD:H52	1.42	0.97
2:B:2:DC:OP2	8:B:244:HOH:O	1.93	0.85
2:E:2:DC:OP2	8:E:263:HOH:O	1.95	0.85
8:C:385:HOH:O	4:E:101:MPD:H13	1.81	0.79

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:101:MPD:H12	4:E:101:MPD:C5	2.16	0.75

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:172[B]:GLU:OE2	8:F:393:HOH:O[1_556]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	61/64 (95%)	61 (100%)	0	0	100	100
3	F	62/64 (97%)	62 (100%)	0	0	100	100
All	All	123/128 (96%)	123 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	54/55 (98%)	54 (100%)	0	100	100
3	F	56/55 (102%)	56 (100%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	110/110 (100%)	110 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	5CM	A	7	1,2	17,21,22	1.74	5 (29%)	24,30,33	1.64	4 (16%)
1	5CM	D	7	1,2	17,21,22	1.71	5 (29%)	24,30,33	1.65	4 (16%)
2	1CC	B	5	1,2	20,23,24	2.08	7 (35%)	26,33,36	1.48	6 (23%)
2	1CC	E	5	1,2	20,23,24	1.94	6 (30%)	26,33,36	1.29	4 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5CM	A	7	1,2	-	2/7/21/22	0/2/2/2
1	5CM	D	7	1,2	-	2/7/21/22	0/2/2/2
2	1CC	B	5	1,2	-	0/11/25/26	0/2/2/2
2	1CC	E	5	1,2	-	0/11/25/26	0/2/2/2

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	5	1CC	C5-C4	4.79	1.51	1.43
2	B	5	1CC	C5-C4	4.76	1.51	1.43
2	B	5	1CC	C2-N3	3.29	1.43	1.36
1	A	7	5CM	C2-N3	3.10	1.42	1.36
2	E	5	1CC	C2-N3	3.09	1.42	1.36

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	7	5CM	C5-C6-N1	-4.15	119.07	123.34
1	A	7	5CM	C5-C6-N1	-4.13	119.09	123.34
1	A	7	5CM	O4'-C1'-N1	3.84	114.73	107.86
2	B	5	1CC	O22-C21-C5	-3.73	114.26	122.46
1	D	7	5CM	O4'-C1'-N1	3.41	113.96	107.86

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	7	5CM	O4'-C1'-N1-C6
1	D	7	5CM	O4'-C1'-N1-C6
1	A	7	5CM	C2'-C1'-N1-C6
1	D	7	5CM	C2'-C1'-N1-C6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	5	1CC	2	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 6 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	MPD	C	203	-	7,7,7	0.35	0	9,10,10	0.72	0
4	MPD	E	101	-	7,7,7	0.47	0	9,10,10	1.07	1 (11%)
5	ACT	E	102	-	3,3,3	0.81	0	3,3,3	1.11	0
5	ACT	B	102	-	3,3,3	0.82	0	3,3,3	1.05	0
4	MPD	F	203	-	7,7,7	0.21	0	9,10,10	0.78	0
5	ACT	C	204	-	3,3,3	0.83	0	3,3,3	1.24	0
4	MPD	B	101	-	7,7,7	0.74	0	9,10,10	0.95	0
5	ACT	C	205	-	3,3,3	0.78	0	3,3,3	1.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	E	101	-	-	1/5/5/5	-
4	MPD	F	203	-	-	1/5/5/5	-
4	MPD	B	101	-	-	3/5/5/5	-
4	MPD	C	203	-	-	0/5/5/5	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	101	MPD	CM-C2-C1	-2.66	105.03	110.57

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	101	MPD	C1-C2-C3-C4
4	B	101	MPD	CM-C2-C3-C4
4	E	101	MPD	C2-C3-C4-C5
4	F	203	MPD	C2-C3-C4-C5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	B	101	MPD	C2-C3-C4-O4

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	101	MPD	7	0
5	E	102	ACT	1	0
5	B	102	ACT	3	0
4	F	203	MPD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9	
1	A	10/11 (90%)	-0.35	0	100	100	11, 13, 16, 16	0
1	D	10/11 (90%)	-0.39	0	100	100	11, 13, 17, 17	0
2	B	10/11 (90%)	-0.82	0	100	100	8, 12, 15, 15	0
2	E	10/11 (90%)	-0.70	0	100	100	8, 11, 14, 15	0
3	C	61/64 (95%)	-0.12	4 (6%)	18	16	7, 9, 22, 27	1 (1%)
3	F	61/64 (95%)	0.11	4 (6%)	18	16	7, 9, 22, 27	1 (1%)
All	All	162/172 (94%)	-0.14	8 (4%)	29	22	7, 10, 22, 27	2 (1%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	132	GLY	8.1
3	C	132	GLY	7.0
3	F	133	PRO	6.4
3	C	133	PRO	4.5
3	C	136	SER	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	5CM	D	7	20/21	0.94	0.09	10,16,21,22	0
1	5CM	A	7	20/21	0.95	0.08	11,16,21,22	0
2	1CC	B	5	22/23	0.99	0.05	7,8,9,9	0
2	1CC	E	5	22/23	0.99	0.05	7,8,9,9	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	ACT	E	102	4/4	0.73	0.17	20,22,26,26	0
5	ACT	B	102	4/4	0.76	0.21	19,21,26,26	0
4	MPD	C	203	8/8	0.80	0.13	13,33,60,100	0
4	MPD	F	203	8/8	0.81	0.12	18,38,89,107	0
5	ACT	C	204	4/4	0.87	0.14	18,20,24,24	0
4	MPD	E	101	8/8	0.90	0.12	14,25,64,120	0
5	ACT	C	205	4/4	0.91	0.16	18,20,24,24	0
4	MPD	B	101	8/8	0.92	0.10	10,28,77,91	0
6	CA	B	103	1/1	1.00	0.02	8,8,8,8	0
6	CA	E	103	1/1	1.00	0.04	8,8,8,8	0
7	ZN	C	201	1/1	1.00	0.03	7,7,7,7	0
7	ZN	C	202	1/1	1.00	0.02	8,8,8,8	0
7	ZN	F	201	1/1	1.00	0.05	7,7,7,7	0
7	ZN	F	202	1/1	1.00	0.04	8,8,8,8	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.