



wwPDB EM Validation Summary Report ⓘ

Nov 20, 2022 – 09:06 AM EST

PDB ID : 7MBT
EMDB ID : EMD-23746
Title : Cryo-EM structure of zebrafish TRPM5 E337A mutant in the presence of 5 mM calcium (low calcium occupancy in the transmembrane domain)
Authors : Ruan, Z.; Lu, W.; Du, J.; Haley, E.
Deposited on : 2021-04-01
Resolution : Not provided

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is unknown.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	1165	
1	B	1165	
1	C	1165	
1	D	1165	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 30760 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transient receptor potential melastatin 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	996	7587	4970	1298	1275	44	0	0
1	B	996	7587	4970	1298	1275	44	0	0
1	C	996	7587	4970	1298	1275	44	0	0
1	D	996	7587	4970	1298	1275	44	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	337	ALA	GLU	engineered mutation	UNP S5UH55
B	337	ALA	GLU	engineered mutation	UNP S5UH55
C	337	ALA	GLU	engineered mutation	UNP S5UH55
D	337	ALA	GLU	engineered mutation	UNP S5UH55

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total	C	N	O	0
			14	8	1	5	
2	B	1	Total	C	N	O	0
			14	8	1	5	
2	C	1	Total	C	N	O	0
			14	8	1	5	
2	D	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 3 is (2R)-2-(hydroxymethyl)-4-{[(25R)-10alpha,14beta,17beta-spirost-5-en-3beta-y]oxy}butyl 4-O-alpha-D-glucopyranosyl-beta-D-glucopyranoside (three-letter code: YUY) (formula: C₄₄H₇₂O₁₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
3	A	1	Total	C	O	0
			59	44	15	
3	B	1	Total	C	O	0
			59	44	15	
3	C	1	Total	C	O	0
			59	44	15	
3	D	1	Total	C	O	0
			59	44	15	

- Molecule 4 is (25R)-14beta,17beta-spirost-5-en-3beta-ol (three-letter code: YUV) (formula: $C_{27}H_{42}O_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
4	A	1	Total	C	O	0
			30	27	3	
4	B	1	Total	C	O	0
			30	27	3	
4	C	1	Total	C	O	0
			30	27	3	
4	D	1	Total	C	O	0
			30	27	3	


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transient receptor potential melastatin 5



- Molecule 1: Transient receptor potential melastatin 5



L1028	L1031	K1049	E1062	R1067	T1071	K1074	V1075	L1078	M1081	V1082	M1092	ALA	ALA	THR	VAL	GLY	THR	GLU	THR	THR	THR	THR	ASP	THR	ASP	GLY	VAL	VAL	LEU	SER	TRP	TRP	ILE	ALA	GLY	CYS	PHE	HIS	LYS	LYS	THR	LEU	THR	CYS	ASP	ASP	ARG	ALA	PRO	LYS	ALA	PRO	ARG	ILE				
ALA	GLY	SER	SER	ARG	ASP	GLN	GLN	PRO	GLN	GLY	HIS	PRO	ALA	ALA	TYR	GLY	THR	THR	THR	ASP	LYS	VAL	VAL	LEU	PRO	PRO	PHI	ILE	ASP	VAL	VAL	LEU	SER	TRP	TRP	ILE	ALA	GLY	CYS	PHE	HIS	LYS	LYS	THR	LEU	THR	CYS	ASP	ASP	ARG	ALA	PRO	LYS	ALA	PRO	ARG	SER	ILE

• Molecule 1: Transient receptor potential melastatin 5



MET	VAL	GLU	LYS	SER	SER	GLU	ARG	PHE	ASP	LYS	GLN	MET	ALA	GLY	R16	R34	W60	L63	P74	W75	L76	R77	D78	T79	S98	T105	L108	A111	G130	I131	P133	I137	D141	L144	L167	D168	H171	I175												
D178	T187	R191	T198	R202	S210	L216	C217	L218	L219	V220	H221	G222	E223	I226	L227	Y231	L241	D78	T79	S98	T105	L108	A111	G130	I131	P133	I137	D141	L144	L167	D168	H171	I175																	
G389	I392	T397	E410	L413	L414	H415	L429	ALA	ARG	LYS	ARG	MET	SER	GLY	ASN	PRO	ASN	ASN	GLU	LEU	K446	H452	K456	V457	L458	L462	C466	L473	PRO	ALA	GLU	ARG	MET	GLY	LYS	GLY	ARG	PHE	HIS	SER	GLN	LYS	H489							
R496	W501	R502	L509	R513	Q514	E516	H524	A531	K536	L537	M538	A542	A545	K555	M556	A557	Y559	A563	L566	S574	E575	V583	C588	T593	V594	L595	A601	L616	V619	M620	M621	L633	S653	GLU	VAL	SER														
LYS	ASP	GLU	GLU	PRO	PHE	ALA	GLU	LEU	ASP	LYS	LEU	LEU	THR	GLY	GLN	ALA	ALA	LEU	THR	THR	ALA	ALA	ARG	CYS	S698	T718	M721	F728	F734	L738	L739	E754	L757	Y758	F759	T763														
M779	K784	L803	R811	M814	T822	V823	L824	A825	M828	M829	T832	L833	L835	T841	H842	Q843	Q844	L845	Q846	R847	K848	L849	L850	L851	R854	M855	Y859	F860	F861	F862	L863	F864	F865	S867	V868	L870	L871	A872	V875	T876	R877	Q878	A879	L880						
P886	W890	P899	Y900	L901	H902	I903	P908	L909	D913	A914	M917	V926	N943	V951	I952	Y953	L954	L955	V956	T957	M958	V959	L960	N963	L964	L965	F969	S970	Y971	V976	Q977	W984	Y989	N990	L991	P999	I1006	K1020	LYS	THR	GLU	GLU	ASN							
THR	GLN	D1027	L1028	R1031	K1049	E1062	R1067	T1071	K1074	V1075	L1078	M1081	V1082	M1092	ALA	ALA	THR	VAL	GLU	THR	THR	THR	THR	GLY	GLY	GLU	VAL	LEU	TRP	ILE	ALA	GLU	CYS	PHE	HIS	LYS	SER	THR	LEU	LYS	ASP	ASP	ASP	PRO	LYS	ALA	PRO	LYS	ALA	PRO
ARG	SER	ILE	ALA	GLY	SER	SER	ARG	ASP	GLN	GLN	PRO	GLN	GLY	ALA	LYS	ARG	GLN	GLN	PRO	GLY	HIS	HIS	PRO	PRO	TYR	GLY	THR	THR	THR	ILE	ILE	ASP	HIS	LYS	LEU	LEU	PRO	PHE	ILE	ASP	HIS									

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	208000	Depositor
Resolution determination method	Not provided	
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	47	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: YUY, YUV, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/7776	0.45	0/10604
1	B	0.27	0/7776	0.45	0/10604
1	C	0.27	0/7776	0.45	0/10604
1	D	0.27	0/7776	0.45	0/10604
All	All	0.27	0/31104	0.45	0/42416

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7587	0	7412	212	0
1	B	7587	0	7412	219	0
1	C	7587	0	7412	217	0
1	D	7587	0	7412	211	0
2	A	14	0	13	0	0
2	B	14	0	13	0	0
2	C	14	0	13	0	0
2	D	14	0	13	0	0
3	A	59	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	59	0	0	1	0
3	C	59	0	0	1	0
3	D	59	0	0	1	0
4	A	30	0	0	0	0
4	B	30	0	0	0	0
4	C	30	0	0	0	0
4	D	30	0	0	0	0
All	All	30760	0	29700	611	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 611 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1081:MET:HE1	1:D:1082:VAL:CB	1.40	1.51
1:A:1082:VAL:CB	1:B:1081:MET:HE1	1.43	1.48
1:C:1082:VAL:CB	1:D:1081:MET:HE1	1.40	1.48
1:B:1082:VAL:CB	1:C:1081:MET:HE1	1.43	1.48
1:B:1078:LEU:HD22	1:C:1078:LEU:CD1	1.62	1.27

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	986/1165 (85%)	955 (97%)	30 (3%)	1 (0%)	51	51
1	B	986/1165 (85%)	955 (97%)	30 (3%)	1 (0%)	51	51
1	C	986/1165 (85%)	955 (97%)	30 (3%)	1 (0%)	51	51
1	D	986/1165 (85%)	955 (97%)	30 (3%)	1 (0%)	51	51

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	3944/4660 (85%)	3820 (97%)	120 (3%)	4 (0%)	54	51

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1006	ILE
1	B	1006	ILE
1	C	1006	ILE
1	D	1006	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	746/1017 (73%)	745 (100%)	1 (0%)	93	93
1	B	746/1017 (73%)	745 (100%)	1 (0%)	93	93
1	C	746/1017 (73%)	745 (100%)	1 (0%)	93	93
1	D	746/1017 (73%)	745 (100%)	1 (0%)	93	93
All	All	2984/4068 (73%)	2980 (100%)	4 (0%)	93	93

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	811	ARG
1	B	811	ARG
1	C	811	ARG
1	D	811	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	514	GLN
1	D	794	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	977	GLN
1	B	514	GLN
1	B	173	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	YUV	B	1503	-	35,35,35	0.11	0	58,58,58	0.18	0
3	YUY	B	1502	-	66,66,66	0.12	0	100,102,102	0.18	0
2	NAG	C	1501	1	14,14,15	0.29	0	17,19,21	0.87	1 (5%)
3	YUY	A	1502	-	66,66,66	0.12	0	100,102,102	0.18	0
2	NAG	A	1501	1	14,14,15	0.28	0	17,19,21	0.87	1 (5%)
2	NAG	D	1501	1	14,14,15	0.27	0	17,19,21	0.87	1 (5%)
4	YUV	A	1503	-	35,35,35	0.11	0	58,58,58	0.18	0
4	YUV	C	1503	-	35,35,35	0.11	0	58,58,58	0.18	0
4	YUV	D	1503	-	35,35,35	0.12	0	58,58,58	0.17	0
3	YUY	C	1502	-	66,66,66	0.12	0	100,102,102	0.18	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1501	1	14,14,15	0.27	0	17,19,21	0.88	1 (5%)
3	YUY	D	1502	-	66,66,66	0.13	0	100,102,102	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	YUV	B	1503	-	-	-	0/6/6/6
3	YUY	B	1502	-	-	11/21/149/149	0/8/8/8
2	NAG	C	1501	1	-	1/6/23/26	0/1/1/1
3	YUY	A	1502	-	-	11/21/149/149	0/8/8/8
2	NAG	A	1501	1	-	1/6/23/26	0/1/1/1
2	NAG	D	1501	1	-	1/6/23/26	0/1/1/1
4	YUV	A	1503	-	-	-	0/6/6/6
4	YUV	C	1503	-	-	-	0/6/6/6
4	YUV	D	1503	-	-	-	0/6/6/6
3	YUY	C	1502	-	-	11/21/149/149	0/8/8/8
2	NAG	B	1501	1	-	1/6/23/26	0/1/1/1
3	YUY	D	1502	-	-	11/21/149/149	0/8/8/8

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1501	NAG	C4-C3-C2	-2.05	108.02	111.02
2	D	1501	NAG	C4-C3-C2	-2.03	108.04	111.02
2	C	1501	NAG	C4-C3-C2	-2.01	108.07	111.02
2	A	1501	NAG	C4-C3-C2	-2.01	108.07	111.02

There are no chirality outliers.

5 of 48 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1502	YUY	C43-C29-C30-O3
3	A	1502	YUY	C28-C29-C43-O14
3	A	1502	YUY	C30-C29-C43-O14
3	B	1502	YUY	C43-C29-C30-O3
3	B	1502	YUY	C28-C29-C43-O14



There are no ring outliers.



4 monomers are involved in 4 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1502	YUY	1	0
3	A	1502	YUY	1	0
3	C	1502	YUY	1	0
3	D	1502	YUY	1	0




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-23746. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections

This section was not generated.

6.2 Central slices

This section was not generated.

6.3 Largest variance slices

This section was not generated.

6.4 Orthogonal surface views

This section was not generated.

6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution

This section was not generated.

7.2 Volume estimate versus contour level

This section was not generated.

7.3 Rotationally averaged power spectrum

This section was not generated. The rotationally averaged power spectrum had issues being displayed.

8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit

This section was not generated.