



# wwPDB X-ray Structure Validation Summary Report

Jan 7, 2024 – 04:14 am GMT

PDB ID : 5MIW  
Title : X-ray structure uridine phosphorylase from *Vibrio cholerae* in complex with uracil at 1.28 Å.  
Authors : Prokofev, I.I.; Lashkov, A.A.; Balaev, V.V.; Gabdoulkhakov, A.G.; Betzel, C.; Mikhailov, A.M.  
Deposited on : 2016-11-29  
Resolution : 1.28 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1850 (1.30-1.26)
Clashscore	141614	1926 (1.30-1.26)
Ramachandran outliers	138981	1860 (1.30-1.26)
Sidechain outliers	138945	1859 (1.30-1.26)
RSRZ outliers	127900	1807 (1.30-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	 92% 7%
1	B	253	 2% 88% 10%
1	C	253	 6% 85% 14%
1	D	253	 6% 85% 13%
1	E	253	 86% 11%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	253	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	URA	B	303	-	X	-	-
4	URA	C	304	-	X	-	-
4	URA	D	304	-	X	-	-
4	URA	F	302	-	X	-	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 14909 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	252	2115	1322	372	403	18	0	27	0
1	B	252	2134	1337	378	401	18	0	31	0
1	C	251	2148	1342	383	405	18	0	33	0
1	D	251	2134	1330	380	407	17	0	33	0
1	E	251	2107	1316	377	397	17	0	27	0
1	F	252	2172	1355	384	416	17	0	38	0


- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0
2	C	2	Total Cl 2 2	0	0
2	D	2	Total Cl 2 2	0	0
2	E	1	Total Cl 1 1	0	0
2	F	1	Total Cl 1 1	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Na 2 2	0	0
3	B	1	Total Na 1 1	0	0
3	C	1	Total Na 1 1	0	0
3	D	1	Total Na 1 1	0	0
3	E	1	Total Na 1 1	0	0

- Molecule 4 is URACIL (three-letter code: URA) (formula: C<sub>4</sub>H<sub>4</sub>N<sub>2</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 8 4 2 2	0	0
4	B	1	Total C N O 8 4 2 2	0	0
4	C	1	Total C N O 8 4 2 2	0	0
4	D	1	Total C N O 8 4 2 2	0	0
4	E	1	Total C N O 8 4 2 2	0	0
4	F	1	Total C N O 8 4 2 2	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 12 6 6	0	1
6	B	1	Total C O 12 6 6	0	1

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	C	1	Total	C	O	0	1
			12	6	6		
6	C	1	Total	C	O	0	1
			12	6	6		
6	D	1	Total	C	O	0	1
			12	6	6		
6	D	1	Total	C	O	0	0
			6	3	3		
6	E	1	Total	C	O	0	1
			12	6	6		
6	F	1	Total	C	O	0	0
			6	3	3		
6	F	1	Total	C	O	0	1
			12	6	6		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			4	2	2		
7	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	349	Total 355	O 355	0	6
8	B	312	Total 313	O 313	0	1
8	C	298	Total 302	O 302	0	4
8	D	289	Total 293	O 293	0	4
8	E	315	Total 318	O 318	0	3
8	F	330	Total 336	O 336	0	6




### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uridine phosphorylase

Chain A: 





- Molecule 1: Uridine phosphorylase

Chain B: 





- Molecule 1: Uridine phosphorylase

Chain C: 





- Molecule 1: Uridine phosphorylase

Chain D: 





- Molecule 1: Uridine phosphorylase

Chain E:  86% 11% ..



● Molecule 1: Uridine phosphorylase

Chain F:  3% 87% 11% .



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.51Å 71.95Å 89.33Å 69.22° 72.20° 85.60°	Depositor
Resolution (Å)	32.03 – 1.28 49.45 – 1.28	Depositor EDS
% Data completeness (in resolution range)	94.1 (32.03-1.28) 94.1 (49.45-1.28)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.00 (at 1.28Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.132 , 0.169 0.133 , 0.169	Depositor DCC
$R_{free}$ test set	3630 reflections (1.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	9.0	Xtrriage
Anisotropy	0.117	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 63.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	14909	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, URA, NA, GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/2167	0.82	4/2922 (0.1%)
1	B	0.68	0/2189	0.82	2/2956 (0.1%)
1	C	0.64	0/2190	0.80	2/2958 (0.1%)
1	D	0.69	0/2178	0.81	6/2943 (0.2%)
1	E	0.76	0/2156	0.93	9/2914 (0.3%)
1	F	0.68	0/2209	0.82	2/2986 (0.1%)
All	All	0.69	0/13089	0.84	25/17679 (0.1%)

There are no bond length outliers.

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	174	ARG	NE-CZ-NH2	-9.21	115.70	120.30
1	E	174[A]	ARG	NE-CZ-NH2	7.83	124.22	120.30
1	E	174[C]	ARG	NE-CZ-NH2	7.83	124.22	120.30
1	E	174[B]	ARG	NE-CZ-NH2	7.83	124.22	120.30
1	E	174[A]	ARG	NE-CZ-NH1	-6.77	116.92	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2115	0	2135	25	0
1	B	2134	0	2150	20	0
1	C	2148	0	2156	40	0
1	D	2134	0	2144	34	0
1	E	2107	0	2114	30	0
1	F	2172	0	2190	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	2	0	0	1	0
2	D	2	0	0	1	0
2	E	1	0	0	0	0
2	F	1	0	0	1	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
4	A	8	0	3	0	0
4	B	8	0	3	0	0
4	C	8	0	3	0	0
4	D	8	0	3	2	0
4	E	8	0	3	0	0
4	F	8	0	3	0	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	12	0	16	0	0
6	B	18	0	24	0	0
6	C	24	0	32	2	0
6	D	18	0	23	2	0
6	E	12	0	16	0	0
6	F	18	0	24	0	0
7	B	4	0	6	1	0
7	C	4	0	6	3	0
8	A	355	0	0	13	0
8	B	313	0	0	6	0
8	C	302	0	0	19	0
8	D	293	0	0	21	0
8	E	318	0	0	7	0
8	F	336	0	0	19	0
All	All	14909	0	13054	176	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 176 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:93[A]:THR:HG23	8:D:402:HOH:O	1.24	1.27
1:E:214[A]:CYS:SG	8:E:601:HOH:O	1.95	1.20
1:F:93[A]:THR:HG23	8:F:403:HOH:O	1.48	1.10
1:A:185[A]:GLU:OE2	8:A:401:HOH:O	1.69	1.08
1:A:214[B]:CYS:SG	8:A:636:HOH:O	2.06	1.08

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/253 (110%)	273 (98%)	5 (2%)	1 (0%)	34	10
1	B	281/253 (111%)	279 (99%)	1 (0%)	1 (0%)	34	10
1	C	283/253 (112%)	279 (99%)	3 (1%)	1 (0%)	34	10
1	D	283/253 (112%)	279 (99%)	3 (1%)	1 (0%)	34	10
1	E	277/253 (110%)	274 (99%)	2 (1%)	1 (0%)	34	10
1	F	288/253 (114%)	284 (99%)	3 (1%)	1 (0%)	41	16
All	All	1691/1518 (111%)	1668 (99%)	17 (1%)	6 (0%)	34	10

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	162	TYR
1	A	162	TYR
1	B	162	TYR
1	C	162	TYR
1	D	162	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/203 (113%)	227 (99%)	3 (1%)	69	34
1	B	232/203 (114%)	225 (97%)	7 (3%)	41	6
1	C	231/203 (114%)	226 (98%)	5 (2%)	52	14
1	D	230/203 (113%)	227 (99%)	3 (1%)	69	34
1	E	229/203 (113%)	219 (96%)	10 (4%)	28	2
1	F	235/203 (116%)	224 (95%)	11 (5%)	26	2
All	All	1387/1218 (114%)	1348 (97%)	39 (3%)	52	7

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	3	LYS
1	F	235[A]	GLU
1	F	93[A]	THR
1	F	204[A]	MET
1	F	239[A]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 41 ligands modelled in this entry, 14 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	GOL	F	304[B]	-	5,5,5	0.35	0	5,5,5	0.22	0
6	GOL	C	305[B]	-	5,5,5	0.34	0	5,5,5	0.46	0
7	EDO	C	303	-	3,3,3	0.45	0	2,2,2	0.61	0
4	URA	A	304	-	8,8,8	3.79	5 (62%)	9,10,10	1.51	2 (22%)
5	SO4	A	305	-	4,4,4	0.60	0	6,6,6	0.50	0
6	GOL	A	306[A]	-	5,5,5	0.35	0	5,5,5	0.36	0
4	URA	C	304	-	8,8,8	4.85	7 (87%)	9,10,10	2.14	4 (44%)
4	URA	F	302	-	8,8,8	4.79	7 (87%)	9,10,10	2.73	4 (44%)
6	GOL	B	305[A]	-	5,5,5	0.36	0	5,5,5	0.27	0
6	GOL	C	306[B]	-	5,5,5	0.81	0	5,5,5	1.33	1 (20%)
6	GOL	D	305[B]	-	5,5,5	0.66	0	5,5,5	2.06	3 (60%)
6	GOL	D	306	-	5,5,5	0.31	0	5,5,5	0.43	0
7	EDO	B	302	-	3,3,3	0.45	0	2,2,2	0.29	0
6	GOL	B	306	-	5,5,5	0.33	0	5,5,5	0.59	0
6	GOL	E	304[B]	-	5,5,5	1.32	1 (20%)	5,5,5	1.39	1 (20%)
6	GOL	F	304[A]	-	5,5,5	0.34	0	5,5,5	0.27	0
4	URA	B	303	-	8,8,8	3.17	7 (87%)	9,10,10	2.78	3 (33%)
6	GOL	C	305[A]	-	5,5,5	0.34	0	5,5,5	0.32	0
4	URA	E	303	-	8,8,8	2.62	4 (50%)	9,10,10	1.51	1 (11%)
6	GOL	A	306[B]	-	5,5,5	0.34	0	5,5,5	0.42	0
6	GOL	C	306[A]	-	5,5,5	0.38	0	5,5,5	0.88	0
5	SO4	B	304	-	4,4,4	0.14	0	6,6,6	0.23	0
6	GOL	D	305[A]	-	5,5,5	0.36	0	5,5,5	0.17	0
4	URA	D	304	-	8,8,8	5.18	8 (100%)	9,10,10	2.91	5 (55%)
6	GOL	B	305[B]	-	5,5,5	0.34	0	5,5,5	0.34	0
6	GOL	E	304[A]	-	5,5,5	0.58	0	5,5,5	1.05	0
6	GOL	F	303	-	5,5,5	0.31	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	F	304[B]	-	-	2/4/4/4	-
6	GOL	C	305[B]	-	-	2/4/4/4	-
7	EDO	C	303	-	-	0/1/1/1	-
4	URA	A	304	-	-	-	0/1/1/1
6	GOL	A	306[A]	-	-	0/4/4/4	-
4	URA	C	304	-	-	-	0/1/1/1
4	URA	F	302	-	-	-	0/1/1/1
6	GOL	B	305[A]	-	-	0/4/4/4	-
6	GOL	C	306[B]	-	-	3/4/4/4	-
6	GOL	D	305[B]	-	-	1/4/4/4	-
6	GOL	D	306	-	-	0/4/4/4	-
7	EDO	B	302	-	-	1/1/1/1	-
6	GOL	B	306	-	-	0/4/4/4	-
6	GOL	F	304[A]	-	-	0/4/4/4	-
6	GOL	E	304[B]	-	-	2/4/4/4	-
6	GOL	C	305[A]	-	-	2/4/4/4	-
4	URA	B	303	-	-	-	0/1/1/1
4	URA	E	303	-	-	-	0/1/1/1
6	GOL	A	306[B]	-	-	2/4/4/4	-
6	GOL	C	306[A]	-	-	0/4/4/4	-
6	GOL	D	305[A]	-	-	0/4/4/4	-
4	URA	D	304	-	-	-	0/1/1/1
6	GOL	B	305[B]	-	-	2/4/4/4	-
6	GOL	E	304[A]	-	-	2/4/4/4	-
6	GOL	F	303	-	-	0/4/4/4	-

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	304	URA	C2-N1	8.53	1.48	1.36
4	D	304	URA	C2-N1	8.23	1.47	1.36
4	F	302	URA	C2-N1	7.40	1.46	1.36
4	A	304	URA	C2-N1	7.24	1.46	1.36
4	F	302	URA	C2-N3	6.64	1.48	1.37

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	303	URA	C6-N1-C2	-7.40	117.86	122.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	304	URA	C4-N3-C2	-4.96	120.78	125.70
4	F	302	URA	O2-C2-N1	-4.22	118.14	122.79
4	C	304	URA	C6-N1-C2	-4.12	119.87	122.40
4	F	302	URA	C6-N1-C2	-3.98	119.95	122.40

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	306[B]	GOL	O1-C1-C2-C3
6	C	305[A]	GOL	O1-C1-C2-O2
6	C	305[A]	GOL	O1-C1-C2-C3
6	C	305[B]	GOL	O1-C1-C2-C3
6	C	306[B]	GOL	O1-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	303	EDO	3	0
6	D	305[B]	GOL	2	0
7	B	302	EDO	1	0
6	C	306[A]	GOL	2	0
4	D	304	URA	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	252/253 (99%)	-0.34	1 (0%) 92 90	6, 11, 21, 37	0
1	B	252/253 (99%)	-0.20	6 (2%) 59 54	6, 11, 26, 51	2 (0%)
1	C	251/253 (99%)	-0.08	14 (5%) 24 20	6, 12, 31, 59	1 (0%)
1	D	251/253 (99%)	0.05	15 (5%) 21 17	6, 13, 37, 68	1 (0%)
1	E	251/253 (99%)	-0.30	0 100 100	6, 10, 21, 36	0
1	F	252/253 (99%)	-0.18	7 (2%) 53 48	6, 11, 32, 54	1 (0%)
All	All	1509/1518 (99%)	-0.17	43 (2%) 53 48	6, 11, 29, 68	5 (0%)

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	227	ILE	7.5
1	D	227	ILE	6.7
1	D	219[A]	ILE	5.4
1	C	219[A]	ILE	5.3
1	D	225	LYS	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	SO4	A	305	5/5	0.74	0.17	52,52,52,53	5
6	GOL	C	305[A]	6/6	0.83	0.15	14,14,15,15	6
6	GOL	C	305[B]	6/6	0.83	0.15	24,27,28,28	6
6	GOL	D	306	6/6	0.89	0.11	27,30,30,32	0
6	GOL	F	303	6/6	0.89	0.12	24,25,27,28	0
7	EDO	B	302	4/4	0.89	0.10	44,44,45,45	0
7	EDO	C	303	4/4	0.91	0.12	45,45,46,47	0
6	GOL	C	306[A]	6/6	0.92	0.12	20,21,22,23	6
6	GOL	C	306[B]	6/6	0.92	0.12	15,17,18,18	6
6	GOL	D	305[A]	6/6	0.93	0.13	9,10,11,14	6
6	GOL	D	305[B]	6/6	0.93	0.13	19,21,22,22	6
6	GOL	B	305[B]	6/6	0.95	0.09	21,24,24,25	6
6	GOL	B	306	6/6	0.95	0.10	15,17,21,24	0
2	CL	E	301	1/1	0.95	0.08	37,37,37,37	0
6	GOL	E	304[A]	6/6	0.95	0.08	13,14,15,15	6
6	GOL	E	304[B]	6/6	0.95	0.08	12,14,14,14	6
6	GOL	A	306[A]	6/6	0.95	0.10	11,12,13,14	6
6	GOL	A	306[B]	6/6	0.95	0.10	20,20,22,23	6
6	GOL	B	305[A]	6/6	0.95	0.09	9,10,11,14	6
6	GOL	F	304[A]	6/6	0.96	0.10	8,9,10,13	6
6	GOL	F	304[B]	6/6	0.96	0.10	27,28,28,28	6
5	SO4	B	304	5/5	0.96	0.16	10,11,11,16	5
4	URA	D	304	8/8	0.96	0.11	9,14,16,17	8
4	URA	F	302	8/8	0.97	0.08	9,13,14,15	8
4	URA	C	304	8/8	0.97	0.09	9,12,14,15	8
2	CL	B	301	1/1	0.98	0.05	46,46,46,46	0
2	CL	D	301	1/1	0.99	0.05	20,20,20,20	1
4	URA	E	303	8/8	0.99	0.06	7,7,8,8	0
3	NA	A	302	1/1	0.99	0.08	12,12,12,12	1
3	NA	A	303	1/1	0.99	0.06	19,19,19,19	0
4	URA	A	304	8/8	0.99	0.05	9,9,10,11	0
2	CL	D	302	1/1	0.99	0.04	20,20,20,20	1
2	CL	C	302	1/1	1.00	0.05	20,20,20,20	1
4	URA	B	303	8/8	1.00	0.05	6,7,8,8	0
2	CL	F	301	1/1	1.00	0.03	23,23,23,23	1
2	CL	A	301	1/1	1.00	0.05	12,12,12,12	1
2	CL	C	301	1/1	1.00	0.03	18,18,18,18	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NA	B	307	1/1	1.00	0.04	18,18,18,18	0
3	NA	C	307	1/1	1.00	0.06	12,12,12,12	1
3	NA	D	303	1/1	1.00	0.04	21,21,21,21	0
3	NA	E	302	1/1	1.00	0.05	10,10,10,10	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.