



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2023 – 01:24 AM EDT

PDB ID : 6MIY  
Title : Crystal structure of the mCD1d/xxa (JJ239)/iNKTCR ternary complex  
Authors : Zajonc, D.M.; Bitra, A.; Janssens, J.  
Deposited on : 2018-09-20  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : **FAILED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1

## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

## 2 Entry composition i

There are 11 unique types of molecules in this entry. The entry contains 13394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T cell receptor alpha variable 11,T cell receptor alpha variable 11,T cell receptor alpha joining 18,Human nkt tcr alpha chain, CHIMERIC PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	200	1542	956	264	314	8	0	0	0
1	G	200	1533	950	262	313	8	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	initiating methionine	UNP A0A0B4J1J9
C	113	ILE	-	linker	UNP A0A0B4J1J9
G	0	MET	-	initiating methionine	UNP A0A0B4J1J9
G	113	ILE	-	linker	UNP A0A0B4J1J9

- Molecule 2 is a protein called Beta-chain,T cell receptor chain,T cell receptor beta constant 2, CHIMERIC PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	240	1877	1178	334	359	6	0	0	0
2	H	239	1872	1175	334	357	6	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	95	ASP	-	linker	UNP A2NTY6
D	96	GLU	-	linker	UNP A2NTY6
D	97	GLY	-	linker	UNP A2NTY6
D	98	TYR	-	linker	UNP A2NTY6
D	130	ALA	ALA	linker	UNP A0N8J3
D	168	CYS	SER	variant	UNP A0A5B9

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	186	SER	CYS	variant	UNP A0A5B9
H	95	ASP	-	linker	UNP A2NTY6
H	96	GLU	-	linker	UNP A2NTY6
H	97	GLY	-	linker	UNP A2NTY6
H	98	TYR	-	linker	UNP A2NTY6
H	130	ALA	ALA	linker	UNP A0N8J3
H	168	CYS	SER	variant	UNP A0A5B9
H	186	SER	CYS	variant	UNP A0A5B9

- Molecule 3 is a protein called Antigen-presenting glycoprotein CD1d1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	271	Total	C	N	O	S	0	0	0
			2171	1386	371	401	13			
3	E	274	Total	C	N	O	S	0	0	0
			2193	1397	378	405	13			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	HIS	-	expression tag	UNP A0A0R4J090
A	281	HIS	-	expression tag	UNP A0A0R4J090
A	282	HIS	-	expression tag	UNP A0A0R4J090
A	283	HIS	-	expression tag	UNP A0A0R4J090
A	284	HIS	-	expression tag	UNP A0A0R4J090
A	285	HIS	-	expression tag	UNP A0A0R4J090
E	280	HIS	-	expression tag	UNP A0A0R4J090
E	281	HIS	-	expression tag	UNP A0A0R4J090
E	282	HIS	-	expression tag	UNP A0A0R4J090
E	283	HIS	-	expression tag	UNP A0A0R4J090
E	284	HIS	-	expression tag	UNP A0A0R4J090
E	285	HIS	-	expression tag	UNP A0A0R4J090

- Molecule 4 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	B	98	Total	C	N	O	S	0	0	0
			801	511	135	148	7			
4	F	98	Total	C	N	O	S	0	0	0
			801	511	135	148	7			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a

cetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
5	J	2	Total	C	N	O	0	0	0
			28	16	2	10			
5	L	2	Total	C	N	O	0	0	0
			28	16	2	10			
5	N	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	K	3	Total	C	N	O	0	0	0
			38	22	2	14			
6	M	3	Total	C	N	O	0	0	0
			38	22	2	14			


- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	1	Total	Na	0	0
			1	1		
7	D	1	Total	Na	0	0
			1	1		
7	G	1	Total	Na	0	0
			1	1		
7	H	1	Total	Na	0	0
			1	1		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	D	1	Total	Cl	0	0
			1	1		

- Molecule 9 is N-((2S,3S,4R)-1-[(4-O-benzyl-alpha-D-galactopyranosyl)oxy]-3,4-dihydroxyoctadecan-2-yl)hexacosanamide (three-letter code: JTV) (formula: C<sub>57</sub>H<sub>105</sub>NO<sub>9</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
9	A	1	67	57	1	9	0	0
9	E	1	67	57	1	9	0	0

- Molecule 10 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
10	A	1	6	3	3	0	0
10	A	1	6	3	3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	E	1	Total	C	O	0	0
			6	3	3		
10	G	1	Total	C	O	0	0
			6	3	3		

- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	C	17	Total	O	0	0
			17	17		
11	D	35	Total	O	0	0
			35	35		
11	A	46	Total	O	0	0
			46	46		
11	B	14	Total	O	0	0
			14	14		
11	E	46	Total	O	0	0
			46	46		
11	G	34	Total	O	0	0
			34	34		
11	H	43	Total	O	0	0
			43	43		
11	F	18	Total	O	0	0
			18	18		

MolProbity and EDS failed to run properly - this section is therefore empty.

### 3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.86Å 192.48Å 150.42Å 90.00° 90.79° 90.00°	Depositor
Resolution (Å)	47.42 – 2.75	Depositor
% Data completeness (in resolution range)	96.9 (47.42-2.75)	Depositor
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 2.77Å)	Xtrriage
Refinement program	REFMAC 5.8.0222	Depositor
R, $R_{free}$	0.192 , 0.247	Depositor
Wilson B-factor (Å <sup>2</sup> )	56.7	Xtrriage
Anisotropy	0.023	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.360 for h,-k,-l	Xtrriage
Total number of atoms	13394	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 4 Model quality [i](#)

### 4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles [i](#)

#### 4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	I	1	5,3	14,14,15	0.38	0	17,19,21	0.74	0
5	NAG	I	2	5	14,14,15	0.28	0	17,19,21	0.78	0
5	NAG	J	1	5,3	14,14,15	0.38	0	17,19,21	0.78	0
5	NAG	J	2	5	14,14,15	0.34	0	17,19,21	0.78	0
6	NAG	K	1	3,6	14,14,15	0.31	0	17,19,21	0.69	0
6	NAG	K	2	6	14,14,15	0.39	0	17,19,21	0.87	0
6	FUC	K	3	6	10,10,11	0.36	0	14,14,16	0.61	0
5	NAG	L	1	5,3	14,14,15	0.40	0	17,19,21	0.94	1 (5%)
5	NAG	L	2	5	14,14,15	0.34	0	17,19,21	0.90	1 (5%)
6	NAG	M	1	3,6	14,14,15	0.29	0	17,19,21	0.81	1 (5%)
6	NAG	M	2	6	14,14,15	0.41	0	17,19,21	0.78	0
6	FUC	M	3	6	10,10,11	0.36	0	14,14,16	0.59	0
5	NAG	N	1	5,3	14,14,15	0.36	0	17,19,21	0.69	0
5	NAG	N	2	5	14,14,15	0.30	0	17,19,21	0.79	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	I	1	5,3	-	0/6/23/26	0/1/1/1
5	NAG	I	2	5	-	2/6/23/26	0/1/1/1
5	NAG	J	1	5,3	-	0/6/23/26	0/1/1/1
5	NAG	J	2	5	-	2/6/23/26	0/1/1/1
6	NAG	K	1	3,6	-	0/6/23/26	0/1/1/1
6	NAG	K	2	6	-	2/6/23/26	0/1/1/1
6	FUC	K	3	6	-	-	0/1/1/1
5	NAG	L	1	5,3	-	0/6/23/26	0/1/1/1
5	NAG	L	2	5	-	0/6/23/26	0/1/1/1
6	NAG	M	1	3,6	-	0/6/23/26	0/1/1/1
6	NAG	M	2	6	-	2/6/23/26	0/1/1/1
6	FUC	M	3	6	-	-	0/1/1/1
5	NAG	N	1	5,3	-	0/6/23/26	0/1/1/1
5	NAG	N	2	5	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
5	L	1	NAG	C1-O5-C5	2.74	115.90	112.19
5	L	2	NAG	C4-C3-C2	2.31	114.41	111.02
6	M	1	NAG	O5-C5-C6	2.05	110.42	107.20

There are no chirality outliers.


All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	I	2	NAG	C4-C5-C6-O6
6	K	2	NAG	O5-C5-C6-O6
6	K	2	NAG	C4-C5-C6-O6
5	I	2	NAG	O5-C5-C6-O6
5	J	2	NAG	C4-C5-C6-O6
5	J	2	NAG	O5-C5-C6-O6
6	M	2	NAG	O5-C5-C6-O6
6	M	2	NAG	C4-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.
















## 4.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
10	GOL	A	310	-	5,5,5	0.32	0	5,5,5	0.18	0
10	GOL	G	301	-	5,5,5	0.30	0	5,5,5	0.15	0
10	GOL	A	309	-	5,5,5	0.34	0	5,5,5	0.13	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
10	GOL	E	509	-	5,5,5	0.32	0	5,5,5	0.15	0
9	JTV	A	308	-	68,68,68	0.81	2 (2%)	75,79,79	0.78	0
9	JTV	E	508	-	68,68,68	0.80	2 (2%)	75,79,79	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	GOL	A	310	-	-	1/4/4/4	-
10	GOL	G	301	-	-	0/4/4/4	-
10	GOL	A	309	-	-	0/4/4/4	-
10	GOL	E	509	-	-	2/4/4/4	-
9	JTV	A	308	-	-	12/63/83/83	0/2/2/2
9	JTV	E	508	-	-	14/63/83/83	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	308	JTV	CAT-CAU	-4.18	1.40	1.50
9	E	508	JTV	CAT-CAU	-3.97	1.41	1.50
9	E	508	JTV	O1-C1	2.60	1.44	1.40
9	A	308	JTV	O1-C1	2.46	1.44	1.40

There are no bond angle outliers.

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	E	509	GOL	O1-C1-C2-O2
10	E	509	GOL	O1-C1-C2-C3
9	E	508	JTV	CCK-CCL-CCM-CCN
9	A	308	JTV	O6-C2-C3-O2
9	A	308	JTV	CCK-CCL-CCM-CCN
9	E	508	JTV	O6-C2-C3-O2
9	A	308	JTV	CCB-CCC-CCD-CCE
9	E	508	JTV	CCG-CCH-CCI-CCJ
9	E	508	JTV	OAR-CAQ-CBG-CBH
9	A	308	JTV	CCA-CCB-CCC-CCD

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
9	A	308	JTV	CCE-CCF-CCG-CCH
9	E	508	JTV	CBP-CBQ-CBR-CBS
9	E	508	JTV	CBK-CBL-CBM-CBN
9	A	308	JTV	CCF-CCG-CCH-CCI
9	E	508	JTV	CCA-CCB-CCC-CCD
9	E	508	JTV	CBH-CBI-CBJ-CBK
9	A	308	JTV	CAU-CAT-O4-C4
9	E	508	JTV	CCE-CCF-CCG-CCH
9	E	508	JTV	CBW-CBX-CBY-CBZ
9	A	308	JTV	CBZ-CCA-CCB-CCC
9	E	508	JTV	CAU-CAT-O4-C4
9	E	508	JTV	CCL-CCM-CCN-CCO
9	A	308	JTV	CCJ-CCK-CCL-CCM
9	A	308	JTV	CAP-CAN-NAO-CBE
9	E	508	JTV	CAP-CAN-NAO-CBE
10	A	310	GOL	O2-C2-C3-O3
9	A	308	JTV	CBG-CBH-CBI-CBJ
9	E	508	JTV	CCC-CCD-CCE-CCF
9	A	308	JTV	CCL-CCM-CCN-CCO

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





#### 4.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

#### 4.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 5 Fit of model and data

### 5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

### 5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers

EDS failed to run properly - this section is therefore empty.