



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2023 – 09:40 PM EDT

PDB ID : 3MJA
Title : Cricket Paralysis Virus IGR IRES Domain 3 RNA bound to selenate, structure #2
Authors : Kieft, J.S.; Golden, B.L.; Costantino, D.A.; Chase, E.
Deposited on : 2010-04-12
Resolution : 2.80 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
RNA backbone	3102	1227 (3.10-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	B	29	
2	C	14	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SE4	B	100	-	X	-	-
3	SE4	B	103	-	X	-	-
3	SE4	C	101	-	X	-	-
3	SE4	C	102	-	X	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 937 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a RNA chain called Domain 3 of the cricket paralysis virus intergenic region IRES RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	B	29	621	277	110	205	29	0	0	0

- Molecule 2 is a RNA chain called RNA (5'-R(P*UP*AP*AP*GP*AP*AP*AP*UP*UP*UP*AP*CP*CP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	14	296	133	51	98	14	0	0	0

- Molecule 3 is SELENATE ION (three-letter code: SE4) (formula: O₄Se).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			O	Se		
3	B	1	4	1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	Se		
3	B	1	5	4	1	0	0
3	C	1	5	4	1	0	0
3	C	1	5	4	1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Domain 3 of the cricket paralysis virus intergenic region IRES RNA

Chain B:  45% 48% 7%



- Molecule 2: RNA (5'-R(P*UP*AP*AP*GP*AP*AP*AP*UP*UP*UP*AP*CP*CP*U)-3')

Chain C:  57% 29% 14%



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	59.25Å 59.25Å 99.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.80 29.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-2.80) 99.7 (29.00-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 2.31Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.233 , 0.283 0.253 , 0.273	Depositor DCC
R_{free} test set	1428 reflections (9.49%)	wwPDB-VP
Wilson B-factor (Å ²)	33.0	Xtrriage
Anisotropy	0.115	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.16 , -10.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	937	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SE4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.54	1/694 (0.1%)	0.69	0/1078
2	C	0.70	1/330 (0.3%)	0.80	1/509 (0.2%)
All	All	0.60	2/1024 (0.2%)	0.73	1/1587 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	6203	U	OP3-P	-7.18	1.52	1.61
1	B	6174	G	OP3-P	-6.50	1.53	1.61

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6203	U	OP1-P-OP2	-5.20	111.80	119.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	6180	U	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	621	0	311	13	0
2	C	296	0	150	6	0
3	B	10	0	0	0	0
3	C	10	0	0	0	0
All	All	937	0	461	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:6177:U:H3	1:B:6199:G:H1	1.15	0.92
2:C:6211:U:O2'	2:C:6212:U:OP1	2.01	0.78
1:B:6181:C:H3'	1:B:6182:A:C5'	2.26	0.66
1:B:6175:G:N2	1:B:6202:C:H1'	2.11	0.65
1:B:6177:U:H2'	1:B:6178:A:C8	2.32	0.65
1:B:6199:G:H2'	1:B:6200:A:C8	2.38	0.58
1:B:6190:U:H3	2:C:6213:A:H61	1.57	0.53
1:B:6177:U:H2'	1:B:6178:A:H8	1.75	0.51
1:B:6178:A:H2'	1:B:6179:U:C6	2.49	0.48
1:B:6192:G:N3	2:C:6209:A:C2	2.84	0.46
1:B:6185:U:O2'	2:C:6211:U:OP1	2.30	0.44
1:B:6201:C:H2'	1:B:6202:C:O4'	2.18	0.42
1:B:6190:U:H3	2:C:6213:A:N6	2.18	0.42
1:B:6199:G:H2'	1:B:6200:A:H8	1.81	0.42
2:C:6211:U:HO2'	2:C:6212:U:P	2.33	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	B	28/29 (96%)	3 (10%)	0
2	C	13/14 (92%)	2 (15%)	2 (15%)
All	All	41/43 (95%)	5 (12%)	2 (4%)

All (5) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	B	6181	C
1	B	6182	A
1	B	6183	G
2	C	6211	U
2	C	6212	U

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	C	6210	U
2	C	6211	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SE4	B	103	-	4,4,4	4.70	4 (100%)	6,6,6	0.15	0
3	SE4	C	102	-	4,4,4	4.70	4 (100%)	6,6,6	0.14	0
3	SE4	B	100	-	4,4,4	4.78	4 (100%)	6,6,6	0.14	0
3	SE4	C	101	-	4,4,4	4.69	4 (100%)	6,6,6	0.13	0

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	102	SE4	SE-O4	-5.17	1.48	1.65
3	B	100	SE4	SE-O4	-5.17	1.48	1.65
3	C	101	SE4	SE-O3	-5.10	1.49	1.65
3	B	100	SE4	SE-O3	-5.06	1.49	1.65
3	C	101	SE4	SE-O4	-5.04	1.49	1.65
3	B	103	SE4	SE-O3	-4.99	1.49	1.65
3	B	103	SE4	SE-O4	-4.97	1.49	1.65
3	C	102	SE4	SE-O3	-4.82	1.50	1.65
3	B	100	SE4	SE-O2	-4.54	1.48	1.63
3	C	102	SE4	SE-O1	-4.43	1.49	1.63
3	B	103	SE4	SE-O2	-4.42	1.49	1.63
3	B	103	SE4	SE-O1	-4.39	1.49	1.63
3	B	100	SE4	SE-O1	-4.31	1.49	1.63
3	C	102	SE4	SE-O2	-4.31	1.49	1.63
3	C	101	SE4	SE-O2	-4.28	1.49	1.63
3	C	101	SE4	SE-O1	-4.27	1.49	1.63

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.