



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 08:51 PM BST

PDB ID : 4MJ2
Title : Crystal structure of apo-iduronidase in the R3 form
Authors : Bie, H.; Yin, J.; He, X.; Kermode, A.R.; Goddard-Borger, E.D.; Withers, S.G.;
James, M.N.G.
Deposited on : 2013-09-03
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	653	 4% 89% 9%
1	B	653	 2% 90% 7%
2	C	2	 50% 50%
3	D	5	 80% 20%
4	E	3	 67% 33%
5	F	7	 29% 57% 14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BMA	E	3	-	-	-	X

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 10336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-L-iduronidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	595	4745	3036	856	841	12	0	1	0
1	B	607	4851	3099	878	860	14	0	4	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	33	GLN	HIS	SEE REMARK 999	UNP P35475
A	63	PRO	GLN	SEE REMARK 999	UNP P35475
A	105	GLN	ARG	SEE REMARK 999	UNP P35475
B	33	GLN	HIS	SEE REMARK 999	UNP P35475
B	63	PRO	GLN	SEE REMARK 999	UNP P35475
B	105	GLN	ARG	SEE REMARK 999	UNP P35475

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	2	28	16	2	10	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	5	61	34	2	25	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	E	3	39	22	2	15	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	F	7	83	46	2	35	0	0	0

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
6	A	1	14	8	1	5	0	0
6	B	1	14	8	1	5	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	A	1	6	3	3	0	0
7	B	1	6	3	3	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Cl	0	0
			1	1		
8	A	1	Total	Cl	0	0
			1	1		

- Molecule 9 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			10	4	6		
9	B	1	Total	C	O	0	0
			10	4	6		

- Molecule 10 is S,R MESO-TARTARIC ACID (three-letter code: SRT) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	B	1	Total	C O	0	0
			10	4 6		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	168	Total	O	0	0
			168	168		
11	B	265	Total	O	0	0
			265	265		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-L-iduronidase




- Molecule 1: Alpha-L-iduronidase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose






- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	259.36 Å 259.36 Å 71.81 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.01 – 2.10 49.01 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.01-2.10) 99.5 (49.01-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 2.10 Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.191 , 0.206 0.196 , 0.212	Depositor DCC
R_{free} test set	5233 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	29.3	Xtrriage
Anisotropy	0.656	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10336	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BMA, NAG, CL, TLA, SRT, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/4890	0.51	0/6680
1	B	0.29	0/4999	0.53	0/6829
All	All	0.29	0/9889	0.52	0/13509

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4745	0	4644	5	0
1	B	4851	0	4753	9	0
2	C	28	0	25	0	0
3	D	61	0	52	0	0
4	E	39	0	34	0	0
5	F	83	0	70	1	0
6	A	14	0	13	0	0
6	B	14	0	13	0	0
7	A	6	0	8	0	0
7	B	30	0	40	2	0
8	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	1	0	0	0	0
9	B	20	0	8	0	0
10	B	10	0	4	0	0
11	A	168	0	0	0	0
11	B	265	0	0	0	0
All	All	10336	0	9664	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:575:SER:HG	1:B:577[A]:CYS:HG	1.19	0.89
1:B:575:SER:OG	1:B:577[A]:CYS:SG	2.48	0.56
1:A:434:TRP:CH2	1:A:436:ALA:HB2	2.45	0.52
1:B:434:TRP:CH2	1:B:436:ALA:HB2	2.51	0.46
1:A:383:ARG:CG	1:A:387:LEU:HD23	2.48	0.43
1:B:276:GLU:OE1	1:B:329:HIS:NE2	2.44	0.43
1:B:56:LEU:CD1	5:F:7:MAN:H3	2.49	0.42
1:B:334:LEU:HD21	1:B:343:TYR:HB3	2.02	0.41
1:A:383:ARG:HG2	1:A:387:LEU:HD23	2.01	0.41
1:B:356:HIS:HB2	7:B:913:GOL:H32	2.02	0.41
1:B:471:TYR:HA	1:B:539:HIS:O	2.20	0.41
1:A:556:ALA:HB1	1:A:564:LEU:HD11	2.02	0.41
1:A:276:GLU:OE1	1:A:329:HIS:NE2	2.47	0.40
1:B:356:HIS:HB2	7:B:913:GOL:C3	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	588/653 (90%)	569 (97%)	19 (3%)	0	100	100
1	B	605/653 (93%)	586 (97%)	19 (3%)	0	100	100
All	All	1193/1306 (91%)	1155 (97%)	38 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	504/545 (92%)	499 (99%)	5 (1%)	76	82
1	B	516/545 (95%)	510 (99%)	6 (1%)	71	77
All	All	1020/1090 (94%)	1009 (99%)	11 (1%)	73	79

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	41	TRP
1	A	69	ASP
1	A	564	LEU
1	A	598	LYS
1	B	41	TRP
1	B	69	ASP
1	B	99	THR
1	B	426	ARG
1	B	555	ARG
1	B	561	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	B	603	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.54	0	17,19,21	0.87	1 (5%)
2	NAG	C	2	2	14,14,15	0.53	0	17,19,21	0.77	0
3	NAG	D	1	1,3	14,14,15	0.55	0	17,19,21	0.81	0
3	NAG	D	2	3	14,14,15	0.56	0	17,19,21	1.09	1 (5%)
3	BMA	D	3	3	11,11,12	0.30	0	15,15,17	0.76	0
3	MAN	D	4	3	11,11,12	0.61	0	15,15,17	0.65	0
3	MAN	D	5	3	11,11,12	0.58	0	15,15,17	0.80	0
4	NAG	E	1	1,4	14,14,15	0.51	0	17,19,21	1.05	1 (5%)
4	NAG	E	2	4	14,14,15	0.56	0	17,19,21	0.87	0
4	BMA	E	3	4	11,11,12	0.28	0	15,15,17	0.67	0
5	NAG	F	1	1,5	14,14,15	0.56	0	17,19,21	1.03	1 (5%)
5	NAG	F	2	5	14,14,15	0.65	0	17,19,21	0.86	1 (5%)
5	BMA	F	3	5	11,11,12	0.37	0	15,15,17	0.73	0
5	MAN	F	4	5	11,11,12	0.57	0	15,15,17	0.82	1 (6%)
5	MAN	F	5	5	11,11,12	0.73	0	15,15,17	1.13	1 (6%)
5	MAN	F	6	5	11,11,12	0.63	0	15,15,17	0.73	0
5	MAN	F	7	5	11,11,12	0.61	0	15,15,17	1.17	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	2/2/19/22	0/1/1/1
3	MAN	D	5	3	-	2/2/19/22	0/1/1/1
4	NAG	E	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	0/6/23/26	0/1/1/1
4	BMA	E	3	4	-	0/2/19/22	0/1/1/1
5	NAG	F	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	F	2	5	-	0/6/23/26	0/1/1/1
5	BMA	F	3	5	-	0/2/19/22	0/1/1/1
5	MAN	F	4	5	-	2/2/19/22	0/1/1/1
5	MAN	F	5	5	-	2/2/19/22	0/1/1/1
5	MAN	F	6	5	-	0/2/19/22	0/1/1/1
5	MAN	F	7	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	5	MAN	C1-C2-C3	3.31	113.74	109.67
3	D	2	NAG	C1-O5-C5	3.28	116.64	112.19
5	F	7	MAN	C1-C2-C3	3.10	113.47	109.67
4	E	1	NAG	C1-O5-C5	2.99	116.25	112.19
5	F	1	NAG	O5-C1-C2	-2.36	107.57	111.29
5	F	7	MAN	C1-O5-C5	2.31	115.32	112.19
2	C	1	NAG	C1-O5-C5	2.30	115.30	112.19
5	F	4	MAN	C1-O5-C5	2.24	115.22	112.19
5	F	2	NAG	C1-O5-C5	2.08	115.02	112.19

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	4	MAN	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	D	4	MAN	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
5	F	5	MAN	O5-C5-C6-O6
5	F	5	MAN	C4-C5-C6-O6
3	D	5	MAN	O5-C5-C6-O6
3	D	5	MAN	C4-C5-C6-O6
5	F	4	MAN	C4-C5-C6-O6
5	F	4	MAN	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	7	MAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	TLA	B	919	-	3,9,9	0.50	0	6,12,12	0.73	0
7	GOL	B	912	-	5,5,5	0.38	0	5,5,5	0.19	0
6	NAG	A	901	1	14,14,15	0.59	0	17,19,21	1.07	1 (5%)
7	GOL	B	913	-	5,5,5	0.35	0	5,5,5	0.32	0
7	GOL	B	914	-	5,5,5	0.28	0	5,5,5	0.36	0
7	GOL	A	909	-	5,5,5	0.26	0	5,5,5	0.19	0
9	TLA	B	917	-	3,9,9	0.44	0	6,12,12	0.98	0
6	NAG	B	901	1	14,14,15	0.50	0	17,19,21	0.81	0
7	GOL	B	915	-	5,5,5	0.31	0	5,5,5	0.20	0
7	GOL	B	916	-	5,5,5	0.26	0	5,5,5	0.47	0
10	SRT	B	918	-	3,9,9	0.42	0	6,12,12	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	TLA	B	919	-	-	0/4/12/12	-
7	GOL	B	912	-	-	4/4/4/4	-
6	NAG	A	901	1	-	0/6/23/26	0/1/1/1
7	GOL	B	913	-	-	0/4/4/4	-
7	GOL	B	914	-	-	2/4/4/4	-
7	GOL	A	909	-	-	2/4/4/4	-
9	TLA	B	917	-	-	0/4/12/12	-
6	NAG	B	901	1	-	0/6/23/26	0/1/1/1
7	GOL	B	915	-	-	0/4/4/4	-
7	GOL	B	916	-	-	0/4/4/4	-
10	SRT	B	918	-	-	4/4/12/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	901	NAG	C4-C3-C2	2.62	114.86	111.02

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	912	GOL	O1-C1-C2-C3
10	B	918	SRT	C1-C2-C3-C4
10	B	918	SRT	O2-C2-C3-O3
7	B	914	GOL	O1-C1-C2-O2
7	B	912	GOL	C1-C2-C3-O3
7	B	914	GOL	O1-C1-C2-C3
7	A	909	GOL	O1-C1-C2-C3
7	B	912	GOL	O1-C1-C2-O2
7	A	909	GOL	O1-C1-C2-O2
10	B	918	SRT	O2-C2-C3-C4
7	B	912	GOL	O2-C2-C3-O3
10	B	918	SRT	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	913	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	595/653 (91%)	-0.01	28 (4%) 31 37	22, 36, 64, 86	0
1	B	607/653 (92%)	-0.12	16 (2%) 56 61	20, 30, 50, 68	0
All	All	1202/1306 (92%)	-0.07	44 (3%) 41 48	20, 33, 61, 86	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	336	ASN	5.0
1	A	639	LEU	4.7
1	A	588	ASP	4.3
1	B	588	ASP	4.2
1	A	616	GLY	4.0
1	B	336	ASN	3.9
1	B	338	THR	3.9
1	B	589	GLY	3.8
1	B	590	LYS	3.7
1	A	529	ALA	3.6
1	A	338	THR	3.5
1	B	642	PRO	3.4
1	B	641	VAL	3.3
1	B	229	PRO	3.2
1	A	594	PRO	3.1
1	A	468	GLY	3.0
1	A	593	THR	3.0
1	A	521	GLY	2.9
1	B	186	HIS	2.9
1	B	426	ARG	2.9
1	A	451	ASN	2.7
1	B	189	ASP	2.7
1	A	408	ALA	2.7
1	B	103	THR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	626	TRP	2.6
1	A	452	ARG	2.6
1	A	189	ASP	2.6
1	A	586	SER	2.6
1	A	64	TYR	2.4
1	A	481	CYS	2.4
1	A	523	ARG	2.4
1	A	635	PRO	2.3
1	A	470	VAL	2.3
1	B	591	ALA	2.3
1	A	339	SER	2.3
1	A	632	PHE	2.3
1	B	337	THR	2.3
1	A	186	HIS	2.2
1	B	451	ASN	2.2
1	A	455	ALA	2.2
1	A	637	PRO	2.2
1	A	337	THR	2.1
1	B	339	SER	2.1
1	A	411	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MAN	F	7	11/12	0.59	0.25	75,77,79,80	0
4	BMA	E	3	11/12	0.66	0.40	85,86,87,88	0
5	MAN	F	5	11/12	0.71	0.23	72,74,77,79	0
3	MAN	D	4	11/12	0.73	0.34	80,84,86,87	0
4	NAG	E	2	14/15	0.78	0.40	68,74,76,81	0
3	MAN	D	5	11/12	0.80	0.25	73,73,74,75	0
2	NAG	C	2	14/15	0.81	0.37	66,72,73,74	0
3	BMA	D	3	11/12	0.85	0.12	60,65,70,74	0

Continued on next page...

Continued from previous page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	MAN	F	6	11/12	0.87	0.13	58,61,66,72	0
5	MAN	F	4	11/12	0.88	0.12	57,58,61,66	0
4	NAG	E	1	14/15	0.89	0.23	43,48,55,62	0
5	BMA	F	3	11/12	0.91	0.12	42,46,51,53	0
2	NAG	C	1	14/15	0.92	0.16	44,47,53,59	0
3	NAG	D	2	14/15	0.93	0.12	47,48,55,55	0
5	NAG	F	2	14/15	0.94	0.13	33,35,38,40	0
3	NAG	D	1	14/15	0.94	0.11	37,41,44,44	0
5	NAG	F	1	14/15	0.96	0.13	27,30,32,32	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	TLA	B	919	10/10	0.77	0.19	66,68,71,71	0
7	GOL	A	909	6/6	0.77	0.22	46,48,49,52	0
7	GOL	B	912	6/6	0.81	0.19	42,43,43,43	0
10	SRT	B	918	10/10	0.86	0.23	50,53,53,54	0
7	GOL	B	915	6/6	0.88	0.15	48,50,53,54	0
6	NAG	A	901	14/15	0.89	0.21	51,57,60,60	0
7	GOL	B	913	6/6	0.89	0.13	39,40,41,41	0
7	GOL	B	914	6/6	0.92	0.14	37,38,38,38	0
9	TLA	B	917	10/10	0.93	0.11	37,39,40,42	0
6	NAG	B	901	14/15	0.94	0.13	35,40,43,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	GOL	B	916	6/6	0.94	0.09	31,32,33,34	0
8	CL	B	920	1/1	0.99	0.06	29,29,29,29	0
8	CL	A	910	1/1	0.99	0.10	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.