

Full wwPDB X-ray Structure Validation Report (i)

Oct 1, 2023 – 10:32 PM EDT

PDB ID	:	6MJN
Title	:	Crystal structure of an organic hydroperoxide resistance protein OsmC, pre-
		dicted redox protein, regulator of sulfide bond formation from Legionella pneu- mophila
Authors	:	Seattle Structural Genomics Center for Infectious Disease; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on Resolution		

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	FAILED
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 1.75 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4181 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	138	Total	С	Ν	0	S	0	6	0
	A	130	1027	645	180	198	4	0		0
1	В	3 140	Total	С	Ν	Ο	S	0	4	0
	I D		1026	644	180	197	5			
1	C	C 137	Total	С	Ν	0	S	0	1	0
	U		970	605	165	196	4	0	1	0
1 D	139	Total	С	Ν	0	S	0	1	0	
		952	594	164	189	5			0	

• Molecule 1 is a protein called Organic hydroperoxide resistance protein.

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-7	MET	-	expression tag	UNP A0A2S6F4M7
А	-6	ALA	-	expression tag	UNP A0A2S6F4M7
А	-5	HIS	-	expression tag	UNP A0A2S6F4M7
А	-4	HIS	-	expression tag	UNP A0A2S6F4M7
А	-3	HIS	-	expression tag	UNP A0A2S6F4M7
А	-2	HIS	-	expression tag	UNP A0A2S6F4M7
А	-1	HIS	-	expression tag	UNP A0A2S6F4M7
А	0	HIS	-	expression tag	UNP A0A2S6F4M7
В	-7	MET	-	expression tag	UNP A0A2S6F4M7
В	-6	ALA	-	expression tag	UNP A0A2S6F4M7
В	-5	HIS	-	expression tag	UNP A0A2S6F4M7
В	-4	HIS	-	expression tag	UNP A0A2S6F4M7
В	-3	HIS	-	expression tag	UNP A0A2S6F4M7
В	-2	HIS	-	expression tag	UNP A0A2S6F4M7
В	-1	HIS	-	expression tag	UNP A0A2S6F4M7
В	0	HIS	-	expression tag	UNP A0A2S6F4M7
С	-7	MET	-	expression tag	UNP A0A2S6F4M7
С	-6	ALA	-	expression tag	UNP A0A2S6F4M7
С	-5	HIS	-	expression tag	UNP A0A2S6F4M7
С	-4	HIS	-	expression tag	UNP A0A2S6F4M7
С	-3	HIS	-	expression tag	UNP A0A2S6F4M7

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	HIS	-	expression tag	UNP A0A2S6F4M7
С	-1	HIS	-	expression tag	UNP A0A2S6F4M7
С	0	HIS	-	expression tag	UNP A0A2S6F4M7
D	-7	MET	-	expression tag	UNP A0A2S6F4M7
D	-6	ALA	-	expression tag	UNP A0A2S6F4M7
D	-5	HIS	-	expression tag	UNP A0A2S6F4M7
D	-4	HIS	-	expression tag	UNP A0A2S6F4M7
D	-3	HIS	-	expression tag	UNP A0A2S6F4M7
D	-2	HIS	-	expression tag	UNP A0A2S6F4M7
D	-1	HIS	-	expression tag	UNP A0A2S6F4M7
D	0	HIS	-	expression tag	UNP A0A2S6F4M7

Continued from previous page...

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Cl 1 1	0	0
2	В	1	Total Cl 1 1	0	0
2	D	1	Total Cl 1 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	80	Total O 82 82	0	2
3	В	72	Total O 72 72	0	0
3	С	30	Total O 30 30	0	0
3	D	19	Total O 19 19	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	76.55Å 51.40Å 82.34Å	Depositor
a, b, c, α , β , γ	90.00° 110.09° 90.00°	Depositor
Resolution (Å)	42.81 - 1.75	Depositor
% Data completeness	92.8 (42.81-1.75)	Depositor
(in resolution range)		-
R _{merge}	0.03	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.95 (at 1.75 \text{\AA})$	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.181 , 0.209	Depositor
Wilson B-factor ($Å^2$)	33.4	Xtriage
Anisotropy	0.088	Xtriage
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4181	wwPDB-VP
Average B, all atoms $(Å^2)$	53.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

