



# Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 11:49 pm BST

PDB ID : 2MKK  
Title : Structural model of tandem RRM domains of cytoplasmic polyadenylation element binding protein 1 (CPEB1) in complex with RNA  
Authors : Afroz, T.; Skrisovska, L.; Belloc, E.; Boixet, J.G.; Mendez, R.; Allain, F.H.-T.  
Deposited on : 2014-02-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 47%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428
RNA backbone	4643	676

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	213	 83% . . 12%
2	B	5	 60% 40%

## 2 Ensemble composition and analysis i

This entry contains 10 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:218-A:267, A:279-A:301, A:310-A:385, A:393-A:430 (187)	1.34	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 5, 6, 7
2	2, 3, 4
3	8, 9
Single-model clusters	10

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3532 atoms, of which 1746 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Cytoplasmic polyadenylation element-binding protein 1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	213	3380	1088	1693	290	301	8	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	218	MET	-	EXPRESSION TAG	UNP Q9BZB8

- Molecule 2 is a RNA chain called RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3').


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
2	B	5	152	46	53	13	36	4	0

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1



- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1




- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')






### 4.2.2 Score per residue for model 2 (medoid)

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1




- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')




### 4.2.3 Score per residue for model 3

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1




- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')




### 4.2.4 Score per residue for model 4

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1







- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')



#### 4.2.5 Score per residue for model 5

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1





- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')



#### 4.2.6 Score per residue for model 6

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1




- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')




### 4.2.7 Score per residue for model 7

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1




- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')



### 4.2.8 Score per residue for model 8

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1




- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')



### 4.2.9 Score per residue for model 9

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1






- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')


Chain B: 




#### 4.2.10 Score per residue for model 10

- Molecule 1: Cytoplasmic polyadenylation element-binding protein 1

Chain A: 





- Molecule 2: RNA (5'-R(\*UP\*UP\*UP\*UP\*A)-3')

Chain B: 



## 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 20 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	
CYANA	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	1463
Number of shifts mapped to atoms	1463
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	47%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

### 5.1 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1490	1496	1493	7±2
2	B	99	53	53	3±1
All	All	15890	15490	15460	75

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:295:LEU:H	1:A:295:LEU:HD13	0.66	1.50	8	2
1:A:405:LEU:HD22	1:A:405:LEU:C	0.60	2.16	3	5
1:A:324:LYS:CD	2:B:1:U:C6	0.60	2.84	5	1
1:A:239:PHE:CE2	2:B:3:U:C6	0.60	2.89	6	4
1:A:405:LEU:C	1:A:405:LEU:HD22	0.59	2.17	7	3
1:A:347:ASP:H	1:A:348:PRO:HD2	0.57	1.59	8	1
1:A:324:LYS:HE3	2:B:1:U:C6	0.55	2.37	9	1
1:A:357:LEU:HD12	1:A:359:GLY:H	0.55	1.62	1	1
1:A:382:ILE:N	1:A:382:ILE:HD13	0.55	2.17	9	1
1:A:285:VAL:HG12	1:A:286:PHE:H	0.53	1.62	9	1
1:A:283:TYR:CD1	2:B:4:U:C4	0.51	2.98	10	5
1:A:266:TRP:CG	1:A:267:PRO:HD2	0.50	2.42	10	1
2:B:5:A:H5'	2:B:5:A:C8	0.50	2.42	1	1
1:A:324:LYS:HE3	2:B:1:U:C1'	0.49	2.38	8	1
1:A:347:ASP:N	1:A:348:PRO:CD	0.49	2.75	5	2
1:A:324:LYS:HD2	2:B:1:U:C6	0.49	2.42	5	2
1:A:353:PHE:CE2	2:B:5:A:H1'	0.48	2.43	2	3
1:A:283:TYR:CD2	2:B:4:U:C2	0.47	3.02	8	1
1:A:361:LEU:H	1:A:361:LEU:CD1	0.47	2.22	10	1
1:A:283:TYR:CD2	2:B:4:U:C4	0.47	3.03	4	2
1:A:231:ASN:N	1:A:232:PRO:CD	0.47	2.78	1	1
1:A:324:LYS:HD3	2:B:1:U:C6	0.47	2.44	5	1
1:A:357:LEU:HD23	1:A:357:LEU:H	0.46	1.69	6	1
1:A:283:TYR:CD2	2:B:4:U:N3	0.46	2.84	8	1
1:A:324:LYS:CE	2:B:1:U:O4'	0.46	2.64	10	1
1:A:283:TYR:CE2	2:B:4:U:C2	0.46	3.04	9	1
1:A:285:VAL:HG12	1:A:286:PHE:N	0.46	2.25	9	1
1:A:283:TYR:CG	2:B:4:U:C4	0.45	3.05	7	1
1:A:361:LEU:H	1:A:361:LEU:HD13	0.45	1.72	10	1
1:A:339:VAL:HG13	1:A:341:SER:H	0.45	1.71	5	2
1:A:285:VAL:HG22	1:A:286:PHE:H	0.45	1.72	6	1
1:A:224:LEU:HD23	1:A:224:LEU:H	0.44	1.73	8	1
1:A:252:LEU:HA	1:A:255:THR:HG22	0.44	1.90	2	1
1:A:281:TYR:N	1:A:281:TYR:CD1	0.43	2.86	10	1
1:A:357:LEU:HD23	1:A:357:LEU:N	0.43	2.29	4	1
1:A:239:PHE:CZ	2:B:3:U:O4'	0.43	2.72	10	2
1:A:239:PHE:CZ	1:A:241:GLY:CA	0.43	3.02	5	1
2:B:5:A:C8	2:B:5:A:H5''	0.42	2.49	8	1
1:A:262:LEU:C	1:A:262:LEU:HD13	0.42	2.35	6	1
1:A:360:MET:SD	1:A:360:MET:N	0.42	2.92	8	2
1:A:339:VAL:CG1	1:A:341:SER:H	0.42	2.27	7	1
1:A:331:TRP:CD2	1:A:384:THR:HG23	0.42	2.49	8	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:283:TYR:CG	2:B:4:U:N3	0.42	2.88	10	1
1:A:266:TRP:CD2	1:A:267:PRO:HD2	0.42	2.50	10	1
1:A:262:LEU:HD13	1:A:262:LEU:C	0.42	2.35	1	1
1:A:338:PHE:CG	1:A:339:VAL:N	0.41	2.89	10	1
1:A:357:LEU:H	1:A:357:LEU:CD2	0.41	2.29	6	1
1:A:283:TYR:CD1	2:B:4:U:N3	0.41	2.89	3	2
1:A:347:ASP:H	1:A:348:PRO:CD	0.40	2.27	8	1
1:A:245:TRP:CZ2	1:A:324:LYS:HE3	0.40	2.51	2	1
1:A:314:PHE:CG	1:A:315:LYS:N	0.40	2.90	8	1
1:A:244:PRO:HA	1:A:245:TRP:CE3	0.40	2.52	4	1

## 5.2 Torsion angles [i](#)

### 5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/213 (87%)	163±6 (88±3%)	19±6 (10±3%)	4±2 (2±1%)	10	50
All	All	1850/2130 (87%)	1626 (88%)	186 (10%)	38 (2%)	10	50

All 26 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	287	GLU	3
1	A	357	LEU	3
1	A	249	GLU	3
1	A	286	PHE	3
1	A	242	GLY	2
1	A	301	HIS	2
1	A	358	HIS	2
1	A	359	GLY	2
1	A	383	ASP	1
1	A	281	TYR	1
1	A	323	CYS	1
1	A	318	SER	1

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	259	PHE	1
1	A	336	SER	1
1	A	243	VAL	1
1	A	261	SER	1
1	A	345	ARG	1
1	A	385	ASP	1
1	A	335	ASP	1
1	A	280	GLY	1
1	A	347	ASP	1
1	A	356	ALA	1
1	A	374	PHE	1
1	A	322	ARG	1
1	A	333	LEU	1
1	A	421	THR	1

### 5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	163/185 (88%)	154±2 (95±1%)	9±2 (5±1%)	25 74
All	All	1630/1850 (88%)	1541 (95%)	89 (5%)	25 74

All 37 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	281	TYR	10
1	A	405	LEU	10
1	A	286	PHE	7
1	A	382	ILE	6
1	A	339	VAL	5
1	A	420	PHE	4
1	A	360	MET	4
1	A	366	LEU	3
1	A	295	LEU	3
1	A	247	ILE	3
1	A	253	VAL	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	357	LEU	2
1	A	361	LEU	2
1	A	262	LEU	2
1	A	321	MET	2
1	A	315	LYS	2
1	A	284	LEU	2
1	A	422	LYS	1
1	A	324	LYS	1
1	A	395	ARG	1
1	A	240	LEU	1
1	A	373	LEU	1
1	A	358	HIS	1
1	A	224	LEU	1
1	A	243	VAL	1
1	A	419	LYS	1
1	A	228	ASN	1
1	A	346	LEU	1
1	A	399	ASN	1
1	A	327	GLN	1
1	A	283	TYR	1
1	A	337	ASN	1
1	A	234	TYR	1
1	A	296	LEU	1
1	A	333	LEU	1
1	A	329	ILE	1
1	A	370	LEU	1

### 5.2.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
2	B	4/5 (80%)	2±1 (55±24%)	0±0 (5±10%)	0.10±0.08
All	All	40/50 (80%)	22 (55%)	2 (5%)	0.10

The overall RNA backbone suiteness is 0.10.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	B	4	U	10
2	B	5	A	6
2	B	2	U	5
2	B	3	U	1

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	B	2	U	1
2	B	4	U	1

### 5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.5 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.6 Other polymers [i](#)

There are no such molecules in this entry.

### 5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 47% for the well-defined parts and 45% for the entire structure.

### 6.1 Chemical shift list 1

File name: input\_cs.cif

Chemical shift list name: *CPEB1RRM12Complex*

#### 6.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1463
Number of shifts mapped to atoms	1463
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 6.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	110	$2.70 \pm 0.13$	Should be applied
$^{13}\text{C}_\beta$	87	$2.79 \pm 0.25$	Should be applied
$^{13}\text{C}'$	0	—	None (insufficient data)
$^{15}\text{N}$	183	$-0.08 \pm 0.35$	None needed ( $< 0.5$ ppm)

#### 6.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 47%, i.e. 1149 atoms were assigned a chemical shift out of a possible 2425. 26 out of 37 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	532/917 (58%)	266/365 (73%)	101/374 (27%)	165/178 (93%)
Sidechain	463/1188 (39%)	271/699 (39%)	185/431 (43%)	7/58 (12%)

*Continued on next page...*



Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	154/233 (66%)	95/123 (77%)	55/104 (53%)	4/6 (67%)
Overall	1149/2425 (47%)	632/1234 (51%)	341/944 (36%)	176/247 (71%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 45%, i.e. 1241 atoms were assigned a chemical shift out of a possible 2743. 26 out of 39 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	586/1035 (57%)	293/411 (71%)	110/426 (26%)	183/198 (92%)
Sidechain	497/1366 (36%)	292/809 (36%)	198/492 (40%)	7/65 (11%)
Aromatic	158/255 (62%)	97/135 (72%)	57/112 (51%)	4/8 (50%)
Overall	1241/2743 (45%)	682/1402 (49%)	365/1065 (34%)	194/276 (70%)

#### 6.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 6.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

