



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 3, 2024 – 06:49 PM EST

PDB ID : 1MUY
Title : CATALYTIC DOMAIN OF MUTY FROM ESCHERICHIA COLI
Authors : Guan, Y.; Tainer, J.A.
Deposited on : 1998-08-20
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	225	 84% 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	303	-	X	X	-
4	IMD	A	301	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2027 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADENINE GLYCOSYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	225	1762	1133	306	312	11	0	0	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	6	3	3	0	0

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe S	0	0
			8	4 4		

- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C N	0	0
			5	3 2		
4	A	1	Total	C N	0	0
			5	3 2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	241	Total 241	O 241	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: ADENINE GLYCOSYLASE



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	83.06Å 48.95Å 69.03Å 90.00° 123.37° 90.00°	Depositor
Resolution (Å)	20.00 – 1.40	Depositor
% Data completeness (in resolution range)	92.0 (20.00-1.40)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.153 , 0.231	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2027	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, IMD, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/1809	1.36	18/2456 (0.7%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	119	ARG	CD-NE-CZ	15.79	145.70	123.60
1	A	119	ARG	NE-CZ-NH2	15.22	127.91	120.30
1	A	119	ARG	NE-CZ-NH1	-12.30	114.15	120.30
1	A	58	ARG	NE-CZ-NH2	-11.73	114.44	120.30
1	A	143	ARG	CD-NE-CZ	11.57	139.81	123.60
1	A	143	ARG	NE-CZ-NH1	10.86	125.73	120.30
1	A	147	ARG	NE-CZ-NH2	10.25	125.42	120.30
1	A	194	ARG	NE-CZ-NH1	9.92	125.26	120.30
1	A	87	ARG	NE-CZ-NH1	-9.13	115.73	120.30
1	A	27	ASP	CB-CG-OD2	8.26	125.73	118.30
1	A	179	ARG	CD-NE-CZ	6.96	133.34	123.60
1	A	82	TYR	CB-CG-CD1	-6.17	117.30	121.00
1	A	143	ARG	NH1-CZ-NH2	-6.03	112.77	119.40
1	A	147	ARG	NE-CZ-NH1	-5.91	117.34	120.30
1	A	87	ARG	NE-CZ-NH2	5.64	123.12	120.30
1	A	87	ARG	CD-NE-CZ	-5.55	115.83	123.60
1	A	54	ARG	NE-CZ-NH2	-5.39	117.60	120.30
1	A	82	TYR	CB-CG-CD2	5.26	124.16	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1762	0	1771	26	0
2	A	6	0	7	5	0
3	A	8	0	0	0	0
4	A	10	0	10	4	0
5	A	241	0	0	3	0
All	All	2027	0	1788	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:303:GOL:C2	2:A:303:GOL:C3	1.93	1.46
2:A:303:GOL:O1	2:A:303:GOL:C1	1.78	1.31
2:A:303:GOL:C3	2:A:303:GOL:C1	2.61	0.78
1:A:19:ARG:HD3	1:A:21:THR:H	1.51	0.74
1:A:20:LYS:HG3	1:A:25:GLN:HB3	1.70	0.71
1:A:222:LYS:HD3	5:A:1058:HOH:O	1.95	0.66
1:A:194:ARG:HG3	5:A:1035:HOH:O	1.97	0.64
1:A:225:LYS:HB3	1:A:225:LYS:NZ	2.11	0.63
1:A:154:TRP:CZ2	1:A:223:LYS:HG2	2.37	0.58
1:A:19:ARG:CD	1:A:21:THR:H	2.18	0.55
1:A:19:ARG:HH11	1:A:21:THR:HB	1.71	0.54
1:A:19:ARG:HD3	1:A:21:THR:N	2.21	0.53
1:A:21:THR:HA	1:A:26:ILE:HD11	1.89	0.53
1:A:140:ASN:HB3	1:A:188:GLY:HA3	1.91	0.52
2:A:303:GOL:C2	2:A:303:GOL:O3	2.50	0.52
1:A:1:MET:HG3	1:A:5:GLN:HB3	1.92	0.51
1:A:224:PRO:O	1:A:225:LYS:HB2	2.12	0.49
1:A:87:ARG:HD3	5:A:1115:HOH:O	2.13	0.49
2:A:303:GOL:C3	2:A:303:GOL:O2	2.58	0.49
1:A:16:LYS:HD2	1:A:16:LYS:N	2.28	0.49
1:A:138:ASP:HB3	4:A:301:IMD:C4	2.45	0.47
1:A:120:SER:OG	4:A:301:IMD:N1	2.48	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:ILE:HB	4:A:301:IMD:H2	1.96	0.47
1:A:19:ARG:HH11	1:A:21:THR:CB	2.28	0.46
1:A:21:THR:HG22	1:A:21:THR:O	2.16	0.46
1:A:20:LYS:CG	1:A:25:GLN:HB3	2.43	0.44
1:A:120:SER:CB	4:A:301:IMD:HN1	2.31	0.44
1:A:24:TRP:O	1:A:28:LYS:HG2	2.17	0.43
1:A:15:ASP:C	1:A:16:LYS:HD2	2.41	0.41
1:A:154:TRP:CE2	1:A:223:LYS:HG2	2.56	0.41
1:A:45:VAL:HG13	1:A:46:ALA:N	2.35	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	223/225 (99%)	222 (100%)	1 (0%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	185/185 (100%)	178 (96%)	7 (4%)	33 6

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	16	LYS
1	A	19	ARG
1	A	41	GLN
1	A	76	LEU
1	A	223	LYS
1	A	225	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	88	ASN
1	A	95	GLN
1	A	162	ASN
1	A	205	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	IMD	A	301	-	3,5,5	0.12	0	4,5,5	0.80	0
2	GOL	A	303	-	5,5,5	6.32	3 (60%)	5,5,5	2.56	3 (60%)
3	SF4	A	300	1	0,12,12	-	-	-	-	-
4	IMD	A	302	-	3,5,5	0.22	0	4,5,5	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IMD	A	301	-	-	-	0/1/1/1
2	GOL	A	303	-	-	1/4/4/4	-
3	SF4	A	300	1	-	-	0/6/5/5
4	IMD	A	302	-	-	-	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	303	GOL	C3-C2	10.15	1.93	1.51
2	A	303	GOL	O1-C1	8.52	1.78	1.42
2	A	303	GOL	O3-C3	-4.59	1.23	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	303	GOL	C3-C2-C1	-4.16	95.54	111.70
2	A	303	GOL	O2-C2-C1	2.35	119.48	109.12
2	A	303	GOL	O2-C2-C3	-2.08	99.98	109.12

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	303	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	301	IMD	4	0
2	A	303	GOL	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.