



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:40 am BST

PDB ID : 4MV2  
Title : Crystal structure of plu4264 protein from Photorhabdus luminescens  
Authors : Michalska, K.; Li, H.; Jedrzejczak, R.; Babnigg, G.; Bingman, C.A.; Yen-  
namalli, R.; Weerth, S.; Thomas, M.G.; Phillips Jr., G.N.; Joachimiak, A.;  
Midwest Center for Structural Genomics (MCSG); Enzyme Discovery for Nat-  
ural Product Biosynthesis (NatPro)  
Deposited on : 2013-09-23  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1385 (1.36-1.32)
Clashscore	141614	1417 (1.36-1.32)
Ramachandran outliers	138981	1397 (1.36-1.32)
Sidechain outliers	138945	1397 (1.36-1.32)
RSRZ outliers	127900	1369 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	128	 9% 87% 7% • 5%
1	B	128	 9% 79% 15% 6%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2638 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called plu4264.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	121	1116	718	188	206	1	3	0	12	0
1	B	120	1226	780	210	231	1	4	0	25	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	HIS	-	EXPRESSION TAG	UNP Q7MZL9
A	124	HIS	-	EXPRESSION TAG	UNP Q7MZL9
A	125	HIS	-	EXPRESSION TAG	UNP Q7MZL9
A	126	HIS	-	EXPRESSION TAG	UNP Q7MZL9
A	127	HIS	-	EXPRESSION TAG	UNP Q7MZL9
A	128	HIS	-	EXPRESSION TAG	UNP Q7MZL9
B	123	HIS	-	EXPRESSION TAG	UNP Q7MZL9
B	124	HIS	-	EXPRESSION TAG	UNP Q7MZL9
B	125	HIS	-	EXPRESSION TAG	UNP Q7MZL9
B	126	HIS	-	EXPRESSION TAG	UNP Q7MZL9
B	127	HIS	-	EXPRESSION TAG	UNP Q7MZL9
B	128	HIS	-	EXPRESSION TAG	UNP Q7MZL9

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ni	0	0
			1	1		
2	A	1	Total	Ni	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	152	Total O 154 154	0	7
4	B	137	Total O 139 139	0	13

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: plu4264

Chain A: 



- Molecule 1: plu4264

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.74Å 147.69Å 83.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.90 – 1.35 24.90 – 1.35	Depositor EDS
% Data completeness (in resolution range)	99.7 (24.90-1.35) 99.5 (24.90-1.35)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 1.35Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_1446)	Depositor
R, $R_{free}$	0.136 , 0.159 0.151 , 0.172	Depositor DCC
$R_{free}$ test set	1113 reflections (1.46%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.9	Xtrriage
Anisotropy	0.323	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 52.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2638	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 72.55 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.1574e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/1144	0.76	2/1553 (0.1%)
1	B	0.58	0/1260	0.74	0/1708
All	All	0.59	0/2404	0.75	2/3261 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	57[A]	LEU	CA-CB-CG	5.19	127.24	115.30
1	A	57[B]	LEU	CA-CB-CG	5.19	127.24	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1116	0	1066	9	0
1	B	1226	0	1135	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
4	A	154	0	0	0	0
4	B	139	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2638	0	2201	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64:ASN:ND2	1:B:77[B]:THR:HG22	2.09	0.66
1:A:57[A]:LEU:HD23	1:A:57[A]:LEU:C	2.25	0.56
1:A:2:MSE:HE1	1:B:83:ILE:CG2	2.36	0.56
1:B:57:LEU:HD12	1:B:57:LEU:C	2.27	0.54
1:A:2:MSE:HB3	1:A:29:GLY:HA3	1.93	0.51
1:B:68:ARG:O	1:B:90[B]:HIS:HA	2.12	0.49
1:B:51[B]:SER:HB2	1:B:87:ASP:HA	1.95	0.49
1:B:13[B]:MSE:SE	1:B:40:ILE:HD12	2.64	0.47
1:A:82[A]:ILE:HG13	1:B:4[A]:ILE:HG22	1.98	0.46
1:A:2:MSE:HE1	1:B:83:ILE:HG21	1.99	0.44
1:B:53[B]:ASN:O	1:B:86:LEU:HD22	2.18	0.44
1:A:32:THR:HG22	1:B:107:TRP:CZ2	2.54	0.42
1:A:107:TRP:CZ2	1:B:32:THR:HG22	2.55	0.41
1:B:84:ILE:HG21	1:B:90[A]:HIS:CE1	2.55	0.41
1:B:43:PRO:HD3	1:B:99:ASP:OD1	2.20	0.41
1:A:46:LYS:HE2	1:A:91:HIS:CD2	2.55	0.41
1:A:57[A]:LEU:HD21	1:A:59:ILE:HG13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/128 (102%)	131 (100%)	0	0	100	100
1	B	143/128 (112%)	137 (96%)	6 (4%)	0	100	100
All	All	274/256 (107%)	268 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/116 (108%)	122 (98%)	3 (2%)	49	14
1	B	137/116 (118%)	135 (98%)	2 (2%)	65	32
All	All	262/232 (113%)	257 (98%)	5 (2%)	69	22

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	A	57[A]	LEU
1	A	57[B]	LEU
1	B	50[A]	HIS
1	B	50[B]	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN
1	B	3	ASN
1	B	64	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	118/128 (92%)	0.51	12 (10%) <b>6</b> <b>8</b>	9, 17, 33, 51	0
1	B	117/128 (91%)	0.44	11 (9%) <b>8</b> <b>9</b>	10, 17, 35, 44	0
All	All	235/256 (91%)	0.48	23 (9%) <b>7</b> <b>9</b>	9, 17, 34, 51	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	122	ASP	5.6
1	A	48	PHE	5.5
1	A	71	ASP	4.3
1	B	53[A]	ASN	4.2
1	A	49	ARG	3.9
1	B	71[A]	ASP	3.9
1	A	53	ASN	3.6
1	B	3	ASN	3.2
1	B	51[A]	SER	3.1
1	B	105[A]	ILE	3.1
1	A	17	TYR	2.8
1	A	3	ASN	2.7
1	A	50	HIS	2.6
1	A	121	GLN	2.6
1	B	87	ASP	2.5
1	B	17[A]	TYR	2.5
1	B	120	GLU	2.4
1	A	120	GLU	2.4
1	B	121	GLN	2.3
1	A	89	GLU	2.2
1	B	52[A]	HIS	2.1
1	B	50[A]	HIS	2.0
1	A	105[A]	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NI	B	201	1/1	0.99	0.06	23,23,23,23	1
3	NA	A	202	1/1	1.00	0.08	8,8,8,8	1
2	NI	A	201	1/1	1.00	0.04	17,17,17,17	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.