



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 7, 2024 – 03:24 am GMT

PDB ID : 5MXJ  
Title : Structure of the Y108F mutant of vanillyl alcohol oxidase  
Authors : Ewing, T.A.; Nguyen, Q.-T.; Allan, R.C.; Gygli, G.; Romero, E.; Binda, C.;  
Fraaije, M.W.; Mattevi, A.; van Berkel, W.J.H.  
Deposited on : 2017-01-23  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	560	 86% 11% ..
1	B	560	 84% 12% ..

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 9081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vanillyl-alcohol oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	553	4349	2791	740	795	23	6	0	0
1	B	553	4348	2791	740	793	24	9	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	108	PHE	TYR	engineered mutation	UNP P56216
B	108	PHE	TYR	engineered mutation	UNP P56216

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	53	27	9	15	2	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0
3	B	1	6	3	3	0	0
3	B	1	6	3	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	131	131	131	0	0
4	B	129	129	129	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vanillyl-alcohol oxidase



- Molecule 1: Vanillyl-alcohol oxidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.90Å 137.90Å 132.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.91 – 2.80 54.91 – 2.80	Depositor EDS
% Data completeness (in resolution range)	88.6 (54.91-2.80) 88.6 (54.91-2.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.63 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.180 , 0.259 0.186 , 0.254	Depositor DCC
$R_{free}$ test set	1309 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.1	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 57.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h 0.014 for -l,-k,-h 0.006 for -h,-l,-k 0.000 for -h,l,k 0.021 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9081	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	1/4468 (0.0%)	0.77	3/6076 (0.0%)
1	B	0.97	2/4467 (0.0%)	0.78	4/6074 (0.1%)
All	All	0.80	3/8935 (0.0%)	0.77	7/12150 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	328	SER	CB-OG	-47.96	0.80	1.42
1	B	149	HIS	CB-CG	-22.94	1.08	1.50
1	A	328	SER	CB-OG	-11.69	1.27	1.42

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	177	LEU	CA-CB-CG	7.56	132.69	115.30
1	B	149	HIS	CA-CB-CG	7.03	125.54	113.60
1	B	177	LEU	CA-CB-CG	6.41	130.04	115.30
1	A	526	ARG	NE-CZ-NH1	6.05	123.33	120.30
1	B	312	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	A	312	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	B	183	ARG	NE-CZ-NH1	5.18	122.89	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	463	ARG	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4349	0	4250	23	0
1	B	4348	0	4253	29	1
2	A	53	0	29	0	0
2	B	53	0	29	3	0
3	A	6	0	8	0	0
3	B	12	0	16	0	0
4	A	131	0	0	0	0
4	B	129	0	0	1	0
All	All	9081	0	8585	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:ARG:NH2	1:B:247:GLY:O	2.17	0.78
1:B:448:GLN:NE2	4:B:701:HOH:O	2.19	0.74
1:A:505:THR:HG21	1:A:513:ILE:HD12	1.74	0.70
1:B:278:SER:OG	1:B:399:ASP:OD1	2.10	0.63
1:A:247:GLY:O	1:B:183:ARG:NH2	2.33	0.61
1:A:361:GLU:OE2	1:A:365:ARG:NH1	2.34	0.60
1:B:413:TRP:CZ2	2:B:601:FAD:HM72	2.40	0.57
1:B:423:LEU:HD21	1:B:488:MET:HG3	1.86	0.57
1:A:177:LEU:HD21	1:A:262:VAL:HG12	1.87	0.56
1:A:167:ASP:OD1	1:A:186:GLY:HA3	2.07	0.55
1:A:505:THR:HG21	1:A:513:ILE:CD1	2.38	0.54
1:A:151:LEU:HD23	1:A:166:LEU:HD21	1.90	0.51

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:413:TRP:CE2	2:B:601:FAD:HM72	2.46	0.51
1:A:330:ARG:NH1	1:A:338:GLU:OE1	2.43	0.51
1:B:464:GLU:OE2	1:B:466:HIS:ND1	2.43	0.50
1:A:531:LEU:HD22	1:B:531:LEU:HD22	1.93	0.49
1:A:385:PHE:HB3	1:A:386:PRO:HD2	1.95	0.48
1:B:505:THR:HG21	1:B:513:ILE:CD1	2.45	0.47
1:B:106:SER:HB3	1:B:422:HIS:HE1	1.80	0.46
1:A:10:LEU:HD22	1:A:40:ILE:HG22	1.97	0.46
1:B:205:ALA:HB2	1:B:541:ILE:HD12	1.97	0.46
1:A:296:ILE:O	1:A:297:ARG:C	2.53	0.46
1:B:151:LEU:HG	1:B:166:LEU:HD21	1.97	0.46
1:B:24:PHE:CD2	1:B:95:PHE:CE2	3.05	0.45
1:B:188:THR:HB	1:B:189:PRO:HD2	1.99	0.45
1:B:105:ASN:HB2	2:B:601:FAD:O1A	2.17	0.45
1:B:443:THR:HA	1:B:491:LEU:HD21	1.98	0.45
1:A:238:ILE:HA	1:A:241:LEU:HD12	1.99	0.44
1:A:21:PHE:CZ	1:A:25:ILE:HD12	2.52	0.44
1:A:307:ASN:HB3	1:A:358:TYR:CE1	2.52	0.44
1:A:425:PHE:CE2	1:A:427:PRO:HD3	2.53	0.44
1:A:59:ASP:OD2	1:A:506:HIS:NE2	2.48	0.44
1:A:187:TYR:HA	1:A:308:VAL:HB	2.00	0.44
1:B:358:TYR:OH	1:B:402:MET:O	2.24	0.43
1:B:312:ARG:NH2	1:B:410:GLU:OE1	2.51	0.43
1:B:428:ILE:HD11	1:B:503:TYR:HB3	2.00	0.43
1:A:114:ARG:NH1	1:A:511:ASP:OD2	2.52	0.42
1:B:425:PHE:CE2	1:B:491:LEU:HB3	2.55	0.42
1:B:474:ASN:N	1:B:480:GLN:OE1	2.49	0.42
1:B:321:LEU:HD13	1:B:346:LEU:HD22	2.02	0.42
1:B:416:TRP:CD1	1:B:417:LEU:HD13	2.55	0.41
1:B:425:PHE:CZ	1:B:491:LEU:HB3	2.55	0.41
1:A:188:THR:HB	1:A:189:PRO:CD	2.51	0.41
1:B:114:ARG:NH1	1:B:511:ASP:OD2	2.54	0.41
1:B:61:HIS:CE1	1:B:422:HIS:CE1	3.09	0.41
1:A:165:TRP:CZ3	1:A:403:GLN:HB2	2.56	0.40
1:B:98:TRP:CD2	1:B:113:PRO:HA	2.56	0.40
1:A:242:PHE:HB2	1:B:464:GLU:HG3	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:337:GLU:OE1	1:B:337:GLU:OE1[2_455]	1.82	0.38

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/560 (98%)	519 (94%)	28 (5%)	2 (0%)	34	66
1	B	549/560 (98%)	513 (93%)	34 (6%)	2 (0%)	34	66
All	All	1098/1120 (98%)	1032 (94%)	62 (6%)	4 (0%)	34	66

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	274	ARG
1	B	199	GLY
1	A	390	PRO
1	B	284	PRO

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	467/482 (97%)	434 (93%)	33 (7%)	14	39
1	B	467/482 (97%)	426 (91%)	41 (9%)	10	29
All	All	934/964 (97%)	860 (92%)	74 (8%)	12	34

All (74) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LEU
1	A	16	LEU
1	A	17	SER
1	A	19	SER
1	A	25	ILE
1	A	64	MET
1	A	95	PHE
1	A	114	ARG
1	A	159	ASN
1	A	177	LEU
1	A	183	ARG
1	A	195	MET
1	A	222	ARG
1	A	236	SER
1	A	251	ASP
1	A	297	ARG
1	A	301	LEU
1	A	310	THR
1	A	342	ILE
1	A	348	LEU
1	A	370	THR
1	A	381	VAL
1	A	384	TYR
1	A	388	ASP
1	A	409	ASP
1	A	422	HIS
1	A	452	LEU
1	A	464	GLU
1	A	466	HIS
1	A	488	MET
1	A	505	THR
1	A	515	GLU
1	A	552	GLN
1	B	15	LYS
1	B	16	LEU
1	B	35	GLU
1	B	41	SER
1	B	42	SER
1	B	52	MET
1	B	63	VAL
1	B	64	MET
1	B	78	ARG

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	B	114	ARG
1	B	125	LYS
1	B	162	ASP
1	B	177	LEU
1	B	189	PRO
1	B	197	HIS
1	B	217	LEU
1	B	224	GLU
1	B	230	PRO
1	B	251	ASP
1	B	297	ARG
1	B	301	LEU
1	B	314	ILE
1	B	321	LEU
1	B	329	SER
1	B	332	GLU
1	B	336	ASP
1	B	350	ARG
1	B	361	GLU
1	B	381	VAL
1	B	387	GLU
1	B	409	ASP
1	B	422	HIS
1	B	434	GLU
1	B	448	GLN
1	B	464	GLU
1	B	488	MET
1	B	489	ARG
1	B	503	TYR
1	B	505	THR
1	B	552	GLN
1	B	555	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	B	158	ASN
1	B	485	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	603	-	5,5,5	0.61	0	5,5,5	0.35	0
2	FAD	B	601	1	53,58,58	1.46	10 (18%)	68,89,89	1.46	13 (19%)
3	GOL	A	602	-	5,5,5	0.33	0	5,5,5	0.49	0
2	FAD	A	601	1	53,58,58	1.36	6 (11%)	68,89,89	1.38	12 (17%)
3	GOL	B	602	-	5,5,5	0.38	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	603	-	-	2/4/4/4	-
2	FAD	B	601	1	-	6/30/50/50	0/6/6/6
3	GOL	A	602	-	-	0/4/4/4	-
2	FAD	A	601	1	-	14/30/50/50	0/6/6/6

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	602	-	-	2/4/4/4	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C9A-C5X	5.36	1.50	1.41
2	B	601	FAD	C9A-C5X	5.31	1.50	1.41
2	A	601	FAD	C8-C7	3.69	1.50	1.40
2	B	601	FAD	C8-C7	3.48	1.49	1.40
2	B	601	FAD	C1'-C2'	2.93	1.56	1.52
2	B	601	FAD	C5A-C4A	2.77	1.48	1.40
2	B	601	FAD	C4-N3	-2.66	1.33	1.38
2	A	601	FAD	C5A-C4A	2.41	1.47	1.40
2	A	601	FAD	C1'-C2'	2.39	1.56	1.52
2	B	601	FAD	C4X-N5	2.31	1.35	1.30
2	A	601	FAD	C4X-N5	2.27	1.35	1.30
2	B	601	FAD	C5A-N7A	-2.19	1.31	1.39
2	B	601	FAD	C2A-N3A	2.15	1.35	1.32
2	B	601	FAD	O4B-C1B	2.06	1.44	1.41
2	B	601	FAD	C10-N10	2.04	1.41	1.37
2	A	601	FAD	C4X-C10	2.04	1.50	1.44

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	FAD	C4-C4X-N5	3.39	123.06	118.23
2	B	601	FAD	N3A-C2A-N1A	-3.32	123.48	128.68
2	B	601	FAD	P-O3P-PA	-3.17	121.95	132.83
2	A	601	FAD	N3A-C2A-N1A	-2.97	124.04	128.68
2	B	601	FAD	C4X-C10-N1	-2.94	117.90	124.73
2	A	601	FAD	C8M-C8-C9	-2.91	114.11	119.49
2	A	601	FAD	C8M-C8-C7	2.82	126.52	120.74
2	B	601	FAD	N6A-C6A-N1A	2.69	124.17	118.57
2	B	601	FAD	O2'-C2'-C3'	-2.67	102.60	109.10
2	A	601	FAD	C4X-C10-N1	-2.67	118.53	124.73
2	A	601	FAD	C10-N1-C2	2.49	121.88	116.90
2	A	601	FAD	C4A-C5A-N7A	-2.42	106.88	109.40
2	A	601	FAD	O2-C2-N1	-2.41	117.83	121.83
2	B	601	FAD	C10-N1-C2	2.41	121.72	116.90
2	A	601	FAD	O4-C4-C4X	-2.40	120.22	126.60
2	A	601	FAD	O2'-C2'-C1'	2.33	115.42	109.80
2	B	601	FAD	C4X-C4-N3	2.32	119.09	113.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	FAD	O2'-C2'-C1'	2.26	115.27	109.80
2	A	601	FAD	C4-C4X-N5	2.25	121.43	118.23
2	A	601	FAD	C4X-C10-N10	2.21	119.71	116.48
2	B	601	FAD	C4X-C10-N10	2.20	119.69	116.48
2	B	601	FAD	C5X-N5-C4X	2.18	121.70	118.07
2	B	601	FAD	C10-C4X-N5	-2.07	120.46	124.86
2	A	601	FAD	C2A-N1A-C6A	2.07	122.30	118.75
2	B	601	FAD	C4-N3-C2	-2.02	121.92	125.64

There are no chirality outliers.

All (24) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	FAD	C5B-O5B-PA-O3P
2	A	601	FAD	C3'-C4'-C5'-O5'
2	A	601	FAD	O4'-C4'-C5'-O5'
2	B	601	FAD	N10-C1'-C2'-C3'
3	B	602	GOL	O1-C1-C2-C3
2	A	601	FAD	O4B-C4B-C5B-O5B
2	A	601	FAD	C3B-C4B-C5B-O5B
3	B	603	GOL	O1-C1-C2-C3
2	B	601	FAD	O4'-C4'-C5'-O5'
3	B	602	GOL	O1-C1-C2-O2
2	B	601	FAD	C4'-C5'-O5'-P
2	B	601	FAD	C3'-C4'-C5'-O5'
2	A	601	FAD	C2'-C3'-C4'-O4'
2	A	601	FAD	C4'-C5'-O5'-P
2	A	601	FAD	C2'-C3'-C4'-C5'
2	A	601	FAD	C5B-O5B-PA-O2A
2	A	601	FAD	N10-C1'-C2'-C3'
2	B	601	FAD	N10-C1'-C2'-O2'
3	B	603	GOL	O1-C1-C2-O2
2	A	601	FAD	PA-O3P-P-O1P
2	A	601	FAD	PA-O3P-P-O2P
2	A	601	FAD	C5B-O5B-PA-O1A
2	A	601	FAD	C1'-C2'-C3'-O3'
2	B	601	FAD	C1'-C2'-C3'-O3'

There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	FAD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	553/560 (98%)	-0.14	7 (1%) 77 72	31, 56, 78, 101	2 (0%)
1	B	553/560 (98%)	-0.13	7 (1%) 77 72	32, 48, 71, 86	3 (0%)
All	All	1106/1120 (98%)	-0.13	14 (1%) 77 72	31, 51, 75, 101	5 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	26	GLN	4.7
1	B	19	SER	2.6
1	B	448	GLN	2.6
1	A	158	ASN	2.6
1	A	159	ASN	2.4
1	B	331	THR	2.4
1	A	72	SER	2.2
1	A	7	PHE	2.2
1	B	329	SER	2.2
1	A	16	LEU	2.1
1	A	19	SER	2.1
1	B	345	GLN	2.0
1	B	6	GLU	2.0
1	B	332	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	603	6/6	0.88	0.40	67,70,73,73	0
3	GOL	B	602	6/6	0.90	0.19	57,58,59,60	0
3	GOL	A	602	6/6	0.91	0.24	52,57,58,60	0
2	FAD	A	601	53/53	0.97	0.14	45,59,69,71	0
2	FAD	B	601	53/53	0.98	0.19	30,35,45,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.