



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 27, 2024 – 08:05 PM EDT

PDB ID : 1N33
Title : Structure of the Thermus thermophilus 30S ribosomal subunit bound to codon and near-cognate transfer rna anticodon stem-loop mismatched at the second codon position at the a site with paromomycin
Authors : Ogle, J.M.; Murphy IV, F.V.; Tarry, M.J.; Ramakrishnan, V.
Deposited on : 2002-10-25
Resolution : 3.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.























Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1012 (3.40-3.32)
Clashscore	180529	1035 (3.40-3.32)
Ramachandran outliers	177936	1037 (3.40-3.32)
Sidechain outliers	177891	1037 (3.40-3.32)
RSRZ outliers	164620	1012 (3.40-3.32)
RNA backbone	3690	1010 (3.74-2.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1522	
2	Y	17	
3	Z	6	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	B	256	
5	C	239	
6	D	208	
7	E	161	
8	F	101	
9	G	155	
10	H	138	
11	I	128	
12	J	104	
13	K	129	
14	L	135	
15	M	126	
16	N	60	
17	O	88	
18	P	88	
19	Q	104	
20	R	88	
21	S	92	
22	T	106	
23	V	26	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
25	MG	A	441	-	-	-	X

2 Entry composition [i](#)

There are 26 unique types of molecules in this entry. The entry contains 52140 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 16S RIBOSOMAL RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	1513	32508	14472	6016	10509	1511	22	0	0

- Molecule 2 is a RNA chain called ANTICODON STEM-LOOP OF SER TRANSFER RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	Y	11	235	106	45	74	10	0	0	0

- Molecule 3 is a RNA chain called A-SITE MESSENGER RNA FRAGMENT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	Z	4	77	36	8	30	3	0	0	0

- Molecule 4 is a protein called 30S RIBOSOMAL PROTEIN S2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	B	234	1900	1213	341	341	5	0	0	0

- Molecule 5 is a protein called 30S RIBOSOMAL PROTEIN S3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	C	206	1612	1016	314	281	1	0	0	0

- Molecule 6 is a protein called 30S RIBOSOMAL PROTEIN S4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	D	208	1703	1066	339	291	7	0	0	0

- Molecule 7 is a protein called 30S RIBOSOMAL PROTEIN S5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	E	150	1146	724	217	201	4	0	0	0

- Molecule 8 is a protein called 30S RIBOSOMAL PROTEIN S6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	F	101	843	531	155	154	3	0	0	0

- Molecule 9 is a protein called 30S RIBOSOMAL PROTEIN S7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	G	155	1257	781	252	218	6	0	0	0

- Molecule 10 is a protein called 30S RIBOSOMAL PROTEIN S8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	H	138	1116	705	215	193	3	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	25	ASP	GLU	conflict	UNP P24319
H	37	ARG	LYS	conflict	UNP P24319
H	52	ASP	GLU	conflict	UNP P24319
H	61	VAL	ILE	conflict	UNP P24319
H	62	TYR	HIS	conflict	UNP P24319
H	81	HIS	LYS	conflict	UNP P24319
H	88	LYS	ARG	conflict	UNP P24319
H	115	SER	PRO	conflict	UNP P24319

- Molecule 11 is a protein called 30S RIBOSOMAL PROTEIN S9.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
11	I	127	1011	639	198	174	0	0	0

- Molecule 12 is a protein called 30S RIBOSOMAL PROTEIN S10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
12	J	98	792	498	156	137	1	0	0	0

- Molecule 13 is a protein called 30S RIBOSOMAL PROTEIN S11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
13	K	119	885	549	168	165	3	0	0	0

- Molecule 14 is a protein called 30S RIBOSOMAL PROTEIN S12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
14	L	124	970	611	195	163	1	0	0	0

- Molecule 15 is a protein called 30S RIBOSOMAL PROTEIN S13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
15	M	118	937	579	193	163	2	0	0	0

- Molecule 16 is a protein called 30S RIBOSOMAL PROTEIN S14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
16	N	60	492	312	104	72	4	0	0	0

- Molecule 17 is a protein called 30S RIBOSOMAL PROTEIN S15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
17	O	88	734	459	147	126	2	0	0	0

- Molecule 18 is a protein called 30S RIBOSOMAL PROTEIN S16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
18	P	83	700	443	139	117	1	0	0	0

- Molecule 19 is a protein called 30S RIBOSOMAL PROTEIN S17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
19	Q	104	Total	C	N	O	S	0	0	0
			857	547	161	147	2			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Q	50	LYS	ARG	conflict	UNP P24321
Q	53	LEU	VAL	conflict	UNP P24321
Q	62	SER	ALA	conflict	UNP P24321
Q	79	SER	GLU	conflict	UNP P24321
Q	82	MET	LEU	conflict	UNP P24321
Q	90	ILE	VAL	conflict	UNP P24321
Q	96	GLN	ALA	conflict	UNP P24321

- Molecule 20 is a protein called 30S RIBOSOMAL PROTEIN S18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
20	R	73	Total	C	N	O	S	0	0	0
			597	380	118	99				

- Molecule 21 is a protein called 30S RIBOSOMAL PROTEIN S19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
21	S	80	Total	C	N	O	S	0	0	0
			647	414	119	112	2			


- Molecule 22 is a protein called 30S RIBOSOMAL PROTEIN S20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
22	T	99	Total	C	N	O	S	0	0	0
			763	470	162	129	2			

- Molecule 23 is a protein called 30S RIBOSOMAL PROTEIN THX.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
23	V	24	Total	C	N	O		0	0	0
			208	128	50	30				

- Molecule 24 is PAROMOMYCIN (three-letter code: PAR) (formula: C₂₃H₄₅N₅O₁₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
24	A	1	42	23	5	14	0	0

- Molecule 25 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
25	A	103	103	103	0	0
25	Y	1	1	1	0	0
25	Z	2	2	2	0	0


- Molecule 26 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
26	D	1	1	1	0	0
26	N	1	1	1	0	0

3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 16S RIBOSOMAL RNA










● Molecule 2: ANTICODON STEM-LOOP OF SER TRANSFER RNA








- Molecule 3: A-SITE MESSENGER RNA FRAGMENT




- Molecule 4: 30S RIBOSOMAL PROTEIN S2




- Molecule 5: 30S RIBOSOMAL PROTEIN S3




- Molecule 6: 30S RIBOSOMAL PROTEIN S4




- Molecule 7: 30S RIBOSOMAL PROTEIN S5




- Molecule 8: 30S RIBOSOMAL PROTEIN S6




- Molecule 9: 30S RIBOSOMAL PROTEIN S7








● Molecule 10: 30S RIBOSOMAL PROTEIN S8




● Molecule 11: 30S RIBOSOMAL PROTEIN S9




● Molecule 12: 30S RIBOSOMAL PROTEIN S10



● Molecule 13: 30S RIBOSOMAL PROTEIN S11








● Molecule 14: 30S RIBOSOMAL PROTEIN S12



● Molecule 15: 30S RIBOSOMAL PROTEIN S13





● Molecule 16: 30S RIBOSOMAL PROTEIN S14




● Molecule 17: 30S RIBOSOMAL PROTEIN S15





● Molecule 18: 30S RIBOSOMAL PROTEIN S16





- Molecule 19: 30S RIBOSOMAL PROTEIN S17



- Molecule 20: 30S RIBOSOMAL PROTEIN S18



- Molecule 21: 30S RIBOSOMAL PROTEIN S19




- Molecule 22: 30S RIBOSOMAL PROTEIN S20





• Molecule 23: 30S RIBOSOMAL PROTEIN THX



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	401.60Å 401.60Å 176.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.54 – 3.35 74.54 – 3.35	Depositor EDS
% Data completeness (in resolution range)	88.6 (74.54-3.35) 88.6 (74.54-3.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 3.33Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.225 , 0.284 0.230 , 0.281	Depositor DCC
R_{free} test set	9128 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	104.3	Xtrriage
Anisotropy	0.103	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 339.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	52140	wwPDB-VP
Average B, all atoms (Å ²)	101.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PSU, PAR, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.58	1/36387 (0.0%)	0.76	33/56789 (0.1%)
2	Y	0.59	0/241	0.92	2/375 (0.5%)
3	Z	0.48	0/84	0.78	0/128
4	B	0.36	0/1935	0.66	0/2609
5	C	0.38	0/1636	0.65	0/2205
6	D	0.42	0/1733	0.68	0/2318
7	E	0.48	0/1162	0.77	0/1564
8	F	0.37	0/856	0.66	0/1154
9	G	0.35	0/1276	0.60	0/1709
10	H	0.53	0/1136	0.83	1/1527 (0.1%)
11	I	0.35	0/1029	0.64	0/1378
12	J	0.38	0/805	0.71	1/1082 (0.1%)
13	K	0.41	0/900	0.69	0/1213
14	L	0.44	0/986	0.79	0/1320
15	M	0.35	0/947	0.61	0/1270
16	N	0.39	0/501	0.66	0/664
17	O	0.41	0/745	0.63	0/992
18	P	0.53	0/716	0.84	0/963
19	Q	0.55	0/870	0.80	0/1159
20	R	0.39	0/603	0.68	0/799
21	S	0.35	0/661	0.68	0/890
22	T	0.45	0/765	0.81	2/1007 (0.2%)
23	V	0.37	0/212	0.59	0/277
All	All	0.53	1/56186 (0.0%)	0.74	39/83392 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	3	46
10	H	0	1
All	All	3	47

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	758	G	C5-C6	-5.26	1.37	1.42

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	A	C2'-C3'-O3'	10.26	132.07	109.50
1	A	366	C	C2'-C3'-O3'	8.95	129.19	109.50
1	A	1498	U	C2'-C3'-O3'	8.91	129.10	109.50
1	A	1528	U	C2'-C3'-O3'	8.77	128.80	109.50
1	A	181	G	C2'-C3'-O3'	8.66	128.56	109.50

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	243	A	C3'
1	A	1504	G	C3'
1	A	1528	U	C3'

5 of 47 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	106	C	Sidechain
1	A	115	G	Sidechain
1	A	190(L)	U	Sidechain
1	A	51	A	Sidechain
1	A	54	C	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	32508	0	16414	1353	0
2	Y	235	0	121	11	0
3	Z	77	0	42	1	0
4	B	1900	0	1951	303	0
5	C	1612	0	1677	285	0
6	D	1703	0	1764	212	0
7	E	1146	0	1207	129	0
8	F	843	0	857	94	0
9	G	1257	0	1296	132	0
10	H	1116	0	1177	143	0
11	I	1011	0	1043	177	0
12	J	792	0	835	169	2
13	K	885	0	904	103	0
14	L	970	0	1057	138	0
15	M	937	0	995	121	0
16	N	492	0	531	89	0
17	O	734	0	771	78	0
18	P	700	0	720	86	0
19	Q	857	0	930	130	0
20	R	597	0	668	93	0
21	S	647	0	673	94	0
22	T	763	0	861	96	0
23	V	208	0	221	37	0
24	A	42	0	45	4	0
25	A	103	0	0	0	0
25	Y	1	0	0	0	0
25	Z	2	0	0	0	0
26	D	1	0	0	0	0
26	N	1	0	0	0	0
All	All	52140	0	36760	3828	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 3828 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:D:36:ARG:H	6:D:37:PRO:HD3	1.05	1.11
1:A:1443:G:H5''	1:A:1446:A:H5''	1.28	1.09
1:A:1356:G:H2'	1:A:1357:A:C8	1.90	1.06
6:D:63:LYS:HD2	6:D:198:VAL:HG23	1.34	1.06
1:A:1250:A:H4'	11:I:68:GLY:H	1.18	1.05

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
12:J:79:ARG:NH1	12:J:79:ARG:NH1[8_665]	1.67	0.53
12:J:80:LYS:NZ	12:J:80:LYS:NZ[8_665]	1.96	0.24

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	B	232/256 (91%)	132 (57%)	57 (25%)	43 (18%)	0	0
5	C	204/239 (85%)	115 (56%)	46 (22%)	43 (21%)	0	0
6	D	206/208 (99%)	145 (70%)	46 (22%)	15 (7%)	1	5
7	E	148/161 (92%)	115 (78%)	20 (14%)	13 (9%)	0	3
8	F	99/101 (98%)	73 (74%)	21 (21%)	5 (5%)	1	10
9	G	153/155 (99%)	93 (61%)	39 (26%)	21 (14%)	0	0
10	H	136/138 (99%)	102 (75%)	24 (18%)	10 (7%)	1	5
11	I	125/128 (98%)	70 (56%)	37 (30%)	18 (14%)	0	0
12	J	96/104 (92%)	54 (56%)	22 (23%)	20 (21%)	0	0
13	K	117/129 (91%)	77 (66%)	29 (25%)	11 (9%)	0	3
14	L	122/135 (90%)	81 (66%)	23 (19%)	18 (15%)	0	0
15	M	116/126 (92%)	63 (54%)	35 (30%)	18 (16%)	0	0
16	N	58/60 (97%)	34 (59%)	14 (24%)	10 (17%)	0	0
17	O	86/88 (98%)	57 (66%)	26 (30%)	3 (4%)	3	16
18	P	81/88 (92%)	52 (64%)	24 (30%)	5 (6%)	1	8
19	Q	102/104 (98%)	81 (79%)	11 (11%)	10 (10%)	0	3
20	R	71/88 (81%)	47 (66%)	14 (20%)	10 (14%)	0	0
21	S	78/92 (85%)	46 (59%)	21 (27%)	11 (14%)	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
22	T	97/106 (92%)	51 (53%)	33 (34%)	13 (13%)	0	0
23	V	22/26 (85%)	16 (73%)	4 (18%)	2 (9%)	0	3
All	All	2349/2532 (93%)	1504 (64%)	546 (23%)	299 (13%)	0	1

5 of 299 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	B	8	LYS
4	B	9	GLU
4	B	15	VAL
4	B	16	HIS
4	B	21	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	B	202/220 (92%)	177 (88%)	25 (12%)	4	14
5	C	160/188 (85%)	149 (93%)	11 (7%)	13	38
6	D	180/180 (100%)	160 (89%)	20 (11%)	5	19
7	E	115/122 (94%)	101 (88%)	14 (12%)	4	15
8	F	90/90 (100%)	81 (90%)	9 (10%)	6	23
9	G	126/126 (100%)	118 (94%)	8 (6%)	15	40
10	H	119/119 (100%)	101 (85%)	18 (15%)	2	9
11	I	98/99 (99%)	86 (88%)	12 (12%)	4	15
12	J	87/91 (96%)	73 (84%)	14 (16%)	2	8
13	K	90/99 (91%)	83 (92%)	7 (8%)	10	33
14	L	104/111 (94%)	91 (88%)	13 (12%)	3	14
15	M	94/101 (93%)	88 (94%)	6 (6%)	14	40
16	N	49/49 (100%)	44 (90%)	5 (10%)	6	22
17	O	79/79 (100%)	73 (92%)	6 (8%)	11	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
18	P	72/74 (97%)	62 (86%)	10 (14%)	3	11
19	Q	96/96 (100%)	86 (90%)	10 (10%)	5	21
20	R	64/77 (83%)	61 (95%)	3 (5%)	22	49
21	S	71/79 (90%)	63 (89%)	8 (11%)	4	18
22	T	76/82 (93%)	67 (88%)	9 (12%)	4	16
23	V	19/21 (90%)	18 (95%)	1 (5%)	19	45
All	All	1991/2103 (95%)	1782 (90%)	209 (10%)	5	21

5 of 209 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
11	I	121	ARG
14	L	67	THR
22	T	8	ARG
12	J	13	HIS
12	J	98	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
9	G	106	GLN
14	L	75	HIS
21	S	56	GLN
9	G	122	HIS
12	J	76	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	1511/1522 (99%)	206 (13%)	66 (4%)
2	Y	10/17 (58%)	2 (20%)	0
3	Z	3/6 (50%)	0	0
All	All	1524/1545 (98%)	208 (13%)	66 (4%)

5 of 208 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	8	A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	9	G
1	A	31	G
1	A	32	A
1	A	39	G

5 of 66 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	1281	U
1	A	1347	G
1	A	1528	U
1	A	429	U
1	A	428	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PSU	Y	40	2	18,21,22	2.03	6 (33%)	21,30,33	1.29	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PSU	Y	40	2	-	5/7/25/26	0/2/2/2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Y	40	PSU	C2-N1	5.11	1.43	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Y	40	PSU	O3'-C3'	3.47	1.51	1.43
2	Y	40	PSU	O4'-C1'	-2.58	1.40	1.43
2	Y	40	PSU	C6-N1	2.43	1.40	1.36
2	Y	40	PSU	C4-N3	2.06	1.42	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Y	40	PSU	O2-C2-N1	2.78	125.65	122.79
2	Y	40	PSU	C6-C5-C4	2.16	119.63	118.17
2	Y	40	PSU	O4'-C4'-C3'	-2.10	100.98	105.15

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Y	40	PSU	C2'-C1'-C5-C4
2	Y	40	PSU	C2'-C1'-C5-C6
2	Y	40	PSU	O4'-C4'-C5'-O5'
2	Y	40	PSU	C3'-C4'-C5'-O5'
2	Y	40	PSU	C4'-C5'-O5'-P

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Y	40	PSU	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 109 ligands modelled in this entry, 108 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
24	PAR	A	1545	-	44,45,45	2.06	13 (29%)	63,67,67	1.02	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
24	PAR	A	1545	-	-	1/18/94/94	0/4/4/4

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
24	A	1545	PAR	C31-C21	6.48	1.61	1.53
24	A	1545	PAR	C34-C24	4.47	1.59	1.53
24	A	1545	PAR	O43-C13	4.23	1.49	1.41
24	A	1545	PAR	O54-C14	3.55	1.51	1.41
24	A	1545	PAR	C44-C54	3.20	1.59	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
24	A	1545	PAR	O11-C11-C21	2.83	112.71	108.08
24	A	1545	PAR	O54-C54-C64	2.44	110.77	106.07

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
24	A	1545	PAR	C24-C14-O33-C33

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
24	A	1545	PAR	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1512/1522 (99%)	0.76	196 (12%) 9 11	25, 86, 178, 198	0
2	Y	10/17 (58%)	1.17	1 (10%) 14 16	91, 117, 190, 194	0
3	Z	4/6 (66%)	0.81	1 (25%) 2 4	87, 92, 96, 197	0
4	B	234/256 (91%)	1.03	38 (16%) 5 8	22, 132, 191, 198	0
5	C	206/239 (86%)	0.83	26 (12%) 9 12	55, 126, 183, 198	0
6	D	208/208 (100%)	0.57	25 (12%) 10 13	16, 91, 157, 197	0
7	E	150/161 (93%)	0.49	8 (5%) 33 29	31, 83, 138, 198	0
8	F	101/101 (100%)	0.60	8 (7%) 20 20	56, 117, 174, 194	0
9	G	155/155 (100%)	0.90	17 (10%) 12 14	64, 127, 175, 198	0
10	H	138/138 (100%)	0.23	7 (5%) 34 30	4, 65, 145, 197	0
11	I	127/128 (99%)	1.27	30 (23%) 2 4	57, 140, 187, 198	0
12	J	98/104 (94%)	1.32	21 (21%) 3 5	51, 146, 197, 198	0
13	K	119/129 (92%)	0.89	18 (15%) 6 9	48, 99, 169, 198	0
14	L	124/135 (91%)	0.60	15 (12%) 10 12	37, 88, 157, 181	0
15	M	118/126 (93%)	1.19	20 (16%) 5 7	60, 120, 175, 196	0
16	N	60/60 (100%)	1.32	12 (20%) 3 6	60, 117, 174, 198	0
17	O	88/88 (100%)	0.46	7 (7%) 20 20	21, 83, 147, 198	0
18	P	83/88 (94%)	0.28	3 (3%) 46 38	10, 62, 121, 172	0
19	Q	104/104 (100%)	0.55	8 (7%) 21 21	14, 75, 157, 198	0
20	R	73/88 (82%)	0.40	4 (5%) 32 29	46, 99, 178, 194	0
21	S	80/92 (86%)	1.21	15 (18%) 4 6	74, 138, 191, 198	0
22	T	99/106 (93%)	0.48	8 (8%) 19 20	24, 70, 136, 186	0
23	V	24/26 (92%)	1.71	6 (25%) 2 4	44, 122, 183, 198	0
All	All	3915/4077 (96%)	0.78	494 (12%) 9 12	4, 98, 179, 198	0

The worst 5 of 494 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
21	S	3	ARG	15.0
21	S	2	PRO	9.7
11	I	128	ARG	9.3
12	J	58	ASP	7.8
1	A	1129	C	7.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PSU	Y	40	20/21	0.90	0.32	25,25,25,25	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
25	MG	A	1575	1/1	0.66	0.30	34,34,34,34	1
25	MG	A	1597	1/1	0.71	0.16	25,25,25,25	1
25	MG	A	441	1/1	0.73	0.55	34,34,34,34	1
25	MG	A	1584	1/1	0.77	0.31	34,34,34,34	0
25	MG	A	1620	1/1	0.78	0.25	34,34,34,34	1
25	MG	A	1609	1/1	0.81	1.08	34,34,34,34	1
25	MG	A	1588	1/1	0.81	0.51	34,34,34,34	0
25	MG	A	1606	1/1	0.81	0.52	34,34,34,34	1
25	MG	A	210	1/1	0.82	0.13	34,34,34,34	1
25	MG	A	1622	1/1	0.82	0.43	34,34,34,34	1
25	MG	A	1601	1/1	0.83	0.19	34,34,34,34	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
25	MG	A	1594	1/1	0.83	0.27	34,34,34,34	0
25	MG	A	1562	1/1	0.83	0.40	34,34,34,34	0
25	MG	A	1585	1/1	0.84	0.46	34,34,34,34	1
25	MG	A	1626	1/1	0.84	0.12	34,34,34,34	1
25	MG	A	473	1/1	0.84	0.19	34,34,34,34	1
25	MG	A	1618	1/1	0.85	0.35	34,34,34,34	1
25	MG	A	1599	1/1	0.85	0.15	34,34,34,34	0
25	MG	A	1615	1/1	0.85	0.37	34,34,34,34	1
25	MG	A	1548	1/1	0.86	0.31	34,34,34,34	1
25	MG	A	1561	1/1	0.86	0.10	34,34,34,34	1
25	MG	A	1603	1/1	0.86	0.22	34,34,34,34	1
25	MG	A	1549	1/1	0.87	0.23	34,34,34,34	0
25	MG	A	1610	1/1	0.87	0.09	34,34,34,34	0
25	MG	A	466	1/1	0.87	0.25	34,34,34,34	1
25	MG	A	214	1/1	0.87	0.48	34,34,34,34	1
25	MG	A	1572	1/1	0.88	0.21	34,34,34,34	0
25	MG	A	1629	1/1	0.88	0.30	34,34,34,34	0
25	MG	A	1630	1/1	0.88	0.34	34,34,34,34	0
25	MG	A	1574	1/1	0.88	0.26	34,34,34,34	0
25	MG	A	1590	1/1	0.88	0.24	34,34,34,34	0
25	MG	A	1633	1/1	0.88	0.26	34,34,34,34	1
25	MG	Z	501	1/1	0.88	0.14	34,34,34,34	1
25	MG	A	1616	1/1	0.89	0.18	34,34,34,34	1
25	MG	A	469	1/1	0.89	0.18	34,34,34,34	1
25	MG	A	1580	1/1	0.89	0.30	34,34,34,34	0
25	MG	A	1586	1/1	0.89	0.31	34,34,34,34	0
25	MG	Y	500	1/1	0.89	0.14	25,25,25,25	1
25	MG	A	1560	1/1	0.89	0.31	34,34,34,34	0
25	MG	A	1550	1/1	0.90	0.12	34,34,34,34	1
25	MG	A	1627	1/1	0.90	0.23	34,34,34,34	1
25	MG	A	1557	1/1	0.90	0.21	34,34,34,34	0
24	PAR	A	1545	42/42	0.90	0.10	30,30,30,30	0
25	MG	A	1624	1/1	0.90	0.34	34,34,34,34	0
25	MG	A	1547	1/1	0.91	0.26	34,34,34,34	0
25	MG	A	1551	1/1	0.91	0.45	34,34,34,34	0
25	MG	A	87	1/1	0.91	0.23	34,34,34,34	0
25	MG	A	1595	1/1	0.91	0.24	34,34,34,34	1
25	MG	A	1614	1/1	0.92	0.27	34,34,34,34	1
25	MG	A	1567	1/1	0.92	0.32	34,34,34,34	0
25	MG	A	1587	1/1	0.92	0.26	34,34,34,34	0
25	MG	A	1617	1/1	0.92	0.14	34,34,34,34	0
25	MG	A	1605	1/1	0.92	0.21	34,34,34,34	0

Continued on next page...

Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
25	MG	A	1559	1/1	0.92	0.27	34,34,34,34	0
25	MG	A	471	1/1	0.92	0.33	34,34,34,34	1
25	MG	A	1607	1/1	0.92	0.77	34,34,34,34	1
25	MG	A	1556	1/1	0.92	0.37	34,34,34,34	0
25	MG	A	1623	1/1	0.92	0.12	34,34,34,34	1
25	MG	Z	400	1/1	0.92	0.17	34,34,34,34	1
25	MG	A	1565	1/1	0.92	0.23	34,34,34,34	1
25	MG	A	1596	1/1	0.93	0.19	34,34,34,34	1
25	MG	A	1591	1/1	0.93	0.15	34,34,34,34	0
25	MG	A	1576	1/1	0.93	0.21	34,34,34,34	1
25	MG	A	1558	1/1	0.93	0.28	34,34,34,34	0
25	MG	A	493	1/1	0.93	0.24	34,34,34,34	1
25	MG	A	1552	1/1	0.93	0.25	34,34,34,34	0
25	MG	A	1621	1/1	0.93	0.27	34,34,34,34	1
25	MG	A	1568	1/1	0.93	0.10	34,34,34,34	0
25	MG	A	1631	1/1	0.94	0.06	34,34,34,34	1
25	MG	A	1632	1/1	0.94	0.08	34,34,34,34	1
25	MG	A	1619	1/1	0.94	0.07	34,34,34,34	1
25	MG	A	1592	1/1	0.94	0.07	34,34,34,34	0
25	MG	A	1581	1/1	0.94	0.35	34,34,34,34	0
25	MG	A	1612	1/1	0.94	0.31	34,34,34,34	0
25	MG	A	1604	1/1	0.94	0.17	34,34,34,34	0
25	MG	A	1573	1/1	0.95	0.21	34,34,34,34	0
25	MG	A	1555	1/1	0.95	0.34	34,34,34,34	0
25	MG	A	1589	1/1	0.95	0.33	34,34,34,34	0
25	MG	A	1582	1/1	0.95	0.27	34,34,34,34	0
25	MG	A	1553	1/1	0.95	0.28	34,34,34,34	0
25	MG	A	86	1/1	0.95	0.17	34,34,34,34	0
25	MG	A	1577	1/1	0.95	0.17	34,34,34,34	1
25	MG	A	1628	1/1	0.96	0.25	34,34,34,34	0
25	MG	A	1600	1/1	0.96	0.13	34,34,34,34	1
25	MG	A	1579	1/1	0.96	0.17	34,34,34,34	0
25	MG	A	211	1/1	0.96	0.52	34,34,34,34	1
25	MG	A	1564	1/1	0.96	0.33	34,34,34,34	0
25	MG	A	1593	1/1	0.96	0.17	34,34,34,34	0
25	MG	A	1546	1/1	0.96	0.24	34,34,34,34	0
25	MG	A	1566	1/1	0.96	0.29	34,34,34,34	0
25	MG	A	1608	1/1	0.96	0.24	34,34,34,34	0
25	MG	A	1583	1/1	0.96	0.21	34,34,34,34	0
25	MG	A	1571	1/1	0.96	0.08	34,34,34,34	0
25	MG	A	1611	1/1	0.96	0.17	34,34,34,34	0
25	MG	A	1598	1/1	0.96	0.08	34,34,34,34	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
25	MG	A	71	1/1	0.96	0.10	34,34,34,34	0
25	MG	A	1569	1/1	0.97	0.28	34,34,34,34	0
25	MG	A	1570	1/1	0.97	0.12	34,34,34,34	0
25	MG	A	1602	1/1	0.97	0.16	34,34,34,34	1
25	MG	A	1634	1/1	0.97	0.28	34,34,34,34	1
25	MG	A	1613	1/1	0.97	0.12	34,34,34,34	0
25	MG	A	1625	1/1	0.97	0.18	34,34,34,34	0
25	MG	A	1563	1/1	0.97	0.17	34,34,34,34	0
25	MG	A	1578	1/1	0.97	0.16	34,34,34,34	0
25	MG	A	467	1/1	0.98	0.07	34,34,34,34	0
25	MG	A	1554	1/1	0.98	0.34	34,34,34,34	0
25	MG	A	470	1/1	0.99	0.28	34,34,34,34	0
26	ZN	D	306	1/1	0.99	0.10	34,34,34,34	0
26	ZN	N	307	1/1	0.99	0.04	34,34,34,34	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.