



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 20, 2024 – 06:31 pm GMT

PDB ID : 7NB1  
Title : Crystal structure of human choline alpha in complex with an inhibitor  
Authors : Casale, E.; Fasolini, M.  
Deposited on : 2021-01-25  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	384	 2% 80% 11% 9%
1	BBB	384	 2% 83% 9% 8%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Choline kinase alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	351	Total 2909	C 1884	N 487	O 522	S 16	0	0	0
1	BBB	353	Total 2879	C 1868	N 478	O 517	S 16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	74	GLY	-	expression tag	UNP P35790
BBB	74	GLY	-	expression tag	UNP P35790

- Molecule 2 is 4-(6-aminopurin-9-yl)- {N}-[4-(trifluoromethylsulfonyl)phenyl]cyclohexane-1-carboxamide (three-letter code: U6T) (formula: C<sub>19</sub>H<sub>19</sub>F<sub>3</sub>N<sub>6</sub>O<sub>3</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
2	AAA	1	32	19	3	6	3	1	0	0
2	BBB	1	32	19	3	6	3	1	0	0

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	AAA	1	15	8	2	4	1	0	0
3	AAA	1	15	8	2	4	1	0	0
3	BBB	1	15	8	2	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	78	Total	O	0	0
			78	78		
4	BBB	64	Total	O	0	0
			64	64		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Choline kinase alpha



- Molecule 1: Choline kinase alpha



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.85Å 122.00Å 132.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.86 – 2.30 44.82 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (44.86-2.30) 100.0 (44.82-2.30)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.03 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.195 , 0.249 0.195 , 0.249	Depositor DCC
$R_{free}$ test set	2094 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.4	Xtrriage
Anisotropy	0.487	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 31.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6039	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: U6T, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	AAA	0.40	0/2985	0.73	0/4020
1	BBB	0.41	0/2955	0.71	0/3989
All	All	0.41	0/5940	0.72	0/8009

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2909	0	2880	27	0
1	BBB	2879	0	2811	16	0
2	AAA	32	0	0	3	0
2	BBB	32	0	0	0	0
3	AAA	30	0	35	1	0
3	BBB	15	0	17	1	0
4	AAA	78	0	0	1	0
4	BBB	64	0	0	1	0
All	All	6039	0	5743	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:371:THR:H	1:AAA:374:GLN:HE21	1.31	0.76
1:AAA:192:LEU:CD1	1:AAA:238:MET:SD	2.84	0.66
1:AAA:267:GLU:CG	1:AAA:270:ARG:HG3	2.29	0.63
1:AAA:292:LEU:HD22	1:AAA:457:VAL:HG11	1.84	0.60
1:AAA:267:GLU:HG3	1:AAA:270:ARG:HG3	1.86	0.57
1:AAA:371:THR:H	1:AAA:374:GLN:NE2	2.01	0.56
1:BBB:192:LEU:CD1	1:BBB:238:MET:SD	2.96	0.53
1:AAA:267:GLU:HG2	1:AAA:270:ARG:HG3	1.90	0.52
1:BBB:217:GLU:H	1:BBB:217:GLU:CD	2.11	0.52
1:BBB:352:TYR:O	4:BBB:601:HOH:O	2.19	0.52
1:AAA:250:PHE:CE1	1:AAA:253:MET:HE1	2.45	0.51
1:AAA:86:ARG:O	1:AAA:89:ARG:HB3	2.12	0.50
1:AAA:88:ARG:NH2	1:AAA:111:PHE:O	2.44	0.50
1:BBB:212:ARG:NH2	1:BBB:314:LEU:HD22	2.26	0.50
1:AAA:457:VAL:O	1:AAA:457:VAL:HG12	2.13	0.49
1:AAA:423:TRP:O	1:AAA:427:GLN:HG2	2.13	0.49
2:AAA:501:U6T:O20	2:AAA:501:U6T:C23	2.61	0.49
1:BBB:390:PHE:C	1:BBB:392:ASN:H	2.16	0.49
1:AAA:252:THR:HG22	1:AAA:256:TYR:CE2	2.49	0.48
1:BBB:282:LEU:N	1:BBB:283:PRO:CD	2.77	0.47
1:BBB:423:TRP:CH2	3:BBB:502:EPE:H91	2.50	0.47
1:AAA:304:HIS:O	1:AAA:305:ASN:HB2	2.15	0.46
1:AAA:149:GLY:O	1:AAA:150:ALA:C	2.54	0.46
1:BBB:303:CYS:SG	1:BBB:337:ASN:HB3	2.56	0.45
1:BBB:452:LYS:HG2	1:BBB:457:VAL:HG23	1.98	0.45
1:AAA:116:ILE:HG23	2:AAA:501:U6T:C27	2.47	0.45
1:AAA:282:LEU:N	1:AAA:283:PRO:CD	2.80	0.44
1:BBB:308:GLN:HB3	1:BBB:349:GLU:HG3	2.00	0.44
1:BBB:125:PHE:HB2	1:BBB:145:LEU:HB3	2.00	0.43
1:AAA:192:LEU:HD12	1:AAA:238:MET:SD	2.59	0.43
1:BBB:192:LEU:HD12	1:BBB:238:MET:SD	2.59	0.42
1:AAA:373:LYS:HE2	4:AAA:674:HOH:O	2.19	0.42
1:AAA:192:LEU:HD11	1:AAA:238:MET:HE1	2.01	0.42
1:BBB:383:LEU:N	1:BBB:384:PRO:HD2	2.35	0.42
1:AAA:123:MET:HE1	1:AAA:149:GLY:HA3	2.02	0.42
1:AAA:87:THR:HA	1:AAA:90:ARG:HG3	2.01	0.42
1:AAA:265:PHE:O	1:AAA:271:ILE:HD11	2.20	0.41
1:AAA:244:LYS:HD3	1:AAA:336:TYR:CZ	2.54	0.41
1:AAA:431:SER:O	3:AAA:503:EPE:H51	2.20	0.41
1:AAA:116:ILE:HG23	2:AAA:501:U6T:C22	2.51	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:267:GLU:CD	1:AAA:267:GLU:H	2.22	0.41
1:BBB:256:TYR:O	1:BBB:260:VAL:HG23	2.21	0.41
1:BBB:211:SER:HA	1:BBB:316:GLU:HG3	2.02	0.40
1:BBB:368:LYS:HD3	1:BBB:368:LYS:HA	1.92	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	347/384 (90%)	336 (97%)	11 (3%)	0	100	100
1	BBB	349/384 (91%)	338 (97%)	11 (3%)	0	100	100
All	All	696/768 (91%)	674 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	314/342 (92%)	303 (96%)	11 (4%)	36	50
1	BBB	305/342 (89%)	295 (97%)	10 (3%)	38	53
All	All	619/684 (90%)	598 (97%)	21 (3%)	37	51

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	87	THR
1	AAA	110	GLU
1	AAA	114	SER
1	AAA	121	SER
1	AAA	128	SER
1	AAA	175	GLU
1	AAA	179	LEU
1	AAA	204	ARG
1	AAA	279	SER
1	AAA	388	ASN
1	AAA	399	SER
1	BBB	96	LYS
1	BBB	112	HIS
1	BBB	131	ASP
1	BBB	217	GLU
1	BBB	269	SER
1	BBB	291	SER
1	BBB	321	SER
1	BBB	342	ASP
1	BBB	389	ASP
1	BBB	407	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	U6T	BBB	501	-	32,35,35	2.69	4 (12%)	39,53,53	2.33	6 (15%)
3	EPE	AAA	502	-	15,15,15	0.68	1 (6%)	18,20,20	1.02	1 (5%)
3	EPE	BBB	502	-	15,15,15	1.07	1 (6%)	18,20,20	1.43	2 (11%)
2	U6T	AAA	501	-	32,35,35	2.62	4 (12%)	39,53,53	2.59	7 (17%)
3	EPE	AAA	503	-	15,15,15	1.11	1 (6%)	18,20,20	1.47	3 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	U6T	BBB	501	-	-	1/23/37/37	0/4/4/4
3	EPE	AAA	502	-	-	5/9/19/19	0/1/1/1
3	EPE	BBB	502	-	-	5/9/19/19	0/1/1/1
2	U6T	AAA	501	-	-	9/23/37/37	0/4/4/4
3	EPE	AAA	503	-	-	5/9/19/19	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	BBB	501	U6T	C25-S1	-11.95	1.62	1.76
2	AAA	501	U6T	C25-S1	-11.78	1.63	1.76
2	BBB	501	U6T	C29-S1	-7.61	1.70	1.85
2	AAA	501	U6T	C29-S1	-6.85	1.72	1.85
3	BBB	502	EPE	O2S-S	3.91	1.56	1.45
3	AAA	503	EPE	O2S-S	3.48	1.55	1.45
2	AAA	501	U6T	C22-N21	-2.88	1.35	1.41
2	BBB	501	U6T	C22-N21	-2.65	1.36	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	501	U6T	C4-N5	2.36	1.35	1.32
3	AAA	502	EPE	O3S-S	2.33	1.55	1.47
2	BBB	501	U6T	C4-N5	2.31	1.35	1.32

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	501	U6T	O28-S1-O2	-11.72	106.10	118.98
2	BBB	501	U6T	O28-S1-O2	-10.50	107.44	118.98
2	AAA	501	U6T	O2-S1-C29	5.98	112.87	104.69
2	BBB	501	U6T	O28-S1-C29	5.98	112.87	104.69
2	AAA	501	U6T	N5-C4-N3	-4.81	121.16	128.68
2	BBB	501	U6T	N5-C4-N3	-4.56	121.56	128.68
3	BBB	502	EPE	O2S-S-C10	-4.36	101.66	106.92
3	AAA	503	EPE	O3S-S-O1S	4.03	121.12	111.27
2	BBB	501	U6T	O2-S1-C29	3.71	109.77	104.69
2	AAA	501	U6T	C26-C25-S1	3.61	122.49	119.37
2	AAA	501	U6T	O28-S1-C29	3.44	109.39	104.69
3	BBB	502	EPE	O3S-S-O1S	3.07	118.78	111.27
2	AAA	501	U6T	C14-C15-C16	2.90	116.04	111.18
2	AAA	501	U6T	F32-C29-S1	2.74	114.64	110.30
3	AAA	502	EPE	O3S-S-C10	-2.49	101.74	105.77
2	BBB	501	U6T	O2-S1-C25	2.15	110.88	108.27
3	AAA	503	EPE	O3S-S-O2S	-2.09	106.17	111.27
2	BBB	501	U6T	N12-C6-N3	2.08	122.89	118.57
3	AAA	503	EPE	O2S-S-O1S	-2.07	106.77	113.95

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	501	U6T	C24-C25-S1-C29
3	AAA	502	EPE	C9-C10-S-O2S
3	AAA	503	EPE	C10-C9-N1-C2
3	BBB	502	EPE	S-C10-C9-N1
3	AAA	503	EPE	N4-C7-C8-O8
2	AAA	501	U6T	C24-C25-S1-O2
2	AAA	501	U6T	C26-C25-S1-O2
2	AAA	501	U6T	C26-C25-S1-C29
3	AAA	502	EPE	C10-C9-N1-C6
3	AAA	503	EPE	C10-C9-N1-C6
3	BBB	502	EPE	C10-C9-N1-C2

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	AAA	502	EPE	C8-C7-N4-C3
3	AAA	502	EPE	C8-C7-N4-C5
3	AAA	503	EPE	C8-C7-N4-C3
3	AAA	503	EPE	C8-C7-N4-C5
2	AAA	501	U6T	C17-C16-C19-O20
2	AAA	501	U6T	C24-C25-S1-O28
2	BBB	501	U6T	C17-C16-C19-O20
3	AAA	502	EPE	C10-C9-N1-C2
3	BBB	502	EPE	C10-C9-N1-C6
3	BBB	502	EPE	C8-C7-N4-C3
3	BBB	502	EPE	C8-C7-N4-C5
2	AAA	501	U6T	C17-C16-C19-N21
2	AAA	501	U6T	C26-C25-S1-O28
2	AAA	501	U6T	C15-C16-C19-O20

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	BBB	502	EPE	1	0
2	AAA	501	U6T	3	0
3	AAA	503	EPE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	351/384 (91%)	0.09	9 (2%) 56 63	31, 49, 89, 113	1 (0%)
1	BBB	353/384 (91%)	-0.07	8 (2%) 60 67	32, 53, 98, 158	0
All	All	704/768 (91%)	0.01	17 (2%) 59 66	31, 51, 91, 158	1 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	322	GLU	4.8
1	BBB	317	GLY	4.3
1	BBB	320	ASN	3.9
1	AAA	87	THR	3.4
1	BBB	323	LYS	3.4
1	BBB	321	SER	3.3
1	BBB	318	ARG	3.0
1	BBB	324	GLN	2.7
1	AAA	89	ARG	2.5
1	AAA	111	PHE	2.5
1	AAA	120	LEU	2.4
1	AAA	390	PHE	2.3
1	AAA	85	PRO	2.3
1	BBB	82	GLN	2.2
1	AAA	265	PHE	2.1
1	AAA	320	ASN	2.1
1	AAA	92	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EPE	BBB	502	15/15	0.81	0.28	96,111,127,131	0
3	EPE	AAA	503	15/15	0.88	0.27	63,103,129,132	0
3	EPE	AAA	502	15/15	0.88	0.22	86,97,108,122	0
2	U6T	BBB	501	32/32	0.90	0.19	36,45,110,119	0
2	U6T	AAA	501	32/32	0.94	0.13	39,49,121,129	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.