



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 12, 2022 – 02:13 pm GMT

PDB ID : 7NBZ
Title : Crystal structure of ligand free open conformation of sulfoquinovosyl binding protein (SQBP) from *Agrobacterium tumefaciens*
Authors : Snow, A.; Sharma, M.; Davies, G.J.
Deposited on : 2021-01-28
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.24
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.24


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	 5% 89% 6% 5%
1	B	396	 % 93% . .
1	C	396	 % 94% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	A	401	-	-	X	-

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 19395 atoms, of which 8916 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sulfoquinovosyl binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	377	5767	1864	2859	471	562	11	95	5	0
1	B	385	6033	1937	3003	496	585	12	87	10	0
1	C	390	6110	1962	3045	502	589	12	89	8	0

There are 27 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP A0A083ZKV5
A	389	LEU	-	expression tag	UNP A0A083ZKV5
A	390	GLU	-	expression tag	UNP A0A083ZKV5
A	391	HIS	-	expression tag	UNP A0A083ZKV5
A	392	HIS	-	expression tag	UNP A0A083ZKV5
A	393	HIS	-	expression tag	UNP A0A083ZKV5
A	394	HIS	-	expression tag	UNP A0A083ZKV5
A	395	HIS	-	expression tag	UNP A0A083ZKV5
A	396	HIS	-	expression tag	UNP A0A083ZKV5
B	1	MET	-	initiating methionine	UNP A0A083ZKV5
B	389	LEU	-	expression tag	UNP A0A083ZKV5
B	390	GLU	-	expression tag	UNP A0A083ZKV5
B	391	HIS	-	expression tag	UNP A0A083ZKV5
B	392	HIS	-	expression tag	UNP A0A083ZKV5
B	393	HIS	-	expression tag	UNP A0A083ZKV5
B	394	HIS	-	expression tag	UNP A0A083ZKV5
B	395	HIS	-	expression tag	UNP A0A083ZKV5
B	396	HIS	-	expression tag	UNP A0A083ZKV5
C	1	MET	-	initiating methionine	UNP A0A083ZKV5
C	389	LEU	-	expression tag	UNP A0A083ZKV5
C	390	GLU	-	expression tag	UNP A0A083ZKV5
C	391	HIS	-	expression tag	UNP A0A083ZKV5
C	392	HIS	-	expression tag	UNP A0A083ZKV5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	393	HIS	-	expression tag	UNP A0A083ZKV5
C	394	HIS	-	expression tag	UNP A0A083ZKV5
C	395	HIS	-	expression tag	UNP A0A083ZKV5
C	396	HIS	-	expression tag	UNP A0A083ZKV5

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	Total 7	C 2	H 3	O 2	0	0
2	C	1	Total 7	C 2	H 3	O 2	0	0
2	C	1	Total 7	C 2	H 3	O 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	389	Total 389	O 389	0	0
3	B	518	Total 518	O 518	0	0
3	C	557	Total 557	O 557	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.32Å 78.21Å 83.46Å 109.04° 106.86° 104.77°	Depositor
Resolution (Å)	72.01 – 1.35 72.01 – 1.35	Depositor EDS
% Data completeness (in resolution range)	93.3 (72.01-1.35) 90.6 (72.01-1.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.8.0266	Depositor
R, R_{free}	0.183 , 0.209 0.186 , 0.211	Depositor DCC
R_{free} test set	11842 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	12.3	Xtriage
Anisotropy	0.343	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	19395	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.83	1/2969 (0.0%)	0.90	0/4033
1	B	0.79	0/3096	0.89	0/4204
1	C	0.83	0/3128	0.88	0/4246
All	All	0.82	1/9193 (0.0%)	0.89	0/12483

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	130	LYS	C-N	7.34	1.48	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2908	2859	2802	16	0
1	B	3030	3003	2968	10	0
1	C	3065	3045	2998	12	0
2	A	4	3	3	2	0
2	C	8	6	6	0	0
3	A	389	0	0	4	0
3	B	518	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	557	0	0	5	0
All	All	10479	8916	8777	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:255:LYS:CE	3:C:805:HOH:O	2.30	0.78
1:C:99[B]:ASN:ND2	1:C:277[B]:GLU:OE1	2.17	0.77
1:C:255:LYS:HE2	3:C:805:HOH:O	1.85	0.76
1:A:370:MET:HE2	1:A:375:GLY:HA2	1.69	0.74
1:C:99[B]:ASN:OD1	1:C:277[B]:GLU:OE2	2.06	0.74
1:C:255:LYS:HE3	3:C:805:HOH:O	1.94	0.67
1:A:370:MET:CE	3:A:802:HOH:O	2.45	0.64
1:A:370:MET:HE1	3:A:802:HOH:O	1.98	0.61
1:B:94:THR:HG22	1:B:273:LEU:HD22	1.84	0.59
1:C:83:PHE:CE2	1:C:291[C]:LYS:HD3	2.37	0.59
1:C:173:LEU:HB2	1:C:174:PRO:HD3	1.90	0.54
1:B:94:THR:HG23	3:B:498:HOH:O	2.08	0.53
1:B:82:ASP:OD1	1:B:106:VAL:HG12	2.11	0.50
1:A:10:SER:OG	2:A:401:ACT:H2	2.11	0.50
1:B:173:LEU:HB2	1:B:174:PRO:HD3	1.94	0.49
1:C:83:PHE:CE2	1:C:291[C]:LYS:CD	2.95	0.49
1:B:313[B]:LEU:HD22	1:B:320:TYR:HE2	1.78	0.49
1:A:224:ARG:HD2	1:A:224:ARG:C	2.34	0.48
1:A:58[B]:ASP:C	1:A:58[B]:ASP:OD1	2.51	0.48
1:A:173:LEU:HB2	1:A:174:PRO:HD3	1.95	0.48
1:B:313[B]:LEU:HD22	1:B:320:TYR:CE2	2.49	0.47
1:A:139:LYS:HG2	1:A:210:ILE:HD13	1.97	0.47
1:A:99:ASN:ND2	1:A:107:ALA:HB3	2.30	0.46
1:A:221:ASP:OD2	1:A:225:LYS:HE3	2.15	0.46
1:C:114:SER:OG	1:C:271:THR:HG21	2.15	0.46
1:C:360:THR:HG22	3:C:512:HOH:O	2.14	0.46
1:A:228:GLN:OE1	1:A:244:HIS:HB3	2.17	0.45
1:C:140:GLU:HG3	3:C:774:HOH:O	2.15	0.45
1:B:228:GLN:OE1	1:B:244:HIS:HB3	2.18	0.44
1:B:313[B]:LEU:CD2	1:B:320:TYR:HE2	2.31	0.44
2:A:401:ACT:H1	3:A:756:HOH:O	2.18	0.43
1:B:48:GLN:HE22	1:B:51:ASN:HD22	1.65	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:ARG:N	1:A:71:PRO:CD	2.82	0.43
1:A:370:MET:HE3	3:A:802:HOH:O	2.16	0.42
1:C:228:GLN:OE1	1:C:244:HIS:HB3	2.20	0.41
1:A:176:TRP:CZ3	1:A:272[A]:CYS:SG	3.10	0.40
1:A:134:THR:HA	1:A:263:ALA:O	2.20	0.40
1:B:70:ARG:N	1:B:71:PRO:CD	2.85	0.40
1:A:178:GLU:HG2	1:A:264:VAL:HG21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	372/396 (94%)	371 (100%)	1 (0%)	0	100	100
1	B	391/396 (99%)	383 (98%)	8 (2%)	0	100	100
1	C	397/396 (100%)	393 (99%)	4 (1%)	0	100	100
All	All	1160/1188 (98%)	1147 (99%)	13 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/332 (90%)	297 (100%)	1 (0%)	92	83
1	B	318/332 (96%)	316 (99%)	2 (1%)	86	69
1	C	316/332 (95%)	313 (99%)	3 (1%)	78	53
All	All	932/996 (94%)	926 (99%)	6 (1%)	84	69

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	LEU
1	B	64	LEU
1	B	237	ASN
1	C	64	LEU
1	C	89	SER
1	C	237	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	286	GLN
1	B	14	GLN
1	B	48	GLN
1	B	253	ASN
1	C	253	ASN
1	C	286	GLN
1	C	363	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACT	A	401	-	1,3,3	2.41	1 (100%)	0,3,3	-	-
2	ACT	C	402	-	1,3,3	4.10	1 (100%)	0,3,3	-	-
2	ACT	C	401	-	1,3,3	5.00	1 (100%)	0,3,3	-	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	ACT	CH3-C	5.00	1.55	1.48
2	C	402	ACT	CH3-C	4.10	1.54	1.48
2	A	401	ACT	CH3-C	2.41	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	ACT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	377/396 (95%)	0.31	20 (5%) 26 29	8, 15, 32, 40	2 (0%)
1	B	385/396 (97%)	0.04	4 (1%) 82 85	7, 12, 22, 34	1 (0%)
1	C	390/396 (98%)	-0.04	2 (0%) 91 92	7, 12, 21, 31	0
All	All	1152/1188 (96%)	0.10	26 (2%) 60 66	7, 13, 27, 40	3 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	3	ALA	4.0
1	C	87	ASP	3.5
1	A	123	LEU	3.4
1	B	1	MET	3.4
1	B	183	VAL	3.2
1	A	387	LEU	3.1
1	A	229	ALA	2.8
1	A	127	TYR	2.7
1	A	210	ILE	2.6
1	A	131	PRO	2.5
1	A	154	LEU	2.5
1	A	119	TYR	2.5
1	A	252	VAL	2.5
1	B	2	ASP	2.4
1	A	126	LYS	2.4
1	A	233	LEU	2.4
1	A	230	GLY	2.3
1	A	245	PHE	2.2
1	A	386	VAL	2.1
1	A	124	LEU	2.1
1	C	88[A]	LEU	2.1
1	A	122	ASP	2.1
1	A	227	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	148	GLY	2.1
1	A	272[A]	CYS	2.0
1	A	242	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	C	402	4/4	0.77	0.26	24,26,28,28	0
2	ACT	C	401	4/4	0.79	0.19	30,30,39,40	0
2	ACT	A	401	4/4	0.85	0.19	24,25,29,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.