



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 19, 2022 – 04:10 pm BST

PDB ID : 5NC7
Title : ENAH EVH1 in complex with Ac-WPPPTEDEL-NH2
Authors : Barone, M.; Roske, Y.
Deposited on : 2017-03-03
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	113	
1	B	113	
1	C	113	
1	D	113	
2	I	12	
2	J	12	
2	K	12	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	L	12	 83% 8% 8%
2	W	12	 50% 17% 33%
2	X	12	 25% 50% 25%
2	Y	12	 25% 8% 67%
2	Z	12	 33% 33% 33%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein enabled homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	109	863	541	162	155	5	0	0	0
1	B	110	869	544	163	157	5	1	0	0
1	C	110	859	538	160	156	5	1	0	0
1	D	110	860	538	160	157	5	2	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q8N8S7
A	0	SER	-	expression tag	UNP Q8N8S7
B	-1	GLY	-	expression tag	UNP Q8N8S7
B	0	SER	-	expression tag	UNP Q8N8S7
C	-1	GLY	-	expression tag	UNP Q8N8S7
C	0	SER	-	expression tag	UNP Q8N8S7
D	-1	GLY	-	expression tag	UNP Q8N8S7
D	0	SER	-	expression tag	UNP Q8N8S7

- Molecule 2 is a protein called ActA-derived 10-mer Ac-FPPPPTEDEL-NH2 with acetylated (Ac) and amidated (NH2) termini. Phe is substituted by Trp to increase affinity for crystallization.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	K	12	87	57	12	18	0	0	1
2	I	12	87	57	12	18	0	0	1
2	J	12	87	57	12	18	0	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	L	12	Total	C	N	O	0	0	1
			87	57	12	18			
2	Z	8	Total	C	N	O	0	0	1
			44	29	7	8			
2	X	9	Total	C	N	O	0	0	1
			62	42	9	11			
2	Y	4	Total	C	N	O	0	0	0
			28	20	4	4			
2	W	8	Total	C	N	O	0	0	0
			52	34	7	11			


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	O	0	0
			3	3		
3	B	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein enabled homolog

Chain A:  87% 8% . .




- Molecule 1: Protein enabled homolog

Chain B:  88% 10% .




- Molecule 1: Protein enabled homolog

Chain C:  79% 17% . .




- Molecule 1: Protein enabled homolog

Chain D:  86% 11% . .



- Molecule 2: ActA-derived 10-mer Ac-FPPPTEDEL-NH2 with acetylated (Ac) and amidated (NH2) termini. Phe is substituted by Trp to increase affinity for crystallization

Chain K:  75% 17% 8%



- Molecule 2: ActA-derived 10-mer Ac-FPPPTEDEL-NH2 with acetylated (Ac) and amidated (NH2) termini. Phe is substituted by Trp to increase affinity for crystallization

Chain I:  100%


There are no outlier residues recorded for this chain.

- Molecule 2: ActA-derived 10-mer Ac-FPPPPTTEDEL-NH2 with acetylated (Ac) and amidated (NH2) termini. Phe is substituted by Trp to increase affinity for crystallization

Chain J:  100%


There are no outlier residues recorded for this chain.

- Molecule 2: ActA-derived 10-mer Ac-FPPPPTTEDEL-NH2 with acetylated (Ac) and amidated (NH2) termini. Phe is substituted by Trp to increase affinity for crystallization

Chain L:  83% 8% 8%



- Molecule 2: ActA-derived 10-mer Ac-FPPPPTTEDEL-NH2 with acetylated (Ac) and amidated (NH2) termini. Phe is substituted by Trp to increase affinity for crystallization

Chain Z:  33% 33% 33%




- Molecule 2: ActA-derived 10-mer Ac-FPPPPTTEDEL-NH2 with acetylated (Ac) and amidated (NH2) termini. Phe is substituted by Trp to increase affinity for crystallization

Chain X:  25% 50% 25%



- Molecule 2: ActA-derived 10-mer Ac-FPPPPTTEDEL-NH2 with acetylated (Ac) and amidated (NH2) termini. Phe is substituted by Trp to increase affinity for crystallization

Chain Y:  25% 8% 67%



- Molecule 2: ActA-derived 10-mer Ac-FPPPPTTEDEL-NH2 with acetylated (Ac) and amidated (NH2) termini. Phe is substituted by Trp to increase affinity for crystallization

Chain W:  50% 17% 33%



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	187.96Å 34.43Å 110.99Å 90.00° 91.53° 90.00°	Depositor
Resolution (Å)	48.34 – 2.70 48.34 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.34-2.70) 99.9 (48.34-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.209 , 0.272 0.210 , 0.272	Depositor DCC
R_{free} test set	1011 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	49.1	Xtrriage
Anisotropy	0.408	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.125 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3989	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NH2, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.55	1/883 (0.1%)	0.68	0/1192
1	B	0.52	0/889	0.69	0/1200
1	C	0.43	0/879	0.65	1/1188 (0.1%)
1	D	0.43	0/880	0.69	0/1189
2	I	0.50	0/89	0.79	0/126
2	J	0.75	0/89	0.64	0/126
2	K	0.34	0/89	0.43	0/126
2	L	0.34	0/89	0.53	0/126
2	W	0.46	0/53	0.54	0/76
2	X	0.49	0/65	0.56	0/94
2	Y	0.41	0/31	0.46	0/44
2	Z	0.38	0/45	0.50	0/66
All	All	0.49	1/4081 (0.0%)	0.66	1/5553 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	7	CYS	CB-SG	-5.24	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	66	LYS	CD-CE-NZ	-6.89	95.86	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	863	0	835	10	0
1	B	869	0	840	7	0
1	C	859	0	817	14	0
1	D	860	0	820	9	0
2	I	87	0	75	0	0
2	J	87	0	75	0	0
2	K	87	0	75	2	0
2	L	87	0	75	1	0
2	W	52	0	46	3	0
2	X	62	0	54	4	0
2	Y	28	0	28	1	0
2	Z	44	0	40	3	0
3	A	3	0	0	0	0
3	B	1	0	0	0	0
All	All	3989	0	3780	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 46 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:51:ARG:NH1	2:X:2:PRO:O	2.12	0.82
1:D:52:LYS:HB3	1:D:55:ASP:OD2	1.87	0.74
1:B:51:ARG:HB2	2:X:0:ACE:H1	1.73	0.71
1:C:46:PHE:CD1	1:C:111:LEU:HD11	2.27	0.69
2:X:6:THR:HG22	2:X:7:GLU:H	1.59	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	107/113 (95%)	104 (97%)	3 (3%)	0	100	100
1	B	108/113 (96%)	103 (95%)	5 (5%)	0	100	100
1	C	108/113 (96%)	105 (97%)	3 (3%)	0	100	100
1	D	108/113 (96%)	105 (97%)	3 (3%)	0	100	100
2	I	10/12 (83%)	10 (100%)	0	0	100	100
2	J	10/12 (83%)	10 (100%)	0	0	100	100
2	K	10/12 (83%)	10 (100%)	0	0	100	100
2	L	10/12 (83%)	10 (100%)	0	0	100	100
2	W	6/12 (50%)	5 (83%)	1 (17%)	0	100	100
2	X	7/12 (58%)	5 (71%)	1 (14%)	1 (14%)	0	0
2	Y	2/12 (17%)	2 (100%)	0	0	100	100
2	Z	6/12 (50%)	5 (83%)	1 (17%)	0	100	100
All	All	492/548 (90%)	474 (96%)	17 (4%)	1 (0%)	47	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	X	1	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/92 (97%)	85 (96%)	4 (4%)	27	55
1	B	90/92 (98%)	87 (97%)	3 (3%)	38	67
1	C	87/92 (95%)	84 (97%)	3 (3%)	37	66
1	D	88/92 (96%)	84 (96%)	4 (4%)	27	55
2	I	10/10 (100%)	10 (100%)	0	100	100
2	J	10/10 (100%)	10 (100%)	0	100	100
2	K	10/10 (100%)	9 (90%)	1 (10%)	7	18
2	L	10/10 (100%)	9 (90%)	1 (10%)	7	18
2	W	6/10 (60%)	6 (100%)	0	100	100
2	X	7/10 (70%)	7 (100%)	0	100	100
2	Y	4/10 (40%)	4 (100%)	0	100	100
2	Z	5/10 (50%)	5 (100%)	0	100	100
All	All	416/448 (93%)	400 (96%)	16 (4%)	33	62

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	K	8	ASP
1	D	81	ARG
1	C	93	SER
1	D	54	GLN
1	C	81	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.