



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 04:33 pm BST

PDB ID : 4NDL  
Title : Computational design and experimental verification of a symmetric homodimer  
Authors : Mou, Y.; Huang, P.S.; Hsu, F.C.; Huang, S.J.; Mayo, S.L.  
Deposited on : 2013-10-26  
Resolution : 3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	72	50% 10% • 39%
1	B	72	44% 17% • 38%
1	C	72	43% 14% • • 38%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2024 atoms, of which 953 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENH-c2b, computational designed homodimer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
1	B	45	676	236	318	62	60	0	0	0
1	A	44	650	230	303	57	60	0	0	0
1	C	45	697	238	332	65	62	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	O	0	0
			1	1		



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.55Å 167.77Å 29.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.60 – 3.50 25.15 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.2 (24.60-3.50) 98.3 (25.15-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.31 (at 2.80Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.312 , 0.358 0.311 , 0.365	Depositor DCC
$R_{free}$ test set	277 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.3	Xtrriage
Anisotropy	0.459	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 68.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.74	EDS
Total number of atoms	2024	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/357	0.87	0/485
1	B	0.42	0/368	0.83	0/499
1	C	0.56	0/374	1.04	3/505 (0.6%)
All	All	0.48	0/1099	0.92	3/1489 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	53	ARG	NE-CZ-NH1	7.49	124.04	120.30
1	C	53	ARG	CG-CD-NE	-7.42	96.22	111.80
1	C	53	ARG	CB-CA-C	-6.01	98.38	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	24	PHE	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	347	303	303	7	0
1	B	358	318	318	9	0
1	C	365	332	332	18	0
2	B	1	0	0	0	0
All	All	1071	953	953	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:50:LEU:HA	1:C:53:ARG:HH12	1.28	0.95
1:C:53:ARG:NH1	1:C:53:ARG:HB2	2.04	0.73
1:C:50:LEU:HA	1:C:53:ARG:NH1	2.04	0.72
1:B:57:ASN:OD1	1:B:60:GLN:NE2	2.29	0.62
1:C:47:ARG:NH1	1:C:58:GLU:OE1	2.37	0.58
1:B:51:SER:OG	1:B:56:LEU:O	2.14	0.58
1:C:49:TYR:CD1	1:C:53:ARG:NH2	2.72	0.57
1:C:28:GLN:O	1:C:32:LEU:N	2.38	0.56
1:C:33:ASP:OD1	1:C:68:LYS:NZ	2.40	0.54
1:C:54:LEU:HD12	1:C:56:LEU:HD12	1.90	0.53
1:C:49:TYR:CG	1:C:53:ARG:NH2	2.78	0.51
1:C:53:ARG:HH11	1:C:53:ARG:HB2	1.73	0.50
1:C:39:ASP:O	1:C:46:TRP:CH2	2.65	0.50
1:B:36:PHE:HB2	1:B:65:PHE:HE2	1.76	0.49
1:C:53:ARG:CZ	1:C:53:ARG:HB2	2.42	0.49
1:A:39:ASP:OD1	1:A:46:TRP:CH2	2.65	0.48
1:B:49:TYR:OH	1:B:53:ARG:HD3	2.14	0.47
1:A:38:PHE:CD2	1:A:38:PHE:C	2.88	0.47
1:B:49:TYR:CZ	1:A:46:TRP:CD1	3.03	0.47
1:C:49:TYR:CE1	1:C:53:ARG:NH2	2.82	0.47
1:B:46:TRP:CD1	1:A:49:TYR:CZ	3.04	0.46
1:C:50:LEU:CA	1:C:53:ARG:HH12	2.13	0.46
1:C:42:LEU:HD23	1:C:46:TRP:CE3	2.52	0.45
1:B:42:LEU:CD1	1:B:65:PHE:HD1	2.31	0.44
1:C:42:LEU:HD22	1:C:47:ARG:HG2	2.00	0.44
1:C:53:ARG:HB3	1:C:54:LEU:HD23	2.00	0.43
1:B:36:PHE:HB2	1:B:65:PHE:CE2	2.54	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:49:TYR:C	1:C:53:ARG:HH22	2.23	0.42
1:A:39:ASP:OD1	1:A:39:ASP:N	2.52	0.42
1:B:42:LEU:HD11	1:B:65:PHE:CD1	2.54	0.42
1:A:51:SER:OG	1:A:56:LEU:O	2.26	0.41
1:A:36:PHE:CE1	1:A:40:ARG:CB	3.05	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	42/72 (58%)	40 (95%)	2 (5%)	0	100	100
1	B	43/72 (60%)	42 (98%)	1 (2%)	0	100	100
1	C	43/72 (60%)	42 (98%)	1 (2%)	0	100	100
All	All	128/216 (59%)	124 (97%)	4 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	30/66 (46%)	29 (97%)	1 (3%)	38	68
1	B	31/66 (47%)	28 (90%)	3 (10%)	8	33

Continued on next page...



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	C	32/66 (48%)	28 (88%)	4 (12%)	4 23
All	All	93/198 (47%)	85 (91%)	8 (9%)	10 38

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	43	THR
1	B	52	GLN
1	B	56	LEU
1	A	38	PHE
1	C	32	LEU
1	C	47	ARG
1	C	53	ARG
1	C	54	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.