



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 10, 2021 – 02:14 PM GMT

PDB ID : 7NFU
Title : Crystal structure of C-terminally truncated *Geobacillus thermoleovorans* nucleoid occlusion protein Noc
Authors : Jalal, A.S.B.; Tran, N.T.; Wu, L.J.; Ramakrishnan, K.; Rejzek, M.; Stevenson, C.E.M.; Lawson, D.M.; Errington, J.; Le, T.B.K.
Deposited on : 2021-02-07
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.16
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	
1	B	252	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	307	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3588 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoid occlusion protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	221	1756	1105	315	332	4	0	0	0
1	B	213	1660	1048	301	308	3	0	0	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	240	LYS	-	expression tag	UNP G8N1K9
A	241	LEU	-	expression tag	UNP G8N1K9
A	242	ALA	-	expression tag	UNP G8N1K9
A	243	ALA	-	expression tag	UNP G8N1K9
A	244	ALA	-	expression tag	UNP G8N1K9
A	245	LEU	-	expression tag	UNP G8N1K9
A	246	GLU	-	expression tag	UNP G8N1K9
A	247	HIS	-	expression tag	UNP G8N1K9
A	248	HIS	-	expression tag	UNP G8N1K9
A	249	HIS	-	expression tag	UNP G8N1K9
A	250	HIS	-	expression tag	UNP G8N1K9
A	251	HIS	-	expression tag	UNP G8N1K9
A	252	HIS	-	expression tag	UNP G8N1K9
B	240	LYS	-	expression tag	UNP G8N1K9
B	241	LEU	-	expression tag	UNP G8N1K9
B	242	ALA	-	expression tag	UNP G8N1K9
B	243	ALA	-	expression tag	UNP G8N1K9
B	244	ALA	-	expression tag	UNP G8N1K9
B	245	LEU	-	expression tag	UNP G8N1K9
B	246	GLU	-	expression tag	UNP G8N1K9
B	247	HIS	-	expression tag	UNP G8N1K9
B	248	HIS	-	expression tag	UNP G8N1K9
B	249	HIS	-	expression tag	UNP G8N1K9
B	250	HIS	-	expression tag	UNP G8N1K9
B	251	HIS	-	expression tag	UNP G8N1K9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	252	HIS	-	expression tag	UNP G8N1K9

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	23	Total O 23 23	0	0
4	B	8	Total O 8 8	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	146.79Å 146.79Å 146.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	84.89 – 2.50 84.75 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (84.89-2.50) 99.9 (84.75-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.209 , 0.240 0.212 , 0.238	Depositor DCC
R_{free} test set	1793 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	75.1	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 56.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.032 for l,-k,h	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3588	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.76	1/1776 (0.1%)	0.89	2/2394 (0.1%)
1	B	0.80	2/1679 (0.1%)	0.88	0/2271
All	All	0.78	3/3455 (0.1%)	0.89	2/4665 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	27	GLU	CD-OE2	11.78	1.38	1.25
1	B	27	GLU	CD-OE1	6.67	1.32	1.25
1	A	24	GLU	CG-CD	5.96	1.60	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	85	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	A	24	GLU	OE1-CD-OE2	5.43	129.81	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1756	0	1782	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1660	0	1675	8	0
2	A	70	0	0	0	0
2	B	65	0	0	1	0
3	A	6	0	8	2	0
4	A	23	0	0	0	0
4	B	8	0	0	0	0
All	All	3588	0	3465	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 14 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:LEU:HD13	1:B:222:LEU:HD13	1.78	0.65
1:B:143:THR:HG22	1:B:145:GLU:H	1.61	0.64
1:A:143:THR:HG22	1:A:145:GLU:H	1.62	0.64
1:A:75:ASN:HB2	3:A:315:GOL:H31	1.81	0.61
1:A:64:ILE:O	1:A:64:ILE:HG23	2.01	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/252 (87%)	213 (97%)	6 (3%)	0	100	100
1	B	209/252 (83%)	204 (98%)	5 (2%)	0	100	100
All	All	428/504 (85%)	417 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/221 (83%)	177 (97%)	6 (3%)	38	64
1	B	170/221 (77%)	168 (99%)	2 (1%)	71	88
All	All	353/442 (80%)	345 (98%)	8 (2%)	50	76

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	GLU
1	B	183	ARG
1	A	186	ARG
1	A	122	GLU
1	A	149	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	140	HIS
1	B	140	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

28 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	303	-	4,4,4	0.36	0	6,6,6	0.10	0
2	SO4	A	314	-	4,4,4	0.39	0	6,6,6	0.04	0
2	SO4	B	307	-	4,4,4	0.36	0	6,6,6	0.05	0
2	SO4	B	304	-	4,4,4	0.35	0	6,6,6	0.08	0
2	SO4	A	305	-	4,4,4	0.39	0	6,6,6	0.06	0
2	SO4	A	311	-	4,4,4	0.34	0	6,6,6	0.07	0
3	GOL	A	315	-	5,5,5	0.11	0	5,5,5	0.40	0
2	SO4	B	308	-	4,4,4	0.37	0	6,6,6	0.05	0
2	SO4	B	310	-	4,4,4	0.37	0	6,6,6	0.11	0
2	SO4	A	312	-	4,4,4	0.34	0	6,6,6	0.07	0
2	SO4	B	305	-	4,4,4	0.34	0	6,6,6	0.08	0
2	SO4	B	311	-	4,4,4	0.33	0	6,6,6	0.12	0
2	SO4	A	313	-	4,4,4	0.37	0	6,6,6	0.06	0
2	SO4	B	312	-	4,4,4	0.35	0	6,6,6	0.08	0
2	SO4	A	310	-	4,4,4	0.34	0	6,6,6	0.08	0
2	SO4	A	308	-	4,4,4	0.34	0	6,6,6	0.08	0
2	SO4	A	307	-	4,4,4	0.36	0	6,6,6	0.07	0
2	SO4	B	313	-	4,4,4	0.36	0	6,6,6	0.05	0
2	SO4	A	302	-	4,4,4	0.43	0	6,6,6	0.26	0
2	SO4	A	309	-	4,4,4	0.26	0	6,6,6	0.18	0
2	SO4	B	303	-	4,4,4	0.37	0	6,6,6	0.04	0
2	SO4	A	306	-	4,4,4	0.35	0	6,6,6	0.07	0
2	SO4	A	301	-	4,4,4	0.50	0	6,6,6	0.12	0
2	SO4	B	309	-	4,4,4	0.31	0	6,6,6	0.16	0
2	SO4	B	302	-	4,4,4	0.47	0	6,6,6	0.14	0
2	SO4	B	306	-	4,4,4	0.36	0	6,6,6	0.08	0
2	SO4	B	301	-	4,4,4	0.41	0	6,6,6	0.14	0
2	SO4	A	304	-	4,4,4	0.36	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	315	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	315	GOL	O2-C2-C3-O3
3	A	315	GOL	O1-C1-C2-C3
3	A	315	GOL	C1-C2-C3-O3
3	A	315	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	315	GOL	2	0
2	B	309	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	221/252 (87%)	0.75	7 (3%) 47 51	53, 83, 112, 124	0
1	B	213/252 (84%)	0.80	11 (5%) 27 29	56, 85, 123, 167	0
All	All	434/504 (86%)	0.77	18 (4%) 37 40	53, 84, 118, 167	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	11	PHE	3.3
1	B	12	GLY	3.2
1	B	9	PHE	2.7
1	B	13	GLU	2.5
1	B	65	ILE	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	B	307	5/5	0.49	0.44	140,155,168,169	0
2	SO4	B	308	5/5	0.74	0.23	158,159,165,169	0
2	SO4	A	312	5/5	0.77	0.23	116,136,150,151	0
2	SO4	B	312	5/5	0.81	0.17	111,119,134,136	0
2	SO4	A	310	5/5	0.85	0.27	115,131,149,152	0
2	SO4	B	304	5/5	0.86	0.17	104,109,132,138	0
2	SO4	B	306	5/5	0.86	0.15	107,109,128,136	0
2	SO4	A	309	5/5	0.88	0.39	89,94,120,123	0
2	SO4	A	313	5/5	0.88	0.34	155,159,165,166	0
2	SO4	A	304	5/5	0.88	0.20	114,118,137,138	0
2	SO4	B	311	5/5	0.89	0.13	108,110,130,136	0
2	SO4	B	309	5/5	0.89	0.14	91,109,116,120	0
2	SO4	A	306	5/5	0.90	0.17	109,117,128,130	0
2	SO4	A	311	5/5	0.90	0.17	75,102,105,111	0
2	SO4	A	307	5/5	0.91	0.12	123,126,132,135	0
2	SO4	B	313	5/5	0.91	0.29	140,140,148,151	0
2	SO4	A	308	5/5	0.91	0.19	115,121,131,132	0
2	SO4	B	303	5/5	0.91	0.17	123,124,137,137	0
3	GOL	A	315	6/6	0.92	0.17	101,108,114,114	0
2	SO4	B	305	5/5	0.93	0.14	85,85,106,108	0
2	SO4	A	314	5/5	0.93	0.10	144,149,155,167	0
2	SO4	B	310	5/5	0.94	0.17	95,97,101,103	0
2	SO4	A	305	5/5	0.94	0.16	93,107,116,122	0
2	SO4	A	303	5/5	0.95	0.10	114,121,127,128	0
2	SO4	B	301	5/5	0.98	0.20	67,68,71,72	0
2	SO4	A	301	5/5	0.99	0.25	62,63,68,70	0
2	SO4	A	302	5/5	0.99	0.23	52,53,59,61	0
2	SO4	B	302	5/5	0.99	0.18	62,66,67,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.