



# Full wwPDB NMR Structure Validation Report ⓘ

May 29, 2020 – 09:32 am BST

PDB ID : 5NIP  
Title : An i-motif containing the neutral cytidine protonated analogue pseudoisocytidine  
Authors : Mir, B.; Soles, X.; Gonzalez, C.; Escaja, N.  
Deposited on : 2017-03-24

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11



## 2 Ensemble composition and analysis

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 692 atoms, of which 258 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D>(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3').


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	11	Total	C	H	N	O	P	0
			346	106	129	32	69	10	
1	B	11	Total	C	H	N	O	P	0
			346	106	129	32	69	10	

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble


These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A: 



- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B: 




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A: 



- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')


Chain B: 



### 4.2.2 Score per residue for model 2


- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A: 9% 73% 18%



- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B: 18% 55% 27%



### 4.2.3 Score per residue for model 3

- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A: 9% 64% 27%



- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B: 9% 64% 27%



### 4.2.4 Score per residue for model 4


- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A: 9% 45% 45%



- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B: 9% 55% 36%



#### 4.2.5 Score per residue for model 5

- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A: 



- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B: 




#### 4.2.6 Score per residue for model 6

- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A: 



- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B: 



#### 4.2.7 Score per residue for model 7

- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A: 




- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B: 




#### 4.2.8 Score per residue for model 8


- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A:  9% 55% 36%




- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B:  9% 64% 27%



#### 4.2.9 Score per residue for model 9

- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A:  9% 64% 27%




- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B:  9% 73% 18%



#### 4.2.10 Score per residue for model 10


- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain A:  9% 55% 36%



- Molecule 1: DNA (5'-D(\*TP\*(DCP)P\*CP\*GP\*TP\*TP\*TP\*CP\*(PSC)P\*GP\*T)-3')

Chain B:  9% 73% 18%





## 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *molecular dynamics*.

Of the 10 calculated structures, 10 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	
DYANA	structure solution	
Sparky	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	75
Number of shifts mapped to atoms	71
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	4
Assignment completeness (well-defined parts)	14%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

### 5.1 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	4340	2580	2338	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

## 5.2 Torsion angles [i](#)

### 5.2.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.2.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	8YN	A	9	1	17,20,21	1.43±0.03	0±0 (0±0%)
1	8YN	B	9	1	17,20,21	1.45±0.05	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	8YN	A	9	1	19,28,31	1.84±0.08	0±0 (0±0%)
1	8YN	B	9	1	19,28,31	1.97±0.11	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	8YN	B	9	1	-	0±0,7,21,22	0±0,2,2,2
1	8YN	A	9	1	-	0±0,7,21,22	0±0,2,2,2

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

#### 5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

#### 5.5 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.6 Other polymers [i](#)

There are no such molecules in this entry.

#### 5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 14% for the well-defined parts and 14% for the entire structure.

### 6.1 Chemical shift list 1

File name: input\_cs.cif

Chemical shift list name: *CCC.str*

#### 6.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	75
Number of shifts mapped to atoms	71
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	4
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atoms found in structure. All 4 occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	9	8YN	2H2'	1.41	0.01	1
A	9	8YN	2H4	8.67	0.06	1
A	9	8YN	1H4	6.1	0.0	1
A	9	8YN	1H2'	1.95	0.0	1

#### 6.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

#### 6.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 14%, i.e. 54 atoms were assigned a chemical

shift out of a possible 386. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	54/386 (14%)	54/224 (24%)	0/136 (0%)	0/26 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 14%, i.e. 54 atoms were assigned a chemical shift out of a possible 386. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	54/386 (14%)	54/224 (24%)	0/136 (0%)	0/26 (0%)

#### 6.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 6.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (CCC.str). RCI is only applicable to proteins.