

wwPDB X-ray Structure Validation Summary Report (i)

Aug 8, 2023 – 04:37 AM EDT

PDB ID : 1NJM

Title : The crystal structure of the 50S Large ribosomal subunit from Deinococcus

radiodurans complexed with a tRNA acceptor stem mimic (ASM) and the

antibiotic sparsomycin

Authors: Bashan, A.; Agmon, I.; Zarivatch, R.; Schluenzen, F.; Harms, J.M.; Berisio,

R.; Bartels, H.; Hansen, H.A.; Yonath, A.

Deposited on : 2003-01-02

Resolution : 3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$			
Clashscore	141614	1353 (3.70-3.50)			
RNA backbone	3102	1017 (4.20-3.00)			

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain					
1	0	2880	33%		51%	11% •		
2	5	35	26%	37%	9%	29%		
3	K	141		88%		12%		
4	Т	237		94%		6%		

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

\mathbf{Mol}	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SPS	0	2881	X	-	-	-



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 60271 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace	
1	0	2766	Total 59359	C 26479	N 10949	O 19166	P 2765	0	0	0

• Molecule 2 is a RNA chain called tRNA acceptor stem mimic.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace			
2	5	25	Total 543	C 249	N 97	O 173	P 24	0	0	0

• Molecule 3 is a protein called 50S ribosomal protein L16.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	K	124	Total C 124 124	0	0	124

• Molecule 4 is a protein called GENERAL STRESS PROTEIN CTC.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	Т	223	Total C 223 223	0	0	223

• Molecule 5 is SPARSOMYCIN (three-letter code: SPS) (formula: $C_{13}H_{19}N_3O_5S_2$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
5	0	1	Total	C 13	N 2	O 1	S	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: 23S ribosomal RNA












• Molecule 2: tRNA acceptor stem mimic

Chain 5: 26% 37% 9% 29%







4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	I 2 2 2	Depositor	
Cell constants	169.60Å 409.40Å 695.10Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	15.00 - 3.60	Depositor	
% Data completeness	(Not available) (15.00-3.60)	Depositor	
(in resolution range)	(100 available) (15.00-5.00)	Depositor	
R_{merge}	(Not available)	Depositor	
R_{sym}	0.17	Depositor	
Refinement program	CNS	Depositor	
R, R_{free}	0.284 , 0.308	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	60271	wwPDB-VP	
Average B, all atoms (Å ²)	74.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SPS, PPU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Chain	Bond	lengths	Bond angles		
			RMSZ	# Z > 5	RMSZ	# Z > 5	
	1	0	0.14	0/66467	0.63	0/103673	
	2	5	0.16	0/563	0.63	0/873	
	All	All	0.14	0/67030	0.63	0/104546	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers		
1	0	0	1		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	0	873	U	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	59359	0	29917	1742	0
2	5	543	0	290	16	0
3	K	124	0	0	0	0
4	Т	223	0	0	0	0
5	0	22	0	19	0	0
All	All	60271	0	30226	1754	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 1754 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)		
1:0:940:G:H3'	1:0:941:U:H5"	1.22	1.14		
1:0:1141:U:H3	1:0:2008:C:H5"	1.20	1.05		
1:0:1073:G:H2'	1:0:1074:G:H4'	1.40	1.00		
1:0:2548:G:H2'	1:0:2549:G:H5"	1.44	1.00		
1:0:2769:C:H2'	1:0:2867:G:H22	1.22	0.98		

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains (i)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	0	2757/2880 (95%)	416 (15%)	44 (1%)
2	5	22/35~(62%)	2 (9%)	0
All	All	2779/2915 (95%)	418 (15%)	44 (1%)

5 of 418 RNA backbone outliers are listed below:



Mol	Chain	Res	Type
1	0	15	G
1	0	35	G
1	0	45	С
1	0	48	A
1	0	49	U

5 of 44 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	0	1807	A
1	0	2245	A
1	0	1820	G
1	0	2093	G
1	0	2377	U

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
				nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	2	PPU	5	35	2,1	32,40,41	2.81	6 (18%)	33,57,60	0.98	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PPU	5	35	2,1	-	0/21/43/44	0/4/4/4

The worst 5 of 6 bond length outliers are listed below:

2 5 35 PPU C-N3' 12.63 1.61	Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(A)$	Ideal(A)
	2	5	35	PPU	C-N3'	12.63	1.61	1.34

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\text{\AA})$
2	5	35	PPU	OC-CM	-4.96	1.27	1.42
2	5	35	PPU	C6-N1	3.86	1.38	1.33
2	5	35	PPU	CE1-CZ	3.70	1.46	1.38
2	5	35	PPU	CE2-CZ	2.70	1.44	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
2	5	35	PPU	CM-OC-CZ	2.93	123.86	117.51
2	5	35	PPU	C9-N6-C6	2.23	126.26	119.51
2	5	35	PPU	C-CA-N	2.22	117.97	109.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	5	35	PPU	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
						Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	5	SPS	0	2881	-	19,22,23	4.43	9 (47%)	17,28,30	4.15	8 (47%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	\mathbf{Type}	Chain	Res	Link	Chirals	Torsions	Rings
5	SPS	0	2881	-	1/1/2/6	9/15/16/18	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
5	0	2881	SPS	O10-C10	9.82	1.43	1.24
5	0	2881	SPS	C9-C8	8.40	1.54	1.33
5	0	2881	SPS	O1-C1	7.67	1.43	1.24
5	0	2881	SPS	C10-N11	6.56	1.50	1.34
5	0	2881	SPS	C5-N4	5.43	1.42	1.34

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	0	2881	SPS	C12-N11-C10	-7.47	112.03	122.57
5	0	2881	SPS	C3-N2-C1	6.40	120.55	115.14
5	0	2881	SPS	C8-C9-C10	-6.14	109.55	121.56
5	0	2881	SPS	C14-S15-C16	6.14	109.42	101.04
5	0	2881	SPS	O10-C10-C9	-5.96	109.44	123.03

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	0	2881	SPS	C12

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	0	2881	SPS	C1-C6-C8-C9
5	0	2881	SPS	C9-C10-N11-C12
5	0	2881	SPS	O10-C10-N11-C12
5	0	2881	SPS	C14-C12-C13-O13
5	0	2881	SPS	N11-C12-C13-O13


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,



bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

