

wwPDB X-ray Structure Validation Summary Report (i)

Aug 8, 2023 – 08:18 AM EDT

PDB ID : 1NJN

Title: The crystal structure of the 50S Large ribosomal subunit from Deinococcus

radiodurans complexed with the antibiotic sparsomycin

Authors: Bashan, A.; Agmon, I.; Zarivatch, R.; Schluenzen, F.; Harms, J.M.; Berisio,

R.; Bartels, H.; Hansen, H.A.; Yonath, A.

Deposited on : 2003-01-02

Resolution : 3.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
Clashscore	141614	1027 (3.86-3.54)
RNA backbone	3102	1027 (4.40-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length		Quality of chain	
1	0	2880	34%	51%	11% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

\mathbf{Mol}	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SPS	0	2881	X	-	-	-



2 Entry composition (i)


There are 2 unique types of molecules in this entry. The entry contains 59381 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues		Atoms		ZeroOcc	AltConf	Trace		
1	0	2766	Total 59359	C 26479	N 10949	O 19166	P 2765	0	0	0

• Molecule 2 is SPARSOMYCIN (three-letter code: SPS) (formula: $C_{13}H_{19}N_3O_5S_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	0	1	Total 22	C 13	N 3	O 4	S 2	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: 23S ribosomal RNA













4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants	169.10Å 409.90Å 696.30Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 - 3.70	Depositor
% Data completeness	(Not available) (15.00-3.70)	Depositor
(in resolution range)	(1100 available) (19.00 9.10)	Берозпог
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.284 , 0.308	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	59381	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SPS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	Bond lengths		Bond angles	
Mol Chain	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	0	0.14	0/66467	0.63	0/103673	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	0	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	0	873	U	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	59359	0	29917	1711	0
2	0	22	0	19	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	59381	0	29936	1711	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 1711 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:0:940:G:H3'	1:0:941:U:H5"	1.24	1.14
1:0:1073:G:H2'	1:0:1074:G:H4'	1.39	1.02
1:0:2769:C:H2'	1:0:2867:G:H22	1.23	1.01
1:0:1141:U:H3	1:0:2008:C:H5"	1.22	1.01
1:0:2548:G:H2'	1:0:2549:G:H5"	1.44	0.99

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

There are no protein molecules in this entry.

5.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	0	2757/2880 (95%)	413 (14%)	39 (1%)

5 of 413 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	0	15	G
1	0	35	G
1	0	45	С
1	0	48	A

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	0	49	Ü

5 of 39 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	0	2093	G
1	0	2426	G
1	0	2161	С
1	0	2261	G
1	0	2759	U

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

1	/Iol	Type	Chain	Ros	Link	Bo	nd leng	$ ag{ths}$	В	ond ang	les
1	/101	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	2	SPS	0	2881	-	19,22,23	4.39	9 (47%)	17,28,30	4.15	8 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SPS	0	2881	-	1/1/2/6	8/15/16/18	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(A)
2	0	2881	SPS	O10-C10	9.77	1.43	1.24
2	0	2881	SPS	C9-C8	8.37	1.54	1.33
2	0	2881	SPS	O1-C1	7.62	1.43	1.24
2	0	2881	SPS	C10-N11	6.53	1.50	1.34
2	0	2881	SPS	C1-N2	5.41	1.42	1.33

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	0	2881	SPS	C12-N11-C10	-7.51	111.98	122.57
2	0	2881	SPS	C14-S15-C16	6.41	109.79	101.04
2	0	2881	SPS	C3-N2-C1	6.34	120.49	115.14
2	0	2881	SPS	C8-C9-C10	-6.10	109.64	121.56
2	0	2881	SPS	O10-C10-C9	-5.85	109.69	123.03

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	0	2881	SPS	C12

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	0	2881	SPS	C9-C10-N11-C12
2	0	2881	SPS	O10-C10-N11-C12
2	0	2881	SPS	C14-C12-C13-O13
2	0	2881	SPS	N11-C12-C13-O13
2	0	2881	SPS	C13-C12-N11-C10


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

