



Full wwPDB EM Validation Report ⓘ

Dec 19, 2022 – 11:14 am GMT

PDB ID : 7NKY
EMDB ID : EMD-12450
Title : RNA Polymerase II-Spt4/5-nucleosome-FACT structure
Authors : Farnung, L.; Ochmann, M.; Engholm, M.; Cramer, P.
Deposited on : 2021-02-19
Resolution : 3.20 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|-----------------------------|-----------------------------|
| Ramachandran outliers | 154571 | 4023 |
| Sidechain outliers | 154315 | 3826 |
| RNA backbone | 4643 | 859 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | P | 16 | |
| 2 | N | 138 | |
| 3 | T | 148 | |
| 4 | A | 1733 | |
| 5 | B | 1224 | |
| 6 | C | 318 | |
| 7 | E | 215 | |
| 8 | F | 155 | |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 9 | H | 146 | 90% 9% |
| 10 | I | 122 | 97% |
| 11 | J | 70 | 93% 7% |
| 12 | K | 120 | 96% |
| 13 | L | 70 | 63% 37% |
| 14 | Y | 102 | 96% |
| 15 | Z | 1064 | 15% 85% |
| 16 | D | 221 | 81% 19% |
| 17 | G | 171 | 99% |
| 18 | O | 552 | 76% 23% |
| 19 | Q | 1035 | 39% 61% |
| 20 | a | 136 | 63% 35% |
| 20 | e | 136 | 68% 31% |
| 21 | b | 103 | 81% 19% |
| 21 | f | 103 | 76% 24% |
| 22 | c | 130 | 78% 22% |
| 22 | g | 130 | 75% 25% |
| 23 | d | 123 | 77% 23% |
| 23 | h | 123 | 75% 24% |

2 Entry composition [i](#)

There are 25 unique types of molecules in this entry. The entry contains 51842 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(P*UP*CP*UP*UP*UP*UP*AP*UP*UP*UP*UP*UP*UP*CP*UP*G)-3').

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|-----|----|---------|-------|
| | | | Total | C | N | O | P | | |
| 1 | P | 16 | 325 | 146 | 40 | 123 | 16 | 0 | 0 |

- Molecule 2 is a DNA chain called DNA (138-MER).

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|-----|---------|-------|
| | | | Total | C | N | O | P | | |
| 2 | N | 138 | 2843 | 1346 | 526 | 833 | 138 | 0 | 0 |

- Molecule 3 is a DNA chain called DNA (148-MER).

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|-----|---------|-------|
| | | | Total | C | N | O | P | | |
| 3 | T | 148 | 3022 | 1434 | 567 | 874 | 147 | 0 | 0 |

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|------|------|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 4 | A | 1410 | 11092 | 6987 | 1941 | 2102 | 62 | 0 | 0 |

- Molecule 5 is a protein called DNA-directed RNA polymerase II subunit RPB2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|------|------|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 5 | B | 1116 | 8865 | 5611 | 1559 | 1640 | 55 | 0 | 0 |

- Molecule 6 is a protein called DNA-directed RNA polymerase II subunit RPB3.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 6 | C | 265 | 2086 | 1312 | 347 | 414 | 13 | 0 | 0 |

- Molecule 7 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 7 | E | 214 | 1752 | 1111 | 309 | 321 | 11 | 0 | 0 |

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 8 | F | 87 | 705 | 451 | 119 | 132 | 3 | 0 | 0 |

- Molecule 9 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 9 | H | 133 | 1068 | 673 | 180 | 211 | 4 | 0 | 0 |

- Molecule 10 is a protein called DNA-directed RNA polymerase II subunit RPB9.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 10 | I | 119 | 971 | 596 | 179 | 186 | 10 | 0 | 0 |

- Molecule 11 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 11 | J | 65 | 532 | 339 | 93 | 94 | 6 | 0 | 0 |

- Molecule 12 is a protein called DNA-directed RNA polymerase II subunit RPB11.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 12 | K | 115 | 920 | 590 | 157 | 171 | 2 | 0 | 1 |

- Molecule 13 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 13 | L | 44 | 351 | 217 | 70 | 60 | 4 | 0 | 0 |

- Molecule 14 is a protein called Chromatin elongation factor SPT4.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 14 | Y | 98 | 739 | 461 | 126 | 142 | 10 | 0 | 0 |

- Molecule 15 is a protein called Transcription elongation factor SPT5.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 15 | Z | 156 | 1252 | 804 | 224 | 221 | 3 | 0 | 0 |

There are 5 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|----------------|
| Z | 18 | PHE | VAL | conflict | UNP A0A6L0ZFJ2 |
| Z | 98 | THR | SER | conflict | UNP A0A6L0ZFJ2 |
| Z | 176 | GLU | ASP | conflict | UNP A0A6L0ZFJ2 |
| Z | 376 | LEU | LYS | conflict | UNP A0A6L0ZFJ2 |
| Z | 1064 | PHE | - | expression tag | UNP A0A6L0ZFJ2 |

- Molecule 16 is a protein called DNA-directed RNA polymerase II subunit RPB4.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 16 | D | 180 | 1444 | 893 | 257 | 291 | 3 | 0 | 0 |

- Molecule 17 is a protein called DNA-directed RNA polymerase II subunit RPB7.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 17 | G | 171 | 1340 | 861 | 222 | 249 | 8 | 0 | 0 |

- Molecule 18 is a protein called FACT complex subunit POB3.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 18 | O | 427 | 3481 | 2204 | 606 | 665 | 6 | 0 | 0 |

- Molecule 19 is a protein called FACT complex subunit SPT16.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 19 | Q | 404 | 3264 | 2074 | 555 | 624 | 11 | 0 | 0 |

- Molecule 20 is a protein called Histone H3.2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 20 | a | 88 | 720 | 456 | 135 | 126 | 3 | 0 | 0 |
| 20 | e | 94 | 775 | 490 | 149 | 133 | 3 | 0 | 0 |

There are 2 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------|------------|
| a | 102 | ALA | GLY | conflict | UNP P84233 |
| e | 102 | ALA | GLY | conflict | UNP P84233 |

- Molecule 21 is a protein called Histone H4.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 21 | b | 83 | 662 | 418 | 129 | 114 | 1 | 0 | 0 |
| 21 | f | 78 | 619 | 391 | 120 | 107 | 1 | 0 | 0 |

- Molecule 22 is a protein called Histone H2A type 1.

| Mol | Chain | Residues | Atoms | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---------|-------|
| | | | Total | C | N | O | | |
| 22 | c | 102 | 786 | 495 | 153 | 138 | 0 | 0 |
| 22 | g | 97 | 747 | 468 | 148 | 131 | 0 | 0 |

There are 4 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------|------------|
| c | 99 | ARG | GLY | conflict | UNP P06897 |
| c | 123 | SER | ALA | conflict | UNP P06897 |
| g | 99 | ARG | GLY | conflict | UNP P06897 |
| g | 123 | SER | ALA | conflict | UNP P06897 |

- Molecule 23 is a protein called Histone H2B 1.1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 23 | d | 95 | 745 | 469 | 134 | 140 | 2 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 23 | h | 93 | 726 | 457 | 130 | 137 | 2 | 0 | 0 |

There are 4 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|-----------------------|------------|
| d | 0 | MET | - | initiating methionine | UNP P02281 |
| d | 29 | THR | SER | conflict | UNP P02281 |
| h | 0 | MET | - | initiating methionine | UNP P02281 |
| h | 29 | THR | SER | conflict | UNP P02281 |

- Molecule 24 is ZINC ION (three-letter code: ZN) (formula: Zn).

| Mol | Chain | Residues | Atoms | | AltConf |
|-----|-------|----------|------------|---------|---------|
| 24 | A | 2 | Total 2 | Zn 2 | 0 |
| 24 | B | 1 | Total 1 | Zn 1 | 0 |
| 24 | C | 1 | Total 1 | Zn 1 | 0 |
| 24 | I | 2 | Total 2 | Zn 2 | 0 |
| 24 | J | 1 | Total 1 | Zn 1 | 0 |
| 24 | L | 1 | Total 1 | Zn 1 | 0 |
| 24 | Y | 1 | Total 1 | Zn 1 | 0 |


- Molecule 25 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

| Mol | Chain | Residues | Atoms | | AltConf |
|-----|-------|----------|------------|---------|---------|
| 25 | A | 1 | Total 1 | Mg 1 | 0 |


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA (5'-R(P*UP*CP*UP*UP*UP*UP*AP*UP*UP*UP*UP*UP*UP*CP*UP*G)-3')




- Molecule 2: DNA (138-MER)




- Molecule 3: DNA (148-MER)





- Molecule 4: DNA-directed RNA polymerase II subunit RPB1







• Molecule 5: DNA-directed RNA polymerase II subunit RPB2




• Molecule 6: DNA-directed RNA polymerase II subunit RPB3




• Molecule 7: DNA-directed RNA polymerases I, II, and III subunit RPABC1




• Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC2




- Molecule 9: DNA-directed RNA polymerases I, II, and III subunit RPABC3




- Molecule 10: DNA-directed RNA polymerase II subunit RPB9




- Molecule 11: DNA-directed RNA polymerases I, II, and III subunit RPABC5




- Molecule 12: DNA-directed RNA polymerase II subunit RPB11




- Molecule 13: DNA-directed RNA polymerases I, II, and III subunit RPABC4




- Molecule 14: Chromatin elongation factor SPT4





- Molecule 15: Transcription elongation factor SPT5











• Molecule 17: DNA-directed RNA polymerase II subunit RPB7

Chain G: 39% 99%




• Molecule 18: FACT complex subunit POB3

Chain O: 34% 76% 23%




• Molecule 19: FACT complex subunit SPT16


Chain Q: 19% 39% 61%




• Molecule 20: Histone H3.2




• Molecule 20: Histone H3.2




• Molecule 21: Histone H4





• Molecule 21: Histone H4





• Molecule 22: Histone H2A type 1





• Molecule 22: Histone H2A type 1



• Molecule 23: Histone H2B 1.1



• Molecule 23: Histone H2B 1.1



4 Experimental information

| Property | Value | Source |
|--------------------------------------|---|-----------|
| EM reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, Not provided | |
| Number of particles used | 47138 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 40.25 | Depositor |
| Minimum defocus (nm) | Not provided | |
| Maximum defocus (nm) | Not provided | |
| Magnification | Not provided | |
| Image detector | GATAN K3 BIOQUANTUM (6k x 4k) | Depositor |
| Maximum map value | 0.041 | Depositor |
| Minimum map value | -0.014 | Depositor |
| Average map value | -0.000 | Depositor |
| Map value standard deviation | 0.001 | Depositor |
| Recommended contour level | 0.0023 | Depositor |
| Map size (Å) | 419.99997, 419.99997, 419.99997 | wwPDB |
| Map dimensions | 400, 400, 400 | wwPDB |
| Map angles (°) | 90.0, 90.0, 90.0 | wwPDB |
| Pixel spacing (Å) | 1.05, 1.05, 1.05 | Depositor |

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|-----------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | P | 0.25 | 0/358 | 0.79 | 0/552 |
| 2 | N | 0.75 | 0/3189 | 1.03 | 1/4922 (0.0%) |
| 3 | T | 0.79 | 0/3392 | 1.02 | 4/5229 (0.1%) |
| 4 | A | 0.32 | 0/11289 | 0.66 | 3/15263 (0.0%) |
| 5 | B | 0.32 | 0/9037 | 0.67 | 2/12184 (0.0%) |
| 6 | C | 0.34 | 0/2124 | 0.65 | 1/2879 (0.0%) |
| 7 | E | 0.32 | 0/1788 | 0.67 | 0/2406 |
| 8 | F | 0.32 | 0/717 | 0.71 | 0/967 |
| 9 | H | 0.33 | 0/1086 | 0.72 | 1/1470 (0.1%) |
| 10 | I | 0.34 | 0/989 | 0.80 | 1/1331 (0.1%) |
| 11 | J | 0.36 | 0/541 | 0.71 | 0/727 |
| 12 | K | 0.33 | 0/938 | 0.68 | 0/1267 |
| 13 | L | 0.28 | 0/353 | 0.74 | 0/468 |
| 14 | Y | 0.31 | 0/755 | 0.68 | 0/1021 |
| 15 | Z | 0.34 | 0/1271 | 0.71 | 0/1702 |
| 16 | D | 0.36 | 0/1454 | 0.74 | 0/1949 |
| 17 | G | 0.31 | 0/1368 | 0.68 | 1/1844 (0.1%) |
| 18 | O | 0.33 | 0/3550 | 0.70 | 1/4788 (0.0%) |
| 19 | Q | 0.33 | 0/3329 | 0.70 | 1/4501 (0.0%) |
| 20 | a | 0.36 | 0/728 | 0.76 | 0/977 |
| 20 | e | 0.30 | 0/786 | 0.70 | 0/1055 |
| 21 | b | 0.36 | 0/669 | 0.74 | 0/894 |
| 21 | f | 0.30 | 0/626 | 0.69 | 0/837 |
| 22 | c | 0.31 | 0/796 | 0.65 | 0/1077 |
| 22 | g | 0.29 | 0/756 | 0.68 | 0/1020 |
| 23 | d | 0.33 | 0/756 | 0.67 | 0/1015 |
| 23 | h | 0.31 | 0/737 | 0.61 | 0/993 |
| All | All | 0.41 | 0/53382 | 0.74 | 16/73338 (0.0%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected

by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 4 | A | 0 | 3 |
| 6 | C | 0 | 1 |
| 8 | F | 0 | 1 |
| 9 | H | 0 | 1 |
| 19 | Q | 0 | 1 |
| 20 | a | 0 | 1 |
| All | All | 0 | 8 |

There are no bond length outliers.

All (16) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|------|------|------------|------|-------------|----------|
| 18 | O | 293 | ASP | CB-CG-OD1 | 7.25 | 124.83 | 118.30 |
| 4 | A | 1157 | ASP | CB-CG-OD1 | 6.65 | 124.29 | 118.30 |
| 2 | N | 16 | DA | O4'-C1'-N9 | 6.08 | 112.25 | 108.00 |
| 19 | Q | 947 | ASP | CB-CG-OD1 | 5.97 | 123.67 | 118.30 |
| 3 | T | 52 | DA | O4'-C1'-N9 | 5.93 | 112.15 | 108.00 |
| 6 | C | 217 | ASP | CB-CG-OD1 | 5.74 | 123.46 | 118.30 |
| 17 | G | 166 | ASP | CB-CG-OD1 | 5.65 | 123.38 | 118.30 |
| 5 | B | 320 | ASP | CB-CG-OD1 | 5.56 | 123.30 | 118.30 |
| 5 | B | 576 | ASP | CB-CG-OD1 | 5.54 | 123.29 | 118.30 |
| 10 | I | 61 | ASP | CB-CG-OD1 | 5.53 | 123.28 | 118.30 |
| 9 | H | 86 | ASP | CB-CG-OD1 | 5.47 | 123.22 | 118.30 |
| 4 | A | 62 | ASP | CB-CG-OD1 | 5.45 | 123.21 | 118.30 |
| 4 | A | 547 | LEU | CA-CB-CG | 5.11 | 127.04 | 115.30 |
| 3 | T | 25 | DG | O4'-C1'-N9 | 5.10 | 111.57 | 108.00 |
| 3 | T | -53 | DG | P-O3'-C3' | 5.09 | 125.81 | 119.70 |
| 3 | T | -2 | DC | O4'-C1'-N1 | 5.02 | 111.52 | 108.00 |

There are no chirality outliers.

All (8) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 4 | A | 12 | ARG | Sidechain |
| 4 | A | 524 | VAL | Peptide |
| 4 | A | 782 | ARG | Sidechain |
| 6 | C | 34 | ARG | Sidechain |
| 8 | F | 135 | ARG | Sidechain |
| 9 | H | 130 | ARG | Sidechain |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 19 | Q | 738 | VAL | Peptide |
| 20 | a | 49 | ARG | Sidechain |

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 4 | A | 1399/1733 (81%) | 1330 (95%) | 69 (5%) | 0 | 100 | 100 |
| 5 | B | 1102/1224 (90%) | 1056 (96%) | 46 (4%) | 0 | 100 | 100 |
| 6 | C | 263/318 (83%) | 248 (94%) | 15 (6%) | 0 | 100 | 100 |
| 7 | E | 212/215 (99%) | 206 (97%) | 6 (3%) | 0 | 100 | 100 |
| 8 | F | 85/155 (55%) | 80 (94%) | 5 (6%) | 0 | 100 | 100 |
| 9 | H | 129/146 (88%) | 120 (93%) | 9 (7%) | 0 | 100 | 100 |
| 10 | I | 117/122 (96%) | 105 (90%) | 12 (10%) | 0 | 100 | 100 |
| 11 | J | 63/70 (90%) | 61 (97%) | 2 (3%) | 0 | 100 | 100 |
| 12 | K | 113/120 (94%) | 109 (96%) | 4 (4%) | 0 | 100 | 100 |
| 13 | L | 42/70 (60%) | 38 (90%) | 4 (10%) | 0 | 100 | 100 |
| 14 | Y | 96/102 (94%) | 92 (96%) | 4 (4%) | 0 | 100 | 100 |
| 15 | Z | 152/1064 (14%) | 143 (94%) | 9 (6%) | 0 | 100 | 100 |
| 16 | D | 176/221 (80%) | 162 (92%) | 14 (8%) | 0 | 100 | 100 |
| 17 | G | 169/171 (99%) | 162 (96%) | 7 (4%) | 0 | 100 | 100 |
| 18 | O | 419/552 (76%) | 390 (93%) | 29 (7%) | 0 | 100 | 100 |
| 19 | Q | 398/1035 (38%) | 367 (92%) | 31 (8%) | 0 | 100 | 100 |
| 20 | a | 86/136 (63%) | 86 (100%) | 0 | 0 | 100 | 100 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 20 | e | 92/136 (68%) | 91 (99%) | 1 (1%) | 0 | 100 | 100 |
| 21 | b | 81/103 (79%) | 80 (99%) | 1 (1%) | 0 | 100 | 100 |
| 21 | f | 76/103 (74%) | 73 (96%) | 3 (4%) | 0 | 100 | 100 |
| 22 | c | 100/130 (77%) | 97 (97%) | 3 (3%) | 0 | 100 | 100 |
| 22 | g | 95/130 (73%) | 95 (100%) | 0 | 0 | 100 | 100 |
| 23 | d | 93/123 (76%) | 92 (99%) | 1 (1%) | 0 | 100 | 100 |
| 23 | h | 91/123 (74%) | 90 (99%) | 1 (1%) | 0 | 100 | 100 |
| All | All | 5649/8302 (68%) | 5373 (95%) | 276 (5%) | 0 | 100 | 100 |

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|-------------|----------|-------------|-----|
| 4 | A | 1233/1520 (81%) | 1230 (100%) | 3 (0%) | 93 | 98 |
| 5 | B | 967/1061 (91%) | 966 (100%) | 1 (0%) | 93 | 98 |
| 6 | C | 233/274 (85%) | 232 (100%) | 1 (0%) | 91 | 95 |
| 7 | E | 196/197 (100%) | 195 (100%) | 1 (0%) | 88 | 95 |
| 8 | F | 77/137 (56%) | 77 (100%) | 0 | 100 | 100 |
| 9 | H | 117/128 (91%) | 116 (99%) | 1 (1%) | 78 | 91 |
| 10 | I | 113/116 (97%) | 113 (100%) | 0 | 100 | 100 |
| 11 | J | 60/65 (92%) | 60 (100%) | 0 | 100 | 100 |
| 12 | K | 99/102 (97%) | 99 (100%) | 0 | 100 | 100 |
| 13 | L | 39/57 (68%) | 39 (100%) | 0 | 100 | 100 |
| 14 | Y | 82/87 (94%) | 82 (100%) | 0 | 100 | 100 |
| 15 | Z | 136/877 (16%) | 135 (99%) | 1 (1%) | 84 | 94 |
| 16 | D | 161/200 (80%) | 159 (99%) | 2 (1%) | 71 | 88 |
| 17 | G | 152/152 (100%) | 152 (100%) | 0 | 100 | 100 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|-------------|----------|-------------|-----|
| 18 | O | 382/489 (78%) | 378 (99%) | 4 (1%) | 76 | 90 |
| 19 | Q | 365/935 (39%) | 364 (100%) | 1 (0%) | 92 | 96 |
| 20 | a | 77/111 (69%) | 75 (97%) | 2 (3%) | 46 | 76 |
| 20 | e | 82/111 (74%) | 81 (99%) | 1 (1%) | 71 | 88 |
| 21 | b | 68/79 (86%) | 68 (100%) | 0 | 100 | 100 |
| 21 | f | 63/79 (80%) | 63 (100%) | 0 | 100 | 100 |
| 22 | c | 81/102 (79%) | 81 (100%) | 0 | 100 | 100 |
| 22 | g | 75/102 (74%) | 75 (100%) | 0 | 100 | 100 |
| 23 | d | 81/103 (79%) | 81 (100%) | 0 | 100 | 100 |
| 23 | h | 79/103 (77%) | 78 (99%) | 1 (1%) | 69 | 87 |
| All | All | 5018/7187 (70%) | 4999 (100%) | 19 (0%) | 91 | 95 |

All (19) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 4 | A | 200 | ARG |
| 4 | A | 393 | ARG |
| 4 | A | 420 | ARG |
| 5 | B | 1129 | ARG |
| 6 | C | 199 | LYS |
| 7 | E | 192 | ARG |
| 9 | H | 130 | ARG |
| 15 | Z | 317 | LYS |
| 16 | D | 36 | LYS |
| 16 | D | 121 | LYS |
| 18 | O | 18 | ARG |
| 18 | O | 42 | ARG |
| 18 | O | 271 | LYS |
| 18 | O | 368 | ARG |
| 19 | Q | 552 | ARG |
| 20 | a | 52 | ARG |
| 20 | a | 55 | GLN |
| 20 | e | 116 | ARG |
| 23 | h | 30 | ARG |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 4 | A | 1011 | GLN |
| 5 | B | 763 | GLN |
| 5 | B | 1025 | HIS |
| 5 | B | 1187 | ASN |
| 5 | B | 1205 | GLN |
| 19 | Q | 682 | GLN |

5.3.3 RNA [i](#)

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers |
|-----|-------|-------------|-------------------|-----------------|
| 1 | P | 15/16 (93%) | 5 (33%) | 1 (6%) |

All (5) RNA backbone outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | P | -4 | C |
| 1 | P | 0 | U |
| 1 | P | 1 | A |
| 1 | P | 8 | C |
| 1 | P | 10 | G |

All (1) RNA pucker outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | P | -1 | U |

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

| Mol | Chain | Number of breaks |
|-----|-------|------------------|
| 18 | O | 3 |
| 2 | N | 1 |

All chain breaks are listed below:

| Model | Chain | Residue-1 | Atom-1 | Residue-2 | Atom-2 | Distance (Å) |
|-------|-------|-----------|--------|-----------|--------|--------------|
| 1 | N | -65:DG | O3' | -54:DT | P | 30.81 |
| 1 | O | 241:ALA | C | 242:ILE | N | 5.38 |
| 1 | O | 239:GLY | C | 240:ASP | N | 5.32 |
| 1 | O | 237:VAL | C | 238:ALA | N | 3.21 |


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-12450. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 201



Y Index: 194




Z Index: 212

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0023. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 679 nm^3 ; this corresponds to an approximate mass of 613 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i



*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-12450 and PDB model 7NKY. Per-residue inclusion information can be found in section [3](#) on page [9](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0023 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0023).

























































9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0023) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion | Q-score |
|-------|--|--|
| All |  0.8459 |  0.3580 |
| A |  0.9506 |  0.4510 |
| B |  0.9661 |  0.4630 |
| C |  0.9495 |  0.4760 |
| D |  0.4676 |  0.2060 |
| E |  0.9555 |  0.4440 |
| F |  0.9389 |  0.4840 |
| G |  0.5121 |  0.2440 |
| H |  0.8913 |  0.4460 |
| I |  0.8256 |  0.3360 |
| J |  0.9807 |  0.4940 |
| K |  0.9261 |  0.4840 |
| L |  0.8997 |  0.3860 |
| N |  0.9121 |  0.2540 |
| O |  0.4988 |  0.1010 |
| P |  0.8062 |  0.2940 |
| Q |  0.4502 |  0.0840 |
| T |  0.9216 |  0.2670 |
| Y |  0.6442 |  0.1750 |
| Z |  0.7365 |  0.2450 |
| a |  0.9296 |  0.3960 |
| b |  0.9151 |  0.4240 |
| c |  0.9382 |  0.4350 |
| d |  0.9093 |  0.4130 |
| e |  0.9558 |  0.4050 |
| f |  0.9748 |  0.4390 |
| g |  0.9141 |  0.3900 |
| h |  0.9041 |  0.3910 |

