

# Full wwPDB X-ray Structure Validation Report (i)

#### Aug 8, 2023 – 08:55 PM EDT

PDB ID	:	1NL2
Title	:	BOVINE PROTHROMBIN FRAGMENT 1 IN COMPLEX WITH CALCIUM
		AND LYSOPHOSPHOTIDYLSERINE
Authors	:	Huang, M.; Huang, G.; Furie, B.; Seaton, B.; Furie, B.C.
Deposited on		
Resolution	:	2.30  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
$\mathrm{EDS}$	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain					
1	А	146	62%	32% 7%				
2	В	2	50%	50%				

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	CGU	А	7	-	-	Х	-
6	LPS	А	451	Х	-	-	-



# 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 1323 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Prothrombin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	146	Total 1179	C 708	N 209	0 251	S 11	0	1	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	7	CGU	GLU	modified residue	UNP P00735
А	8	CGU	GLU	modified residue	UNP P00735
А	15	CGU	GLU	modified residue	UNP P00735
А	17	CGU	GLU	modified residue	UNP P00735
А	20	CGU	GLU	modified residue	UNP P00735
А	21	CGU	GLU	modified residue	UNP P00735
А	26	CGU	GLU	modified residue	UNP P00735
А	27	CGU	GLU	modified residue	UNP P00735
А	30	CGU	GLU	modified residue	UNP P00735
А	33	CGU	GLU	modified residue	UNP P00735


• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



[	Mol	Chain	Residues	A	Atom	IS		ZeroOcc	AltConf	Trace
	2	В	2	Total 28	C 16	N 2	O 10	0	0	0

• Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).





Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
3	А	1	Total 14	C 8	N 1	O 5	0	0

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	7	Total Ca 7 7	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

I	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	5	А	3	Total Cl 3 3	0	0

• Molecule 6 is O-{HYDROXY[((2R)-2-HYDROXY-3-{[(1S)-1-HYDROXYPENTADECY L]OXY}PROPYL)OXY]PHOSPHORYL}-L-SERINE (three-letter code: LPS) (formula: C<sub>21</sub>H<sub>44</sub>NO<sub>9</sub>P).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
6	٨	1	Total	С	Ν	0	Р	0	0
0	A	1	32	21	1	9	1	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	60	Total         O           60         60	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Prothrombin



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:	50%	50%
NAG2 NAG2		



# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	38.89Å 53.22Å 128.96Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	40.98 - 2.30	Depositor
% Data completeness	89.5 (40.98-2.30)	Depositor
(in resolution range)	00.0 (40.00 2.00)	Depositor
$R_{merge}$	0.09	Depositor
R <sub>sym</sub>	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
$R, R_{free}$	0.236 , $0.274$	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1323	wwPDB-VP
Average B, all atoms $(Å^2)$	46.0	wwPDB-VP



# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CGU, NAG, LPS, CL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.37	1/1081~(0.1%)	0.61	0/1450

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
1	А	76	GLY	C-N	5.39	1.46	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1179	0	1041	47	15
2	В	28	0	25	1	0
3	А	14	0	12	0	0
4	А	7	0	0	0	0
5	А	3	0	0	0	1
6	А	32	0	40	2	0
7	А	60	0	0	3	8
All	All	1323	0	1118	48	16

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 20.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A + 1	A +	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:99:GLU:HG2	1:A:135:ARG:HH21	1.33	0.93
1:A:3:LYS:NZ	1:A:7:CGU:HB2	1.85	0.92
1:A:70:VAL:HG13	1:A:117:ASN:ND2	2.03	0.73
1:A:3:LYS:HZ2	1:A:7:CGU:HB2	1.51	0.72
1:A:99:GLU:CG	1:A:135:ARG:HH21	2.03	0.71
1:A:3:LYS:HZ3	1:A:7:CGU:HB2	1.54	0.69
1:A:93:ARG:HG2	1:A:97:LYS:HG2	1.76	0.67
1:A:99:GLU:HG2	1:A:135:ARG:NH2	2.11	0.61
1:A:96:HIS:O	1:A:98:PRO:HD3	2.00	0.60
1:A:52:ARG:HB3	1:A:52:ARG:HH11	1.67	0.59
1:A:141:VAL:HG13	1:A:142:PRO:HD2	1.85	0.58
1:A:16:ARG:HD2	1:A:17:CGU:OE22	2.02	0.58
1:A:51:ALA:HB1	1:A:57:LYS:HG2	1.87	0.57
1:A:113:ASN:HD22	1:A:113:ASN:H	1.52	0.55
1:A:17:CGU:O	1:A:23:CYS:HB3	2.08	0.53
1:A:104:THR:OG1	1:A:105:HIS:HD2	1.92	0.52
1:A:52:ARG:HB3	1:A:52:ARG:NH1	2.25	0.52
1:A:40:ALA:O	1:A:44:LYS:HG3	2.11	0.50
1:A:51:ALA:HA	1:A:57:LYS:HE3	1.93	0.50
1:A:93:ARG:CG	1:A:97:LYS:HG2	2.41	0.50
1:A:7:CGU:N	1:A:7:CGU:OE21	2.45	0.49
1:A:96:HIS:CE1	1:A:135:ARG:HA	2.48	0.49
1:A:74:TYR:CZ	1:A:76:GLY:HA3	2.48	0.48
1:A:111:ARG:HB3	1:A:111:ARG:HH11	1.79	0.48
1:A:14:LEU:HD22	1:A:32:LEU:HD21	1.96	0.47
1:A:2:ASN:HA	1:A:7:CGU:OE12	2.15	0.47
1:A:91:ARG:HG2	7:A:552:HOH:O	2.14	0.46
1:A:111:ARG:HH12	1:A:114:PHE:HE2	1.64	0.46
1:A:25:ARG:NH1	7:A:547:HOH:O	2.45	0.46
1:A:74:TYR:CD1	1:A:141:VAL:HG11	2.51	0.46
1:A:111:ARG:CB	1:A:111:ARG:NH1	2.80	0.45
1:A:100:ILE:HG23	1:A:105:HIS:CD2	2.51	0.45
1:A:111:ARG:HH11	1:A:111:ARG:CB	2.30	0.45
1:A:34:SER:N	7:A:555:HOH:O	2.50	0.45
1:A:3:LYS:HE3	6:A:451:LPS:H321	2.00	0.43
1:A:55:ARG:HG3	1:A:59:ASN:ND2	2.34	0.43
1:A:99:GLU:H	1:A:99:GLU:HG3	1.48	0.43
1:A:111:ARG:NH1	1:A:111:ARG:HB2	2.35	0.42



Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:A:70:VAL:HG13	1:A:117:ASN:HD22	1.82	0.42
1:A:70:VAL:HG13	1:A:70:VAL:O	2.20	0.42
1:A:3:LYS:HD3	1:A:7:CGU:CD2	2.50	0.42
1:A:113:ASN:HD22	1:A:113:ASN:N	2.14	0.41
1:A:7:CGU:HA	6:A:451:LPS:H2	2.02	0.41
1:A:25:ARG:NH2	1:A:39:ASP:OD1	2.54	0.40
1:A:122:ILE:HD12	1:A:122:ILE:HA	1.78	0.40
2:B:2:NAG:O3	2:B:2:NAG:H83	2.22	0.40
1:A:2:ASN:C	1:A:4:GLY:H	2.24	0.40

Continued from previous page...

All (16) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic	Clash
1100111-1	7100m-2	distance $(Å)$	overlap (Å)
1:A:56[B]:GLU:OE1	7:A:541:HOH:O[4_567]	1.01	1.19
1:A:2:ASN:ND2	7:A:558:HOH:O[4_467]	1.51	0.69
1:A:93:ARG:NE	1:A:141:VAL:O[3_656]	1.63	0.57
7:A:502:HOH:O	7:A:541:HOH:O[4_567]	1.71	0.49
1:A:146:GLN:CB	5:A:251:CL:CL[3_646]	1.92	0.28
1:A:2:ASN:ND2	7:A:509:HOH:O[4_467]	1.93	0.27
1:A:95:PRO:CA	7:A:549:HOH:O[3_656]	1.97	0.23
1:A:93:ARG:CD	1:A:141:VAL:O[3_656]	1.98	0.22
1:A:97:LYS:CE	1:A:146:GLN:O[3_656]	1.98	0.22
1:A:95:PRO:CB	7:A:549:HOH:O[3_656]	2.01	0.19
1:A:27:CGU:OE12	7:A:509:HOH:O[4_467]	2.05	0.15
1:A:97:LYS:NZ	1:A:146:GLN:O[3_656]	2.05	0.15
1:A:93:ARG:O	1:A:143:VAL:CG2[3_656]	2.08	0.12
1:A:69:GLY:O	1:A:94:TYR:CB[3_646]	2.13	0.07
1:A:94:TYR:OH	1:A:141:VAL:CB[3_656]	2.14	0.06
1:A:56[B]:GLU:CD	7:A:541:HOH:O[4_567]	2.19	0.01

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	135/146~(92%)	127~(94%)	7 (5%)	1 (1%)	22 26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	5	PHE

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	116/115~(101%)	108~(93%)	8 (7%)	15 20

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	3	LYS
1	А	70	VAL
1	А	93	ARG
1	А	99	GLU
1	А	113	ASN
1	А	129	THR
1	А	134	LEU
1	А	146	GLN

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such side chains are listed below:

Mol	Chain	Res	Type
1	А	105	HIS
1	А	113	ASN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	B	ond leng	$\operatorname{gths}$	В	ond ang	gles
N101	Type	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
1	CGU	А	15	1,4	9,11,12	1.06	0	9,14,16	1.12	1 (11%)
1	CGU	А	21	1,4	9,11,12	1.07	0	9,14,16	1.04	1 (11%)
1	CGU	А	30	1,4	9,11,12	1.01	0	9,14,16	1.09	1 (11%)
1	CGU	А	20	1,4	9,11,12	0.98	0	9,14,16	1.01	1 (11%)
1	CGU	А	17	1,4	9,11,12	0.98	0	9,14,16	1.02	1 (11%)
1	CGU	А	8	1,4	9,11,12	1.06	0	9,14,16	1.02	1 (11%)
1	CGU	А	27	1,4	9,11,12	1.04	0	9,14,16	1.06	2 (22%)
1	CGU	А	33	1	9,11,12	1.00	0	9,14,16	1.26	2 (22%)
1	CGU	А	7	1,4	9,11,12	1.00	0	9,14,16	1.01	1 (11%)
1	CGU	А	26	1,4	9,11,12	0.95	0	9,14,16	1.27	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CGU	А	15	1,4	-	2/13/14/16	-
1	CGU	А	21	1,4	-	1/13/14/16	-
1	CGU	А	30	1,4	-	2/13/14/16	-
1	CGU	А	20	1,4	-	3/13/14/16	-
1	CGU	А	17	1,4	-	7/13/14/16	-
1	CGU	А	8	1,4	-	7/13/14/16	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CGU	А	27	1,4	-	8/13/14/16	-
1	CGU	А	33	1	-	4/13/14/16	-
1	CGU	А	7	1,4	-	4/13/14/16	-
1	CGU	А	26	1,4	-	3/13/14/16	-

Continued from previous page...

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	А	33	CGU	CB-CG-CD2	2.87	118.94	113.11
1	А	26	CGU	CB-CG-CD2	2.68	118.55	113.11
1	А	20	CGU	CB-CG-CD2	2.59	118.37	113.11
1	А	7	CGU	CB-CG-CD2	2.59	118.37	113.11
1	А	15	CGU	CB-CG-CD2	2.57	118.33	113.11
1	А	30	CGU	CB-CG-CD2	2.49	118.17	113.11
1	А	21	CGU	CB-CG-CD2	2.47	118.13	113.11
1	А	26	CGU	CB-CG-CD1	2.37	117.92	113.11
1	А	33	CGU	CB-CG-CD1	2.24	117.67	113.11
1	А	27	CGU	CB-CG-CD2	2.14	117.46	113.11
1	А	8	CGU	CB-CG-CD2	2.11	117.39	113.11
1	А	17	CGU	CB-CG-CD2	2.06	117.31	113.11
1	А	27	CGU	CB-CG-CD1	2.06	117.30	113.11

There are no chirality outliers.

All	(41)	$\operatorname{torsion}$	outliers	are	listed	below:	
-----	------	--------------------------	----------	-----	--------	--------	--

Mol	Chain	Res	Type	Atoms
1	А	7	CGU	O-C-CA-CB
1	А	8	CGU	C-CA-CB-CG
1	А	8	CGU	CA-CB-CG-CD2
1	А	17	CGU	O-C-CA-CB
1	А	17	CGU	CA-CB-CG-CD2
1	А	26	CGU	CA-CB-CG-CD1
1	А	27	CGU	N-CA-CB-CG
1	А	27	CGU	CA-CB-CG-CD1
1	А	33	CGU	O-C-CA-CB
1	А	33	CGU	CA-CB-CG-CD2
1	А	15	CGU	CA-CB-CG-CD1
1	А	20	CGU	CA-CB-CG-CD2
1	А	17	CGU	C-CA-CB-CG



Mol	Chain	Res	Type	Atoms
1	А	20	CGU	C-CA-CB-CG
1	А	27	CGU	C-CA-CB-CG
1	А	30	CGU	C-CA-CB-CG
1	А	8	CGU	OE11-CD1-CG-CB
1	А	8	CGU	OE12-CD1-CG-CB
1	А	8	CGU	OE21-CD2-CG-CB
1	А	17	CGU	OE21-CD2-CG-CB
1	А	27	CGU	OE11-CD1-CG-CB
1	А	27	CGU	OE12-CD1-CG-CB
1	А	27	CGU	OE21-CD2-CG-CB
1	А	33	CGU	OE21-CD2-CG-CB
1	А	33	CGU	OE22-CD2-CG-CB
1	А	17	CGU	OE11-CD1-CG-CD2
1	А	17	CGU	OE12-CD1-CG-CD2
1	А	26	CGU	OE21-CD2-CG-CD1
1	А	26	CGU	OE22-CD2-CG-CD1
1	А	7	CGU	CA-CB-CG-CD1
1	А	7	CGU	OE21-CD2-CG-CB
1	А	7	CGU	OE22-CD2-CG-CB
1	А	8	CGU	N-CA-CB-CG
1	А	8	CGU	OE22-CD2-CG-CB
1	А	17	CGU	OE22-CD2-CG-CB
1	А	20	CGU	N-CA-CB-CG
1	А	27	CGU	OE22-CD2-CG-CB
1	А	30	CGU	N-CA-CB-CG
1	А	15	CGU	OE11-CD1-CG-CD2
1	А	21	CGU	OE12-CD1-CG-CD2
1	А	27	CGU	OE22-CD2-CG-CD1

Continued from previous page...

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	А	17	CGU	2	0
1	А	27	CGU	0	1
1	А	7	CGU	7	0

### 5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul



statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Res Link		Link	Bond lengths			Bond angles			
	Type	Unam	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
2	NAG	В	1	1,2	14,14,15	0.65	0	$17,\!19,\!21$	0.90	1 (5%)
2	NAG	В	2	2	14,14,15	0.76	0	17,19,21	0.89	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	В	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	В	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	В	1	NAG	C2-N2-C7	-2.86	118.83	122.90
2	В	2	NAG	C4-C3-C2	2.01	113.96	111.02

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	В	1	NAG	C8-C7-N2-C2
2	В	1	NAG	O7-C7-N2-C2
2	В	2	NAG	C8-C7-N2-C2
2	В	2	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	2	NAG	1	0





The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

### 5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 10 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	ain Res Link		Bo	ond leng	$\operatorname{ths}$	Bond angles		
WIOI	туре	e Chain Res Link	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
6	LPS	А	451	4	30,31,31	1.60	3 (10%)	32,37,37	2.45	4 (12%)



Mol	Mol Type Chain Res		Res Link		Bond lengths			Bond angles		
	Type	e Chain Res Lin		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2	
3	NAG	А	303	1	14,14,15	0.54	0	17,19,21	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	LPS	А	451	4	1/1/5/5	16/35/35/35	-
3	NAG	А	303	1	-	5/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
6	А	451	LPS	O32-C31	-6.03	1.22	1.39
6	А	451	LPS	C32-C31	3.44	1.58	1.50
6	А	451	LPS	O31-C31	-3.35	1.34	1.40

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$\mathbf{Ideal}(^{o})$
6	А	451	LPS	O32-C31-C32	9.06	125.90	109.12
6	А	451	LPS	O12-C11-C12	8.91	115.83	108.06
6	А	451	LPS	C33-C32-C31	3.31	121.04	113.38
6	А	451	LPS	C11-C12-C	2.31	115.41	110.93

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	А	451	LPS	C31

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	303	NAG	C8-C7-N2-C2
3	А	303	NAG	O7-C7-N2-C2
6	А	451	LPS	C12-C11-O12-P
6	А	451	LPS	C11-O12-P-O13
3	А	303	NAG	O5-C5-C6-O6
3	А	303	NAG	C4-C5-C6-O6
6	А	451	LPS	O11-C1-C2-C3



Mol	Chain	Res	Type	Atoms
6	А	451	LPS	C39-C40-C41-C42
6	А	451	LPS	C36-C37-C38-C39
6	А	451	LPS	C34-C35-C36-C37
6	А	451	LPS	O12-C11-C12-C
6	А	451	LPS	C41-C42-C43-C44
6	А	451	LPS	O21-C2-C3-O31
6	А	451	LPS	O32-C31-C32-C33
6	А	451	LPS	C35-C36-C37-C38
6	А	451	LPS	O12-C11-C12-N
6	А	451	LPS	C32-C31-O31-C3
6	А	451	LPS	O11-C1-C2-O21
6	А	451	LPS	C11-O12-P-O11
3	А	303	NAG	C3-C2-N2-C7
6	А	451	LPS	C11-O12-P-O14

Continued from previous page...


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	А	451	LPS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

#### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

#### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

