

# wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 – 03:07 PM EDT

PDB ID : 6NQC

Title: Crystal structure of a peptidase from an acI-B1 Actinobacterium

Authors: Forest, K.T.; Dwulit-Smith, J.R.; Satyshur, K.A.

Deposited on : 2019-01-20

Resolution : 1.94 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3934 atoms, of which 1868 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Cyanophycinase-like exopeptidase.

Mo	ol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1		A	244	Total 3766	C 1215	H 1868	N 323	O 357	S 3	0	1	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP J9H630
A	-19	GLY	-	expression tag	UNP J9H630
A	-18	SER	-	expression tag	UNP J9H630
A	-17	SER	-	expression tag	UNP J9H630
A	-16	HIS	-	expression tag	UNP J9H630
A	-15	HIS	_	expression tag	UNP J9H630
A	-14	HIS	-	expression tag	UNP J9H630
A	-13	HIS	-	expression tag	UNP J9H630
A	-12	HIS	-	expression tag	UNP J9H630
A	-11	HIS	-	expression tag	UNP J9H630
A	-10	SER	-	expression tag	UNP J9H630
A	-9	SER	-	expression tag	UNP J9H630
A	-8	GLY	-	expression tag	UNP J9H630
A	-7	GLU	-	expression tag	UNP J9H630
A	-6	ASN	-	expression tag	UNP J9H630
A	-5	LEU	-	expression tag	UNP J9H630
A	-4	TYR	-	expression tag	UNP J9H630
A	-3	PHE	-	expression tag	UNP J9H630
A	-2	GLN	-	expression tag	UNP J9H630
A	-1	GLY		expression tag	UNP J9H630
A	0	HIS	-	expression tag	UNP J9H630

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).





Mol	Chain	Residues	Residues Atoms			ZeroOcc	AltConf
2	A	1	Total 5	O 4	S 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	163	Total O 163 163	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



# 3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source	
Space group	C 2 2 21	Depositor	
Cell constants	44.69Å 88.92Å 131.31Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	26.41 - 1.94	Depositor	
% Data completeness	91.1 (26.41-1.94)	Depositor	
(in resolution range)	,		
$R_{merge}$	(Not available)	Depositor	
$R_{sym}$	(Not available)	Depositor	
$< I/\sigma(I) > 1$	3.13  (at  1.93Å)	Xtriage	
Refinement program	PHENIX (1.14_3260: ???)	Depositor	
$R, R_{free}$	0.164 , $0.227$	Depositor	
Wilson B-factor $(A^2)$	21.3	Xtriage	
Anisotropy	0.528	Xtriage	
L-test for twinning <sup>2</sup>	$ < L > = 0.49, < L^2> = 0.32$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	3934	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.28% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 4 Model quality (i)

## 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

## 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 4.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Type	Chain	Res	Link	B	ond leng	$\operatorname{gths}$	Bond angles		
	IVIOI				Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
ſ	2	SO4	A	301	-	4,4,4	0.25	0	6,6,6	0.36	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 4.7 Other polymers (i)

There are no such residues in this entry.

## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 5 Fit of model and data (i)

### 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

## 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

