



wwPDB EM Validation Summary Report ⓘ

Dec 19, 2022 – 02:35 am GMT

PDB ID : 7NVM
EMDB ID : EMD-12606
Title : Human TRiC complex in closed state with nanobody Nb18, actin and PhLP2A bound
Authors : Kelly, J.J.; Chi, G.; Bulawa, C.; Paavilainen, V.O.; Bountra, C.; Huiskonen, J.T.; Yue, W.; Structural Genomics Consortium (SGC)
Deposited on : 2021-03-15
Resolution : 3.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	556	5% (upper red bar), 96% (green), 2% (yellow), 3% (grey)
1	a	556	1% (upper red bar), 95% (green), 2% (yellow), 2% (grey)
2	B	535	5% (upper red bar), 97% (green), 2% (yellow), 2% (grey)
2	b	535	6% (upper red bar), 97% (green), 2% (yellow), 2% (grey)
3	D	539	5% (upper red bar), 96% (green), 2% (yellow), 2% (grey)
3	d	539	6% (upper red bar), 96% (green), 2% (yellow), 2% (grey)
4	E	541	1% (upper red bar), 98% (green), 2% (yellow), 2% (grey)
4	e	541	1% (upper red bar), 99% (green), 2% (yellow), 2% (grey)
5	G	545	1% (upper red bar), 96% (green), 2% (yellow), 2% (grey)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	g	545	<p>5% 98%</p>
6	H	543	<p>96%</p>
6	h	543	<p>96%</p>
7	N	129	<p>78% 92% 8%</p>
7	n	129	<p>79% 92% 8%</p>
8	Q	548	<p>97%</p>
8	q	548	<p>97%</p>
9	Z	531	<p>98%</p>
9	z	531	<p>99%</p>
10	K	375	<p>61% 80% 15%</p>
11	P	239	<p>71% 77% 21%</p>

2 Entry composition [i](#)

There are 15 unique types of molecules in this entry. The entry contains 71008 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-complex protein 1 subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	536	Total	C	N	O	S	0	0
			4069	2548	711	787	23		
1	a	532	Total	C	N	O	S	0	0
			4041	2533	707	778	23		

- Molecule 2 is a protein called T-complex protein 1 subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	526	Total	C	N	O	S	0	0
			3952	2473	696	764	19		
2	b	525	Total	C	N	O	S	0	0
			3943	2467	694	763	19		

- Molecule 3 is a protein called T-complex protein 1 subunit delta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	D	520	Total	C	N	O	S	0	0
			3923	2453	683	764	23		
3	d	520	Total	C	N	O	S	0	0
			3917	2450	680	764	23		

- Molecule 4 is a protein called T-complex protein 1 subunit epsilon.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	E	535	Total	C	N	O	S	1	0
			4132	2590	719	792	31		
4	e	540	Total	C	N	O	S	1	0
			4169	2610	724	804	31		

- Molecule 5 is a protein called T-complex protein 1 subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	G	526	Total	C	N	O	S	0	0
			4089	2548	726	785	30		
5	g	538	Total	C	N	O	S	0	0
			4169	2590	740	809	30		

- Molecule 6 is a protein called T-complex protein 1 subunit eta.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	H	528	Total	C	N	O	S	0	0
			4054	2561	699	769	25		
6	h	525	Total	C	N	O	S	0	0
			4032	2548	696	763	25		

- Molecule 7 is a protein called Nanobody Nb18.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	N	119	Total	C	N	O	S	0	0
			924	570	169	181	4		
7	n	119	Total	C	N	O	S	0	0
			924	570	169	181	4		

- Molecule 8 is a protein called T-complex protein 1 subunit theta.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	Q	538	Total	C	N	O	S	0	0
			4086	2579	696	784	27		
8	q	533	Total	C	N	O	S	0	0
			4053	2558	690	778	27		

- Molecule 9 is a protein called T-complex protein 1 subunit zeta.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	Z	525	Total	C	N	O	S	0	0
			4022	2528	704	769	21		
9	z	527	Total	C	N	O	S	0	0
			4033	2534	706	772	21		


- Molecule 10 is a protein called Actin, cytoplasmic 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	K	317	Total	C	N	O	S	0	0
			2459	1563	409	469	18		

- Molecule 11 is a protein called Phosducin-like protein 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	P	190	1486	955	239	287	5	0	0

- Molecule 12 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
12	A	1	27	10	5	10	2	0
12	B	1	27	10	5	10	2	0
12	D	1	27	10	5	10	2	0
12	E	1	27	10	5	10	2	0
12	G	1	27	10	5	10	2	0
12	H	1	27	10	5	10	2	0
12	Q	1	27	10	5	10	2	0
12	Z	1	27	10	5	10	2	0
12	a	1	27	10	5	10	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
12	b	1	27	10	5	10	2	0
12	d	1	27	10	5	10	2	0
12	e	1	27	10	5	10	2	0
12	g	1	27	10	5	10	2	0
12	h	1	27	10	5	10	2	0
12	q	1	27	10	5	10	2	0
12	z	1	27	10	5	10	2	0

- Molecule 13 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
13	A	1	1	1	0
13	B	1	1	1	0
13	D	1	1	1	0
13	E	1	1	1	0
13	G	1	1	1	0
13	H	1	1	1	0
13	Q	1	1	1	0
13	Z	1	1	1	0
13	a	1	1	1	0
13	b	1	1	1	0
13	d	1	1	1	0
13	e	1	1	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
13	g	1	Total 1	Mg 1	0
13	h	1	Total 1	Mg 1	0
13	q	1	Total 1	Mg 1	0
13	z	1	Total 1	Mg 1	0

- Molecule 14 is ALUMINUM FLUORIDE (three-letter code: AF3) (formula: AlF_3).



Mol	Chain	Residues	Atoms			AltConf
14	A	1	Total 4	Al 1	F 3	0
14	B	1	Total 4	Al 1	F 3	0
14	D	1	Total 4	Al 1	F 3	0
14	E	1	Total 4	Al 1	F 3	0
14	G	1	Total 4	Al 1	F 3	0
14	H	1	Total 4	Al 1	F 3	0
14	Q	1	Total 4	Al 1	F 3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			AltConf
			Total	Al	F	
14	Z	1	4	1	3	0
14	a	1	4	1	3	0
14	b	1	4	1	3	0
14	d	1	4	1	3	0
14	e	1	4	1	3	0
14	g	1	4	1	3	0
14	h	1	4	1	3	0
14	q	1	4	1	3	0
14	z	1	4	1	3	0

- Molecule 15 is water.

Mol	Chain	Residues	Atoms		AltConf
			Total	O	
15	A	2	2	2	0
15	B	2	2	2	0
15	D	1	1	1	0
15	E	1	1	1	0
15	G	1	1	1	0
15	H	1	1	1	0
15	Q	1	1	1	0
15	Z	1	1	1	0
15	a	2	2	2	0
15	b	1	1	1	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
15	d	1	Total 1	O 1	0
15	e	1	Total 1	O 1	0
15	g	1	Total 1	O 1	0
15	h	1	Total 1	O 1	0
15	q	1	Total 1	O 1	0
15	z	1	Total 1	O 1	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: T-complex protein 1 subunit alpha




- Molecule 1: T-complex protein 1 subunit alpha




- Molecule 2: T-complex protein 1 subunit beta




- Molecule 2: T-complex protein 1 subunit beta








• Molecule 3: T-complex protein 1 subunit delta




• Molecule 3: T-complex protein 1 subunit delta




• Molecule 4: T-complex protein 1 subunit epsilon




• Molecule 4: T-complex protein 1 subunit epsilon




• Molecule 5: T-complex protein 1 subunit gamma







● Molecule 8: T-complex protein 1 subunit theta





● Molecule 9: T-complex protein 1 subunit zeta




● Molecule 9: T-complex protein 1 subunit zeta








● Molecule 10: Actin, cytoplasmic 2





• Molecule 11: Phosducin-like protein 3



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	63082	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	43	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.148	Depositor
Minimum map value	-0.075	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.025	Depositor
Map size (\AA)	423.99997, 423.99997, 423.99997	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, AF3, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/4109	0.49	0/5548
1	a	0.31	0/4081	0.49	0/5510
2	B	0.30	0/3995	0.49	0/5386
2	b	0.31	0/3986	0.50	1/5375 (0.0%)
3	D	0.30	0/3955	0.48	0/5338
3	d	0.30	0/3949	0.50	0/5331
4	E	0.32	0/4183	0.51	0/5635
4	e	0.33	0/4220	0.51	0/5684
5	G	0.31	0/4136	0.50	0/5579
5	g	0.31	0/4215	0.49	0/5683
6	H	0.34	0/4111	0.49	0/5550
6	h	0.33	0/4089	0.48	0/5519
7	N	0.26	0/941	0.53	0/1270
7	n	0.28	0/941	0.54	0/1270
8	Q	0.33	0/4147	0.50	1/5606 (0.0%)
8	q	0.33	0/4112	0.49	0/5558
9	Z	0.31	0/4069	0.49	0/5486
9	z	0.31	0/4080	0.49	0/5501
10	K	0.29	0/2510	0.48	0/3401
11	P	0.26	0/1518	0.44	0/2062
All	All	0.31	0/71347	0.49	2/96292 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	e	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	b	316	ASP	CB-CG-OD2	5.20	122.98	118.30
8	Q	480	LEU	CA-CB-CG	5.03	126.88	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	e	8	ALA	Mainchain

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	534/556 (96%)	508 (95%)	26 (5%)	0	100	100
1	a	530/556 (95%)	502 (95%)	28 (5%)	0	100	100
2	B	524/535 (98%)	502 (96%)	21 (4%)	1 (0%)	47	79
2	b	523/535 (98%)	503 (96%)	19 (4%)	1 (0%)	47	79
3	D	518/539 (96%)	507 (98%)	11 (2%)	0	100	100
3	d	518/539 (96%)	487 (94%)	31 (6%)	0	100	100
4	E	534/541 (99%)	511 (96%)	20 (4%)	3 (1%)	25	59
4	e	539/541 (100%)	516 (96%)	22 (4%)	1 (0%)	47	79
5	G	524/545 (96%)	502 (96%)	22 (4%)	0	100	100
5	g	536/545 (98%)	519 (97%)	17 (3%)	0	100	100
6	H	526/543 (97%)	506 (96%)	20 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
6	h	523/543 (96%)	505 (97%)	18 (3%)	0	100	100
7	N	117/129 (91%)	110 (94%)	7 (6%)	0	100	100
7	n	117/129 (91%)	111 (95%)	6 (5%)	0	100	100
8	Q	536/548 (98%)	516 (96%)	18 (3%)	2 (0%)	34	69
8	q	531/548 (97%)	512 (96%)	18 (3%)	1 (0%)	47	79
9	Z	523/531 (98%)	504 (96%)	18 (3%)	1 (0%)	47	79
9	z	525/531 (99%)	505 (96%)	19 (4%)	1 (0%)	47	79
10	K	309/375 (82%)	275 (89%)	32 (10%)	2 (1%)	25	59
11	P	186/239 (78%)	169 (91%)	17 (9%)	0	100	100
All	All	9173/9548 (96%)	8770 (96%)	390 (4%)	13 (0%)	54	83

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	E	10	ASP
9	Z	16	ALA
4	e	10	ASP
8	q	530	ALA
9	z	16	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	447/463 (96%)	442 (99%)	5 (1%)	73	89
1	a	444/463 (96%)	441 (99%)	3 (1%)	84	93
2	B	418/427 (98%)	414 (99%)	4 (1%)	76	90
2	b	417/427 (98%)	413 (99%)	4 (1%)	76	90
3	D	442/452 (98%)	439 (99%)	3 (1%)	84	93
3	d	441/452 (98%)	440 (100%)	1 (0%)	93	97
4	E	452/456 (99%)	449 (99%)	3 (1%)	84	93

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	e	456/456 (100%)	452 (99%)	4 (1%)	78	91
5	G	456/469 (97%)	453 (99%)	3 (1%)	84	93
5	g	463/469 (99%)	461 (100%)	2 (0%)	91	96
6	H	435/443 (98%)	430 (99%)	5 (1%)	73	89
6	h	432/443 (98%)	427 (99%)	5 (1%)	71	88
7	N	96/105 (91%)	96 (100%)	0	100	100
7	n	96/105 (91%)	96 (100%)	0	100	100
8	Q	442/452 (98%)	440 (100%)	2 (0%)	88	94
8	q	438/452 (97%)	437 (100%)	1 (0%)	93	97
9	Z	437/442 (99%)	436 (100%)	1 (0%)	93	97
9	z	438/442 (99%)	436 (100%)	2 (0%)	88	94
10	K	264/318 (83%)	249 (94%)	15 (6%)	20	52
11	P	155/215 (72%)	148 (96%)	7 (4%)	27	60
All	All	7669/7951 (96%)	7599 (99%)	70 (1%)	79	91

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
10	K	325	MET
10	K	336	LYS
11	P	96	GLU
8	Q	537	SER
8	Q	369	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 123 such sidechains are listed below:

Mol	Chain	Res	Type
9	Z	438	GLN
9	z	17	GLN
2	b	285	HIS
9	z	8	ASN
10	K	111	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 48 ligands modelled in this entry, 16 are monoatomic - leaving 32 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
14	AF3	h	603	-	0,3,3	-	-	-		
12	ADP	D	601	13	24,29,29	0.93	1 (4%)	29,45,45	1.55	4 (13%)
14	AF3	Z	603	-	0,3,3	-	-	-		
14	AF3	E	603	-	0,3,3	-	-	-		
12	ADP	G	601	13	24,29,29	0.94	1 (4%)	29,45,45	1.19	3 (10%)
14	AF3	b	603	-	0,3,3	-	-	-		
12	ADP	H	601	13	24,29,29	0.97	1 (4%)	29,45,45	1.37	3 (10%)
12	ADP	Q	601	13	24,29,29	0.93	1 (4%)	29,45,45	1.39	3 (10%)
14	AF3	z	603	-	0,3,3	-	-	-		
14	AF3	Q	603	-	0,3,3	-	-	-		
12	ADP	q	601	13	24,29,29	0.95	1 (4%)	29,45,45	1.41	3 (10%)
12	ADP	a	601	13	24,29,29	0.97	1 (4%)	29,45,45	1.40	5 (17%)
14	AF3	q	603	-	0,3,3	-	-	-		
14	AF3	a	603	-	0,3,3	-	-	-		
12	ADP	Z	601	13	24,29,29	0.93	1 (4%)	29,45,45	1.53	5 (17%)
12	ADP	h	601	13	24,29,29	0.91	1 (4%)	29,45,45	1.33	4 (13%)
12	ADP	E	601	13	24,29,29	0.96	1 (4%)	29,45,45	1.41	6 (20%)
14	AF3	g	603	-	0,3,3	-	-	-		
12	ADP	e	601	13	24,29,29	0.93	1 (4%)	29,45,45	1.41	6 (20%)
12	ADP	b	601	13	24,29,29	0.92	1 (4%)	29,45,45	1.51	5 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
12	ADP	g	601	13	24,29,29	0.94	1 (4%)	29,45,45	1.33	3 (10%)
14	AF3	G	603	-	0,3,3	-	-	-	-	-
14	AF3	d	603	-	0,3,3	-	-	-	-	-
14	AF3	H	603	-	0,3,3	-	-	-	-	-
14	AF3	e	603	-	0,3,3	-	-	-	-	-
12	ADP	d	601	13	24,29,29	0.95	1 (4%)	29,45,45	1.61	5 (17%)
12	ADP	A	601	13	24,29,29	0.97	1 (4%)	29,45,45	1.44	5 (17%)
14	AF3	B	603	-	0,3,3	-	-	-	-	-
14	AF3	A	603	-	0,3,3	-	-	-	-	-
14	AF3	D	603	-	0,3,3	-	-	-	-	-
12	ADP	B	601	13	24,29,29	0.91	1 (4%)	29,45,45	1.50	5 (17%)
12	ADP	z	601	13	24,29,29	0.95	1 (4%)	29,45,45	1.53	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	ADP	H	601	13	-	6/12/32/32	0/3/3/3
12	ADP	Q	601	13	-	3/12/32/32	0/3/3/3
12	ADP	D	601	13	-	2/12/32/32	0/3/3/3
12	ADP	b	601	13	-	2/12/32/32	0/3/3/3
12	ADP	G	601	13	-	6/12/32/32	0/3/3/3
12	ADP	a	601	13	-	1/12/32/32	0/3/3/3
12	ADP	d	601	13	-	0/12/32/32	0/3/3/3
12	ADP	g	601	13	-	6/12/32/32	0/3/3/3
12	ADP	A	601	13	-	0/12/32/32	0/3/3/3
12	ADP	q	601	13	-	3/12/32/32	0/3/3/3
12	ADP	Z	601	13	-	0/12/32/32	0/3/3/3
12	ADP	h	601	13	-	6/12/32/32	0/3/3/3
12	ADP	e	601	13	-	2/12/32/32	0/3/3/3
12	ADP	E	601	13	-	3/12/32/32	0/3/3/3
12	ADP	B	601	13	-	2/12/32/32	0/3/3/3
12	ADP	z	601	13	-	0/12/32/32	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	z	601	ADP	C5-C4	2.40	1.47	1.40
12	E	601	ADP	C5-C4	2.32	1.47	1.40
12	a	601	ADP	C5-C4	2.31	1.47	1.40
12	g	601	ADP	C5-C4	2.30	1.47	1.40
12	G	601	ADP	C5-C4	2.28	1.47	1.40

The worst 5 of 70 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	Z	601	ADP	PA-O3A-PB	-4.32	118.00	132.83
12	z	601	ADP	PA-O3A-PB	-4.20	118.42	132.83
12	d	601	ADP	PA-O3A-PB	-3.73	120.03	132.83
12	Q	601	ADP	N3-C2-N1	-3.53	123.17	128.68
12	g	601	ADP	PA-O3A-PB	-3.51	120.77	132.83

There are no chirality outliers.



5 of 42 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
12	G	601	ADP	PA-O3A-PB-O3B
12	G	601	ADP	C5'-O5'-PA-O2A
12	H	601	ADP	PA-O3A-PB-O2B
12	H	601	ADP	PB-O3A-PA-O5'
12	H	601	ADP	C3'-C4'-C5'-O5'



There are no ring outliers.



No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






















5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-12606. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 169




Y Index: 186




Z Index: 166


6.3.2 Raw map



X Index: 173



Y Index: 212




Z Index: 231

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.025. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.5.1 emd_12606_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 321 nm³; this corresponds to an approximate mass of 290 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.10	-	-
Author-provided FSC curve	3.12	3.63	3.18
Unmasked-calculated*	3.81	6.89	3.91

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.81 differs from the reported value 3.1 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-12606 and PDB model 7NVM. Per-residue inclusion information can be found in section [3](#) on page [11](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.025 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.025).











































9.4 Atom inclusion [i](#)



At the recommended contour level, 70% of all backbone atoms, 69% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.025) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6949	 0.5290
A	 0.7382	 0.5380
B	 0.7219	 0.5310
D	 0.7330	 0.5420
E	 0.7547	 0.5440
G	 0.7389	 0.5360
H	 0.7797	 0.5680
K	 0.2582	 0.4280
N	 0.1696	 0.2840
P	 0.1826	 0.4180
Q	 0.7485	 0.5530
Z	 0.7352	 0.5380
a	 0.7466	 0.5420
b	 0.7289	 0.5270
d	 0.7340	 0.5380
e	 0.7381	 0.5420
g	 0.7282	 0.5360
h	 0.7747	 0.5630
n	 0.1529	 0.2660
q	 0.7667	 0.5560
z	 0.7322	 0.5320

