



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 30, 2023 – 04:10 AM EDT

PDB ID : 3NWA
Title : Glycoprotein B from Herpes simplex virus type 1, W174R mutant, low-pH
Authors : Stampfer, S.D.; Lou, H.; Cohen, G.H.; Eisenberg, R.J.; Heldwein, E.E.
Deposited on : 2010-07-09
Resolution : 2.26 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	703	 62% 23% 14%
1	B	703	 56% 29% 14%
1	C	703	 59% 26% 14%
1	D	703	 59% 26% 14%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 21541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	607	4896	3086	867	920	23	0	1	0
1	A	608	4899	3088	867	922	22	0	1	0
1	C	608	4883	3080	864	917	22	0	0	0
1	D	607	4900	3087	867	923	23	0	1	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	28	ASP	-	expression tag	UNP P06437
B	29	PRO	-	expression tag	UNP P06437
B	58	ALA	PRO	SEE REMARK 999	UNP P06437
B	174	ARG	TRP	engineered mutation	UNP P06437
B	313	SER	THR	SEE REMARK 999	UNP P06437
B	443	LEU	GLN	SEE REMARK 999	UNP P06437
A	28	ASP	-	expression tag	UNP P06437
A	29	PRO	-	expression tag	UNP P06437
A	58	ALA	PRO	SEE REMARK 999	UNP P06437
A	174	ARG	TRP	engineered mutation	UNP P06437
A	313	SER	THR	SEE REMARK 999	UNP P06437
A	443	LEU	GLN	SEE REMARK 999	UNP P06437
C	28	ASP	-	expression tag	UNP P06437
C	29	PRO	-	expression tag	UNP P06437
C	58	ALA	PRO	SEE REMARK 999	UNP P06437
C	174	ARG	TRP	engineered mutation	UNP P06437
C	313	SER	THR	SEE REMARK 999	UNP P06437
C	443	LEU	GLN	SEE REMARK 999	UNP P06437
D	28	ASP	-	expression tag	UNP P06437
D	29	PRO	-	expression tag	UNP P06437
D	58	ALA	PRO	SEE REMARK 999	UNP P06437

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	174	ARG	TRP	engineered mutation	UNP P06437
D	313	SER	THR	SEE REMARK 999	UNP P06437
D	443	LEU	GLN	SEE REMARK 999	UNP P06437

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is MESO-ERYTHRITOL (three-letter code: MRY) (formula: $C_4H_{10}O_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 8 4 4	0	0
3	B	1	Total C O 8 4 4	0	0
3	A	1	Total C O 8 4 4	0	0
3	C	1	Total C O 8 4 4	0	0
3	D	1	Total C O 8 4 4	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	444	Total O 444 444	0	0
4	A	478	Total O 478 478	0	0
4	C	462	Total O 462 462	0	0
4	D	483	Total O 483 483	0	0



● Molecule 1: Envelope glycoprotein B



● Molecule 1: Envelope glycoprotein B



ALA	THR	VAL	ALA	ALA	GLY	HIS	ALA	THR	LEU	ARG	GLU	HIS	LEU	ARG	D103	I104	K105	C116	P119	V124	Q129	P130	R131	R132	R136	P137	T143	E144	G145	I146	A147	E152	P156	D167	V168	T169	V170	S171	R174	H177	R178	Y179	S180	F186	E187	D188	R189																																		
D199	K200	A203	V206	C207	R208	A211	T221	H224	R225	T241	R242	T243	T250	Y254	Y255	P256	V259	E260	A261	Y265	T268	I272	R279	E286	Y296	F300	E305	G306	S307	H308	H311	T312	S313	D317	G324	R328	T332	K333	A334	R335	P339	N343	K349	A353	R360	P361	S362	V363	C364	T365	M366	E379	Y380	F384	R385	F386	S387	S388	I391	S392	T393	T394	L399	T400	E401	Y402	P403	L404	V407	D408	L409	G410	G414	R418	D419	R423	I424	F425	R428	Y429	T432
H433	I434	K435	Q438	P439	Q440	Y441	Y442	L443	A451	L455	L456	S457	N458	T459	L460	L463	R466	E467	H468	L469	R470	E471	Q472	S473	R474	K475	P476	P477	ASN	THR	PRO	PRO	PRO	PRO	GLY	ALA	SER	ALA	ASN	ALA	ALA	SER	VAL	E493	K496	S499	R499	P600	L601	Y602	S603	F604	R605	A604	Y606																										
N511	H512	I513	Q514	R515	M520	R523	V524	L531	H534	E535	N540	R543	K544	S552	V553	V559	S560	A561	R562	M563	L564	Y567	R568	A569	V570	A577	M580	V581	I582	M587	R588	I589	S590	S591	R592	P593	Y597	S599	R599	P600	L601	Y602	S603	F604	R605	A604	Y606																																		
E607	D608	Q609	V613	Q616	L617	E622	L623	R624	T634	V635	G636	H637	R638	R639	Y640	F650	E651	E652	Y655	L659	R660	R661	I664	H681	E682	F683	Y684	R691	S697	D701	E704	R708	H724	ALA	ASP	ALA	ALA	ALA	ALA	ALA																																									

4 Data and refinement statistics i

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	117.09Å 117.09Å 321.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.32 – 2.26 48.35 – 2.26	Depositor EDS
% Data completeness (in resolution range)	87.1 (47.32-2.26) 87.1 (48.35-2.26)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.27Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, R_{free}	0.172 , 0.227 0.171 , 0.227	Depositor DCC
R_{free} test set	11385 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtrriage
Anisotropy	0.493	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.020 for -h,-k,l 0.467 for h,-h-k,-l 0.021 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	21541	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 74.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4096e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MRY, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/5021	0.55	0/6822
1	B	0.37	0/5017	0.54	0/6814
1	C	0.37	0/5002	0.53	0/6798
1	D	0.38	0/5020	0.55	0/6820
All	All	0.37	0/20060	0.54	0/27254

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4899	0	4713	133	0
1	B	4896	0	4717	183	0
1	C	4883	0	4691	134	0
1	D	4900	0	4708	149	0
2	A	14	0	13	0	0
2	B	14	0	13	1	0
2	C	14	0	13	1	0
2	D	14	0	13	1	0
3	A	8	0	10	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	16	0	20	0	0
3	C	8	0	10	0	0
3	D	8	0	10	0	0
4	A	478	0	0	13	0
4	B	444	0	0	12	0
4	C	462	0	0	19	0
4	D	483	0	0	14	0
All	All	21541	0	18931	595	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 595 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:515:ARG:HG3	1:D:515:ARG:HH11	1.27	0.98
1:C:543:ARG:HB3	1:C:550:ILE:HG21	1.55	0.88
1:D:515:ARG:HG3	1:D:515:ARG:NH1	1.82	0.88
1:B:189:ARG:HB2	1:B:349:LYS:HE2	1.59	0.85
1:D:403:PRO:HG3	1:D:476:PRO:HB3	1.60	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	605/703 (86%)	558 (92%)	43 (7%)	4 (1%)	22 21
1	B	604/703 (86%)	555 (92%)	46 (8%)	3 (0%)	29 29
1	C	604/703 (86%)	543 (90%)	55 (9%)	6 (1%)	15 13
1	D	604/703 (86%)	571 (94%)	29 (5%)	4 (1%)	22 21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	2417/2812 (86%)	2227 (92%)	173 (7%)	17 (1%)	22 21

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	259	VAL
1	A	260	GLU
1	D	177	HIS
1	C	118	PRO
1	C	655	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	526/593 (89%)	504 (96%)	22 (4%)	30 34
1	B	526/593 (89%)	501 (95%)	25 (5%)	25 28
1	C	522/593 (88%)	503 (96%)	19 (4%)	35 42
1	D	527/593 (89%)	502 (95%)	25 (5%)	26 29
All	All	2101/2372 (89%)	2010 (96%)	91 (4%)	29 33

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	502	GLU
1	D	305	GLU
1	C	553	VAL
1	C	724	HIS
1	D	418	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	308	HIS
1	D	511	ASN
1	D	706	GLN
1	D	681	HIS
1	A	585	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1141	1	14,14,15	0.57	0	17,19,21	1.12	1 (5%)
3	MRY	C	2000	-	7,7,7	0.20	0	8,8,8	0.95	0
3	MRY	B	2000	-	7,7,7	0.35	0	8,8,8	0.78	0
2	NAG	A	1398	1	14,14,15	0.52	0	17,19,21	1.41	1 (5%)
2	NAG	D	1398	1	14,14,15	0.55	0	17,19,21	0.96	1 (5%)
2	NAG	B	1398	1	14,14,15	0.46	0	17,19,21	1.00	1 (5%)
3	MRY	D	2000	-	7,7,7	0.48	0	8,8,8	0.81	0
3	MRY	A	2000	-	7,7,7	0.49	0	8,8,8	0.65	0
3	MRY	B	3000	-	7,7,7	0.91	0	8,8,8	1.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1141	1	-	2/6/23/26	0/1/1/1
3	MRY	C	2000	-	-	0/8/8/8	-
3	MRY	B	2000	-	-	0/8/8/8	-
2	NAG	A	1398	1	-	0/6/23/26	0/1/1/1
2	NAG	D	1398	1	-	2/6/23/26	0/1/1/1
2	NAG	B	1398	1	-	2/6/23/26	0/1/1/1
3	MRY	D	2000	-	-	0/8/8/8	-
3	MRY	A	2000	-	-	0/8/8/8	-
3	MRY	B	3000	-	-	0/8/8/8	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1398	NAG	C1-O5-C5	4.71	118.58	112.19
2	B	1398	NAG	C1-O5-C5	3.13	116.44	112.19
2	C	1141	NAG	O5-C1-C2	2.56	115.34	111.29
2	D	1398	NAG	C1-O5-C5	2.05	114.97	112.19

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1141	NAG	O5-C5-C6-O6
2	B	1398	NAG	O5-C5-C6-O6
2	D	1398	NAG	C8-C7-N2-C2
2	B	1398	NAG	C4-C5-C6-O6
2	D	1398	NAG	O7-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1141	NAG	1	0
2	D	1398	NAG	1	0
2	B	1398	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	608/703 (86%)	-0.28	1 (0%) 95 96	21, 33, 55, 75	0
1	B	607/703 (86%)	-0.22	6 (0%) 82 84	21, 36, 63, 90	0
1	C	608/703 (86%)	-0.22	5 (0%) 86 87	22, 35, 64, 88	0
1	D	607/703 (86%)	-0.28	1 (0%) 95 96	23, 34, 55, 88	0
All	All	2430/2812 (86%)	-0.25	13 (0%) 91 91	21, 34, 60, 90	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	463	LEU	3.1
1	D	476	PRO	2.9
1	B	492	VAL	2.9
1	C	464	TYR	2.9
1	A	465	VAL	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	D	1398	14/15	0.85	0.13	40,58,64,64	0
2	NAG	A	1398	14/15	0.86	0.25	49,65,76,81	0
2	NAG	C	1141	14/15	0.93	0.23	84,89,92,95	0
2	NAG	B	1398	14/15	0.93	0.09	46,59,65,66	0
3	MRY	B	2000	8/8	0.95	0.13	29,35,39,44	0
3	MRY	B	3000	8/8	0.95	0.10	39,41,43,47	0
3	MRY	A	2000	8/8	0.97	0.14	27,33,36,37	0
3	MRY	C	2000	8/8	0.98	0.12	27,29,33,38	0
3	MRY	D	2000	8/8	0.98	0.12	27,27,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.