



# wwPDB NMR Structure Validation Summary Report ⓘ

Feb 17, 2022 – 08:29 AM EST

PDB ID : 1NXR  
Title : HIV-1 POLYPURINE HYBRID, R(GAGGACUG):D(CAGTCCTC), NMR,  
18 STRUCTURES  
Authors : Fedoroff, O.Y.; Ge, Y.; Reid, B.R.  
Deposited on : 1997-02-21

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.26  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.26



## 2 Ensemble composition and analysis

This entry contains 18 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 510 atoms, of which 181 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D(\*CP\*AP\*GP\*TP\*CP\*CP\*TP\*C)-3').

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	8	248	76	92	26	47	7	0

- Molecule 2 is a RNA chain called RNA (5'-R(\*GP\*AP\*GP\*GP\*AP\*CP\*UP\*G)-3').

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
2	B	8	262	78	89	35	53	7	0


## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble


These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA (5'-D(\*CP\*AP\*GP\*TP\*CP\*CP\*TP\*C)-3')

Chain A: 



- Molecule 2: RNA (5'-R(\*GP\*AP\*GP\*GP\*AP\*CP\*UP\*G)-3')

Chain B: 




### 4.2 Residue scores for the first model from the NMR ensemble

No representative models were identified. Colouring as in section 4.1 above.


- Molecule 1: DNA (5'-D(\*CP\*AP\*GP\*TP\*CP\*CP\*TP\*C)-3')

Chain A: 



- Molecule 2: RNA (5'-R(\*GP\*AP\*GP\*GP\*AP\*CP\*UP\*G)-3')

Chain B: 



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *DISTANCE GEOMETRY, : RESTRAINED MOLECULAR : DYNAMICS*.

Of the 20 calculated structures, 18 were deposited, based on the following criterion: *RMSD*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DISCOVER, WITH AMBER FORCE FIELD	refinement	FIELD
BIOSYM	structure solution	

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.45±0.01	0±0/173 ( 0.0± 0.0%)	2.03±0.03	9±1/264 ( 3.3± 0.4%)
2	B	1.18±0.01	0±0/194 ( 0.0± 0.0%)	2.03±0.01	10±1/302 ( 3.2± 0.3%)
All	All	1.32	0/6606 ( 0.0%)	2.03	330/10188 ( 3.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.9±0.4
2	B	0.0±0.0	1.3±0.5
All	All	0	41

There are no bond-length outliers.

5 of 23 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	1	G	O4'-C1'-N9	10.48	116.58	108.20	15	18
2	B	6	C	O4'-C1'-N1	9.21	115.57	108.20	12	18
1	A	1	DC	O4'-C1'-N1	9.21	114.44	108.00	2	18
1	A	5	DC	O4'-C1'-N1	9.17	114.42	108.00	9	18
2	B	5	A	O4'-C1'-N9	8.37	114.90	108.20	7	18

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	7	U	Sidechain	18

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group	Models (Total)
1	A	6	DC	Sidechain	15
2	B	1	G	Sidechain	6
1	A	7	DT	Sidechain	1
1	A	3	DG	Sidechain	1

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	5922	3258	3258	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
2	B	7/8 (88%)	1±1 (13±10%)	0±0 (0±0%)	0.29±0.01
All	All	126/144 (88%)	17 (13%)	0 (0%)	0.29

The overall RNA backbone suiteness is 0.29.

All unique RNA backbone outliers are listed below:



Mol	Chain	Res	Type	Models (Total)
2	B	4	G	10
2	B	5	A	7

There are no RNA pucker outliers to report.

#### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

#### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided