



# Full wwPDB X-ray Structure Validation Report ⓘ

May 11, 2026 – 02:05 PM EDT

PDB ID : 9NZG / pdb\_00009nzg  
Title : De novo designed exatecan-binding protein, EPIC Q51N  
Authors : Polizzi, N.F.  
Deposited on : 2025-03-31  
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	
1	C	148	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4906 atoms, of which 2492 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called De novo designed protein, EPIC Q51N.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	148	Total	C	H	N	O	S	0	1	0
			2398	749	1237	175	235	2			
1	C	148	Total	C	H	N	O	S	0	1	0
			2353	738	1209	175	229	2			

- Molecule 2 is exatecan (CCD ID: A1B7L) (formula:  $C_{24}H_{22}FN_3O_4$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	H	N	O	0	0
			55	24	1	23	3	4		
2	C	1	Total	C	F	H	N	O	0	0
			55	24	1	23	3	4		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total 21	O 21	0	0
3	C	24	Total 24	O 24	0	0



- Molecule 1: De novo designed protein, EPIC Q51N



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.18Å 66.00Å 177.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.45 – 2.21 44.45 – 2.21	Depositor EDS
% Data completeness (in resolution range)	97.2 (44.45-2.21) 97.1 (44.45-2.21)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.32 (at 2.22Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.275 , 0.325 0.275 , 0.323	Depositor DCC
$R_{free}$ test set	802 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.8	Xtriage
Anisotropy	0.591	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 55.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4906	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1B7L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.07	0/1167	0.22	0/1564
1	C	0.07	0/1150	0.20	0/1542
All	All	0.07	0/2317	0.21	0/3106

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1161	1237	1233	7	0
1	C	1144	1209	1205	6	0
2	A	32	23	0	3	0
2	C	32	23	0	2	0
3	A	21	0	0	0	0
3	C	24	0	0	1	0
All	All	2414	2492	2438	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:97:MET:HE1	1:A:135:ALA:HB2	1.80	0.64
1:C:77:THR:HG21	3:C:323:HOH:O	2.00	0.60
1:C:51:ASN:ND2	2:C:201:A1B7L:O2	2.35	0.59
1:C:97:MET:HE1	1:C:135:ALA:HB2	1.85	0.58
1:A:44:LEU:O	1:A:48:LEU:HD12	2.09	0.53
1:A:54:MET:HE1	2:A:201:A1B7L:N3	2.24	0.52
1:A:97:MET:HE3	2:A:201:A1B7L:C12	2.42	0.49
1:C:97:MET:HE3	2:C:201:A1B7L:C12	2.45	0.46
1:A:69:LEU:HD23	1:A:69:LEU:C	2.40	0.45
1:C:69:LEU:HD23	1:C:69:LEU:C	2.41	0.45
1:A:21:LEU:HD13	2:A:201:A1B7L:C24	2.46	0.45
1:A:37:ILE:N	1:A:37:ILE:HD12	2.33	0.44
1:C:10:ILE:HG21	1:C:61:LEU:HB2	2.01	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/148 (99%)	145 (99%)	1 (1%)	1 (1%)	18	18
1	C	147/148 (99%)	146 (99%)	1 (1%)	0	100	100
All	All	294/296 (99%)	291 (99%)	2 (1%)	1 (0%)	36	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	111	VAL



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/127 (95%)	119 (98%)	2 (2%)	53	68
1	C	116/127 (91%)	115 (99%)	1 (1%)	70	82
All	All	237/254 (93%)	234 (99%)	3 (1%)	61	75

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	LEU
1	A	109	LYS
1	C	55	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	A1B7L	A	201	-	31,37,37	4.71	19 (61%)	29,60,60	3.09	12 (41%)
2	A1B7L	C	201	-	31,37,37	4.70	19 (61%)	29,60,60	2.94	11 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1B7L	A	201	-	-	3/3/37/37	0/6/6/6
2	A1B7L	C	201	-	-	0/3/37/37	0/6/6/6

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	A1B7L	C17-C18	10.30	1.53	1.36
2	C	201	A1B7L	C17-C18	10.22	1.53	1.36
2	A	201	A1B7L	O4-C22	8.89	1.40	1.20
2	C	201	A1B7L	O4-C22	8.87	1.40	1.20
2	A	201	A1B7L	O-C16	8.46	1.40	1.23
2	A	201	A1B7L	C10-N3	8.46	1.50	1.37
2	C	201	A1B7L	O-C16	8.43	1.40	1.23
2	C	201	A1B7L	C10-N3	8.38	1.50	1.37
2	C	201	A1B7L	C11-C3	-8.33	1.27	1.41
2	A	201	A1B7L	C11-C3	-8.33	1.27	1.41
2	C	201	A1B7L	C2-N	-7.09	1.40	1.47
2	A	201	A1B7L	C2-N	-7.07	1.40	1.47
2	A	201	A1B7L	C4-C9	-6.32	1.32	1.43
2	C	201	A1B7L	C4-C9	-6.32	1.32	1.43
2	A	201	A1B7L	C12-C10	-5.89	1.32	1.41
2	C	201	A1B7L	C12-C10	-5.88	1.32	1.41
2	A	201	A1B7L	O3-C22	5.04	1.40	1.34
2	C	201	A1B7L	O3-C22	4.99	1.40	1.34
2	A	201	A1B7L	C24-C18	4.96	1.56	1.43
2	C	201	A1B7L	C24-C18	4.92	1.56	1.43
2	C	201	A1B7L	C11-C	-4.66	1.40	1.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	A1B7L	C11-C	-4.53	1.41	1.46
2	C	201	A1B7L	C8-C14	4.53	1.45	1.38
2	A	201	A1B7L	C8-C14	4.52	1.45	1.38
2	C	201	A1B7L	C8-C9	-4.04	1.37	1.43
2	A	201	A1B7L	C8-C9	-4.04	1.37	1.43
2	A	201	A1B7L	C2-C3	3.70	1.54	1.50
2	C	201	A1B7L	C2-C3	3.67	1.54	1.50
2	C	201	A1B7L	C12-C13	3.63	1.41	1.35
2	A	201	A1B7L	C12-C13	3.60	1.41	1.35
2	A	201	A1B7L	C16-C17	2.68	1.50	1.44
2	C	201	A1B7L	C16-C17	2.67	1.50	1.44
2	C	201	A1B7L	C9-C10	2.61	1.47	1.42
2	A	201	A1B7L	C9-C10	2.55	1.46	1.42
2	A	201	A1B7L	O3-C23	-2.48	1.40	1.45
2	C	201	A1B7L	C24-C	2.21	1.41	1.37
2	C	201	A1B7L	O3-C23	-2.17	1.41	1.45
2	A	201	A1B7L	C24-C	2.08	1.41	1.37

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	A1B7L	C2-N-C	-7.12	109.80	113.10
2	C	201	A1B7L	C2-N-C	-6.85	109.92	113.10
2	C	201	A1B7L	C12-C13-C14	-6.64	119.32	124.52
2	A	201	A1B7L	C12-C13-C14	-6.63	119.32	124.52
2	A	201	A1B7L	O3-C22-O4	-5.87	110.47	118.14
2	C	201	A1B7L	C12-C10-C9	5.29	125.94	120.40
2	A	201	A1B7L	C12-C10-C9	5.19	125.83	120.40
2	C	201	A1B7L	C-C11-N3	-4.97	122.20	125.32
2	C	201	A1B7L	C7-C8-C14	-4.60	115.73	123.15
2	A	201	A1B7L	C7-C8-C14	-4.56	115.80	123.15
2	A	201	A1B7L	C-C11-N3	-4.49	122.50	125.32
2	A	201	A1B7L	C9-C10-N3	-3.59	115.46	122.45
2	C	201	A1B7L	C9-C10-N3	-3.57	115.49	122.45
2	A	201	A1B7L	C11-C-N	3.56	109.08	107.06
2	C	201	A1B7L	O3-C22-O4	-3.35	113.77	118.14
2	C	201	A1B7L	C11-C-N	3.25	108.90	107.06
2	A	201	A1B7L	C2-C3-C11	3.17	111.91	109.56
2	C	201	A1B7L	C2-C3-C11	3.16	111.90	109.56
2	A	201	A1B7L	C23-C17-C16	2.67	121.07	116.63
2	A	201	A1B7L	C24-C18-C17	-2.52	117.56	120.25
2	C	201	A1B7L	C23-C17-C16	2.50	120.80	116.63

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	201	A1B7L	C11-N3-C10	-2.34	112.56	117.15
2	A	201	A1B7L	C11-N3-C10	-2.29	112.67	117.15

There are no chirality outliers.

All (3) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	201	A1B7L	O2-C19-C20-C21
2	A	201	A1B7L	C18-C19-C20-C21
2	A	201	A1B7L	C22-C19-C20-C21

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	A1B7L	3	0
2	C	201	A1B7L	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	148/148 (100%)	1.48	42 (28%) <b>1</b> <b>1</b>	27, 53, 90, 115	1 (0%)
1	C	148/148 (100%)	0.98	25 (16%) <b>4</b> <b>3</b>	24, 45, 101, 118	1 (0%)
All	All	296/296 (100%)	1.23	67 (22%) <b>2</b> <b>1</b>	24, 51, 100, 118	2 (0%)

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	108	LEU	5.2
1	C	28	LEU	5.1
1	A	114	LEU	4.9
1	A	117	LEU	4.8
1	A	3	LEU	4.5
1	A	50	LEU	4.4
1	C	114	LEU	4.1
1	A	43	ALA	4.0
1	C	117	LEU	3.8
1	A	2	ASP	3.8
1	A	113	ASP	3.8
1	C	76	ASP	3.6
1	A	107	ILE	3.6
1	A	70	GLU	3.4
1	A	111	VAL	3.3
1	C	47	LEU	3.3
1	C	50	LEU	3.3
1	A	63	ALA	3.3
1	A	77	THR	3.2
1	C	44	LEU	3.2
1	A	45	ALA	3.1
1	A	24	ALA	3.1
1	C	111	VAL	3.1
1	C	75	ILE	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	75	ILE	3.1
1	A	1	GLU	3.0
1	A	38	GLU	3.0
1	C	93	ALA	2.9
1	A	112	LYS	2.9
1	C	43	ALA	2.9
1	A	72	VAL	2.9
1	A	115	GLU	2.9
1	C	113	ASP	2.9
1	C	32	ALA	2.8
1	C	31	ALA	2.8
1	A	80	ALA	2.7
1	C	24	ALA	2.7
1	A	25	LEU	2.6
1	A	54	MET	2.6
1	C	115	GLU	2.6
1	C	42	GLU	2.5
1	A	124	VAL	2.5
1	A	17	LEU	2.5
1	A	69	LEU	2.5
1	A	11	ILE	2.5
1	C	37	ILE	2.5
1	C	108	LEU	2.5
1	A	104	GLY	2.4
1	C	40	ALA	2.4
1	A	87	LEU	2.4
1	C	17	LEU	2.4
1	A	21	LEU	2.4
1	A	74	LYS	2.3
1	C	35	LYS	2.3
1	A	84	VAL	2.3
1	A	145	TYR	2.3
1	A	76	ASP	2.2
1	A	147	ALA	2.2
1	C	36	ASP	2.1
1	C	123	GLU	2.1
1	A	79	ALA	2.1
1	A	116	GLU	2.1
1	A	138	ILE	2.1
1	A	102	LYS	2.0
1	A	40	ALA	2.0
1	C	100	ILE	2.0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	15	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	A1B7L	C	201	32/32	0.83	0.12	32,40,49,52	0
2	A1B7L	A	201	32/32	0.90	0.10	29,40,52,52	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around A1B7L C 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around A1B7L A 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.