



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 03:33 pm BST

PDB ID : 6O2D
Title : Schizosaccharomyces pombe Cnp3 Cupin Domain
Authors : Chik, J.K.; Cho, U.S.
Deposited on : 2019-02-22
Resolution : 2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5743 (2.54-2.50)
Clashscore	141614	6463 (2.54-2.50)
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	
1	B	158	

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 2251 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inner kinetochore subunit cnp3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	137	1086	694	186	201	2	3	0	0	0
1	B	145	1160	739	198	218	2	3	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	486	SER	-	expression tag	UNP Q9USR9
A	487	ASN	-	expression tag	UNP Q9USR9
A	488	ALA	-	expression tag	UNP Q9USR9
B	486	SER	-	expression tag	UNP Q9USR9
B	487	ASN	-	expression tag	UNP Q9USR9
B	488	ALA	-	expression tag	UNP Q9USR9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total 3	O 3	0	0
2	B	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Inner kinetochore subunit cnp3



- Molecule 1: Inner kinetochore subunit cnp3



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	55.16Å 55.16Å 206.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.07 – 2.52 33.07 – 2.52	Depositor EDS
% Data completeness (in resolution range)	99.6 (33.07-2.52) 99.6 (33.07-2.52)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.04 (at 2.51Å)	Xtrriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.217 , 0.251 0.217 , 0.251	Depositor DCC
R_{free} test set	552 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	59.8	Xtrriage
Anisotropy	0.498	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2251	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1105	0.74	4/1482 (0.3%)
1	B	0.44	0/1182	0.71	3/1587 (0.2%)
All	All	0.42	0/2287	0.73	7/3069 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	593	ARG	NE-CZ-NH2	7.88	124.24	120.30
1	A	633	ARG	NE-CZ-NH1	-7.15	116.73	120.30
1	A	518	GLU	CA-CB-CG	6.46	127.62	113.40
1	A	633	ARG	NE-CZ-NH2	6.40	123.50	120.30
1	A	518	GLU	CB-CA-C	-5.59	99.21	110.40
1	B	540	LYS	CD-CE-NZ	5.53	124.42	111.70
1	B	593	ARG	CB-CG-CD	5.07	124.77	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1086	0	1090	15	0
1	B	1160	0	1148	21	0
2	A	3	0	0	0	0
2	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2251	0	2238	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:638:ASP:HB3	1:A:640:PRO:HD2	1.76	0.66
1:A:502:ILE:HD12	1:B:594:MSE:HE3	1.79	0.63
1:A:574:MSE:O	1:A:601:ILE:HG13	2.00	0.61
1:B:581:GLY:HA3	1:B:618:VAL:HG22	1.83	0.61
1:A:621:TYR:HH	1:B:575:SER:HG	1.47	0.61
1:A:574:MSE:HE1	1:A:608:TYR:OH	2.03	0.59
1:B:634:ARG:HH21	1:B:643:ARG:HH22	1.51	0.59
1:B:563:GLU:HB3	1:B:611:LYS:HE3	1.85	0.58
1:B:588:ASN:ND2	1:B:606:ASN:OD1	2.34	0.58
1:B:582:LYS:HZ3	1:B:614:GLY:HA2	1.72	0.55
1:A:573:ILE:CD1	1:B:547:GLU:HG2	2.39	0.52
1:B:637:GLY:HA2	1:B:643:ARG:HG3	1.91	0.52
1:B:585:VAL:HG13	1:B:592:PHE:CZ	2.46	0.50
1:A:563:GLU:HG2	1:A:611:LYS:HG3	1.94	0.50
1:A:586:THR:O	1:A:608:TYR:HA	2.12	0.49
1:A:639:PHE:CD1	1:A:640:PRO:HD3	2.48	0.49
1:A:630:GLU:HA	1:A:633:ARG:HG3	1.93	0.49
1:B:634:ARG:HH21	1:B:643:ARG:NH2	2.11	0.48
1:A:581:GLY:HA3	1:A:618:VAL:HG22	1.96	0.48
1:A:505:GLU:HA	1:A:518:GLU:O	2.14	0.47
1:A:531:LEU:HD11	1:A:619:ARG:HH12	1.79	0.47
1:A:520:ARG:NH1	1:A:547:GLU:OE2	2.48	0.46
1:B:582:LYS:NZ	1:B:614:GLY:HA2	2.30	0.46
1:B:582:LYS:NZ	1:B:614:GLY:CA	2.80	0.45
1:B:585:VAL:HG13	1:B:592:PHE:CE1	2.52	0.45
1:B:579:LEU:O	1:B:596:LYS:HD2	2.18	0.44
1:B:586:THR:O	1:B:608:TYR:HA	2.19	0.43
1:A:585:VAL:O	1:A:591:THR:HA	2.20	0.42
1:B:577:CYS:HA	1:B:599:VAL:HG12	2.02	0.42
1:B:630:GLU:OE2	1:B:634:ARG:HD3	2.21	0.41
1:B:510:ASN:HB3	1:B:513:ASN:O	2.20	0.41
1:B:541:PHE:HA	1:B:555:VAL:O	2.21	0.41
1:B:557:GLU:HA	1:B:618:VAL:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/158 (83%)	127 (97%)	4 (3%)	0	100	100
1	B	141/158 (89%)	138 (98%)	3 (2%)	0	100	100
All	All	272/316 (86%)	265 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/135 (88%)	115 (97%)	4 (3%)	37	61
1	B	126/135 (93%)	120 (95%)	6 (5%)	25	45
All	All	245/270 (91%)	235 (96%)	10 (4%)	30	53

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	498	TYR
1	A	525	SER
1	A	543	SER
1	A	633	ARG
1	B	494	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	538	GLN
1	B	541	PHE
1	B	592	PHE
1	B	593	ARG
1	B	595	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	532	GLN
1	A	533	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	134/158 (84%)	0.30	12 (8%) 9 9	41, 66, 99, 110	0
1	B	142/158 (89%)	0.30	9 (6%) 20 21	48, 71, 101, 108	0
All	All	276/316 (87%)	0.30	21 (7%) 13 14	41, 68, 101, 110	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	592	PHE	3.9
1	A	498	TYR	3.4
1	A	577	CYS	3.2
1	A	623	THR	3.1
1	B	643	ARG	3.1
1	B	593	ARG	3.0
1	A	575	SER	3.0
1	B	532	GLN	2.8
1	A	633	ARG	2.7
1	A	514	PRO	2.7
1	B	538	GLN	2.5
1	A	576	PHE	2.4
1	B	634	ARG	2.4
1	A	601	ILE	2.3
1	A	553	ALA	2.3
1	B	512	GLN	2.2
1	A	532	GLN	2.2
1	B	577	CYS	2.1
1	A	621	TYR	2.1
1	A	622	TYR	2.1
1	B	621	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.