

wwPDB X-ray Structure Validation Summary Report (i)

Sep 18, 2021 – 08:03 am BST

PDB ID : 7O20

Title : X-ray structure of furin in complex with the guanylhydrazone-based inhibitor

3 (mi300)

Authors : Dahms, S.O.; Brandstetter, H.

Deposited on : 2021-03-30

Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.23.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{resolution range}(ext{Å}))$
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain				
1	Λ	490				_	
Ţ	Α	480	94%	•	•		



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 7841 atoms, of which 3651 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Furin.


\mathbf{Mol}	Chain	Residues			Atom	ıs			ZeroOcc	AltConf	Trace
1	A	466	Total 7267	C 2299	H 3549	N 671	O 731	S 17	80	41	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	575	SER	=	expression tag	UNP P09958
A	576	GLY	-	expression tag	UNP P09958
A	577	SER	-	expression tag	UNP P09958
A	578	LEU	ı	expression tag	UNP P09958
A	579	VAL	_	expression tag	UNP P09958
A	580	PRO	ı	expression tag	UNP P09958
A	581	ARG	ı	expression tag	UNP P09958
A	582	GLY	-	expression tag	UNP P09958
A	583	SER	ı	expression tag	UNP P09958
A	584	HIS	_	expression tag	UNP P09958
A	585	HIS	ı	expression tag	UNP P09958
A	586	HIS	_	expression tag	UNP P09958
A	587	HIS	_	expression tag	UNP P09958

• Molecule 2 is $[azanyl-[(2E)-2-(1-phenylethylidene)hydrazinyl]methylidene]azanium (three-letter code: UYT) (formula: <math>C_9H_{13}N_4$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
9	Λ	1	Total	al C H N	C H N	0	0		
	Λ	1	26	9	13	4	0	0	
9	Λ	1	Total	С	Н	N	0	0	
	Α	1	26	9	13	4			
2	Λ	1	Total	С	Н	N	0	1	
	A	A 1	52	18	26	8	0		

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
3	A	3	Total Ca 3 3	0	0

• Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).


Mo	ol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
4		A	2	Total Na 2 2	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0

• Molecule 6 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
6	A	1	Total	С	Н	О	S	0	0	
0	A	1	10	2	6	1	1	0	J	
6	A	1	Total	С	Η	О	S	0	0	
0	Λ	1	10	2	6	1	1	0	0	
6	A	1	Total	С	Η	О	S	0	0	
	Λ	1	10	2	6	1	1			
6	A	1	Total	С	Н	Ο	S	0	0	
	Λ	1	10	2	6	1	1	U		
6	A	1	Total	С	Η	Ο	S	0	0	
	Λ	1	10	2	6	1	1	U	U	
6	Δ	1	Total	С	Н	О	S	0	0	
	A	1	10	2	6	1	1		U	

 \bullet Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\rm C_8H_{15}NO_6).$





Mol	Chain	Residues		At	oms			ZeroOcc	AltConf
7	Λ	1	Total	С	Н	N	О	0	1
'	A	1	28	8	14	1	5	0	1

• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	370	Total O 376 376	0	7




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Furin





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 65 2 2	Depositor	
Cell constants	130.32Å 130.32Å 156.04Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor	
Resolution (Å)	45.72 - 1.80	Depositor	
% Data completeness	98.5 (45.72-1.80)	Depositor	
(in resolution range)	30.0 (40.12 1.00)	Depositor	
R_{merge}	(Not available)	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	PHENIX 1.18.2_3874	Depositor	
R, R_{free}	0.162 , 0.176	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	7841	wwPDB-VP	
Average B, all atoms (Å ²)	36.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, NA, DMS, CA, CL, UYT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
		RMSZ	# Z >5	RMSZ	# Z > 5	
1	L	A	0.48	0/3965	0.65	0/5393

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3718	3549	3388	4	0
2	A	52	52	0	0	0
3	A	3	0	0	0	0
4	A	2	0	0	0	0
5	A	1	0	0	0	0
6	A	24	36	36	0	0
7	A	14	14	13	0	0
8	A	376	0	0	0	0
All	All	4190	3651	3437	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	$egin{array}{ll} ext{Interatomic} \ ext{distance} \ (ext{\AA}) \end{array}$	Clash overlap (Å)	
1:A:172:SER:HB3	1:A:182:PRO:HG3	1.98	0.46	
1:A:125:SER:HA	1:A:135[A]:LYS:HE3	1.99	0.44	
1:A:117[A]:LYS:NZ	1:A:362:GLU:OE1	2.50	0.44	
1:A:341:TYR:CE2	1:A:428[B]:LEU:HD11	2.55	0.42	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	505/480 (105%)	487 (96%)	17 (3%)	1 (0%)	47 33	

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	153	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

\mathbf{Mol}	Chain	Analysed	Analysed Rotameric Outliers		Percentiles	
1	A	411/388 (106%)	405 (98%)	6 (2%)	65 56	

5 of 6 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	298	ARG
1	A	303	CYS
1	A	566	PHE
1	A	112	GLU
1	A	110	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 17 ligands modelled in this entry, 6 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	T	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	DMS	A	612	-	3,3,3	0.56	0	3,3,3	0.55	0
6	DMS	A	610	_	3,3,3	0.66	0	3,3,3	0.45	0
6	DMS	A	611	_	3,3,3	0.70	0	3,3,3	0.76	0
2	UYT	A	603[A]	-	13,13,13	2.44	2 (15%)	13,16,16	1.12	2 (15%)
6	DMS	A	613	_	3,3,3	0.57	0	3,3,3	0.45	0
2	UYT	A	603[B]	_	13,13,13	2.89	2 (15%)	13,16,16	1.16	2 (15%)



Mol	Tuna	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	DMS	A	615	-	3,3,3	0.64	0	3,3,3	1.09	0
2	UYT	A	601	-	13,13,13	2.13	2 (15%)	13,16,16	1.09	1 (7%)
7	NAG	A	616[A]	1	14,14,15	0.37	0	17,19,21	0.42	0
6	DMS	A	614	-	3,3,3	0.60	0	3,3,3	0.18	0
2	UYT	A	602	-	13,13,13	2.26	3 (23%)	13,16,16	0.98	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UYT	A	603[A]	-	-	0/9/9/9	0/1/1/1
2	UYT	A	603[B]	-	-	0/9/9/9	0/1/1/1
2	UYT	A	601	-	-	0/9/9/9	0/1/1/1
7	NAG	A	616[A]	1	-	0/6/23/26	0/1/1/1
2	UYT	A	602	-	-	0/9/9/9	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	${f Observed(\AA)}$	$\operatorname{Ideal}(ext{\AA})$
2	A	603[B]	UYT	CAB-NAC	8.44	1.37	1.29
2	A	602	UYT	CAH-CAB	-6.82	1.38	1.49
2	A	603[A]	UYT	CAB-NAC	6.58	1.35	1.29
2	A	603[B]	UYT	CAH-CAB	-5.83	1.40	1.49
2	A	601	UYT	CAH-CAB	-5.61	1.40	1.49

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
2	A	602	UYT	CAH-CAB-NAC	2.60	118.85	115.23
2	A	603[B]	UYT	CAI-CAH-CAB	-2.46	118.31	121.26
2	A	603[A]	UYT	CAH-CAB-NAC	2.38	118.54	115.23
2	A	601	UYT	CAH-CAB-NAC	2.25	118.35	115.23
2	A	603[B]	UYT	CAH-CAB-NAC	2.20	118.29	115.23

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

