



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 4, 2023 – 05:23 AM EDT

PDB ID : 6O3E  
Title : mouse aE-catenin 82-883  
Authors : Pokutta, S.; Weis, W.I.  
Deposited on : 2019-02-26  
Resolution : 4.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1087 (4.30-3.70)
Clashscore	141614	1148 (4.30-3.70)
Ramachandran outliers	138981	1108 (4.30-3.70)
Sidechain outliers	138945	1099 (4.30-3.70)
RSRZ outliers	127900	1028 (4.34-3.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	806	 53% 13% 33%
1	B	806	 55% 12% 33%

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 8070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catenin alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	536	4004	2485	707	793	19	0	0	0
1	B	541	4066	2515	724	807	20	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	78	GLY	-	expression tag	UNP P26231
A	79	GLY	-	expression tag	UNP P26231
A	80	ILE	-	expression tag	UNP P26231
A	81	LEU	-	expression tag	UNP P26231
A	116	SER	CYS	engineered mutation	UNP P26231
B	78	GLY	-	expression tag	UNP P26231
B	79	GLY	-	expression tag	UNP P26231
B	80	ILE	-	expression tag	UNP P26231
B	81	LEU	-	expression tag	UNP P26231
B	116	SER	CYS	engineered mutation	UNP P26231


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Catenin alpha-1



#### • Molecule 1: Catenin alpha-1



V583	V605	F611	L617	G621	R626	R633	T634	P635	GLU	GLU	LEU	ASP	ASP	SER	SER	ASP	PHE	PHE	GLU	THR	THR	GLU	ASP	PHE	PHE	ASP	ASP	ASP	VAL	ARG	ARG	SER	ARG	GLN	THR	GLU	ASP	GLN	LEU	ILE	ILE	ALA	GLY	GLN	GLN	ALA	ARG	ALA	ILE	MET	LYS	ALA	GLN	PRO	LEU	PRO	GLN	GLU						
GLN	LYS	ALA	LYS	ILE	ALA	GLU	GLN	VAL	SER	PHE	PHE	GLY	GLY	VAL	THR	THR	ILE	ILE	LYS	TRP	TRP	ASP	HIS	ASP	PRO	SER	GLY	ASN	ASP	ALA	ALA	CYS	ILE	ILE	GLN	VAL	LEU	ALA	LYS	GLN	TYR	ALA	TYR	LEU	GLN	LEU	ILE	GLY	THR	ASP	PHE	THR	ARG	ALA	ILE	GLY	LYS	GLY	PRO	LEU	PRO	GLN	ASN	THR
SER	ASP	VAL	ILE	SER	ALA	ALA	LYS	VAL	SER	ALA	GLY	LEU	LYS	LEU	GLY	THR	ILE	ILE	ASN	ASP	ASP	HIS	CYS	PRO	ALA	VAL	SER	VAL	ALA	ALA	CYS	ILE	LYS	GLN	ASP	LEU	TYR	ALA	TYR	LEU	GLN	LEU	ALA	ILE	TYR	CYS	THR	HIS	ASP	PHE	THR	ARG	ALA	ILE	GLY	TYR	LEU	VAL	VAL	LYS	ALA	ALA	GLU	
VAL	GLN	ASN	LEU	GLY	GLY	LEU	VAL	VAL	SER	GLY	VAL	ASP	SER	SER	ILE	GLN	ALA	ASN	LEU	LEU	MET	ASN	ASN	VAL	VAL	VAL	VAL	GLN	THR	THR	VAL	LYS	GLN	ALA	SER	TYR	VAL	ALA	ALA	SER	THR	THR	LYS	GLN	GLY	MET	ALA	SER	SER	LEU	ASN	ASN	PRO	ALA	VAL	VAL	SER	TRP						
LYS	MET	LYS	ALA	PRO	GLU	LYS	LYS	PRO	LEU	VAL	LYS	ARG	GLU	LYS	GLN	ASP	GLU	THR	GLN	THR	THR	ILE	LYS																																									

## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	145.26Å 145.26Å 136.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.97 – 4.00 19.97 – 4.00	Depositor EDS
% Data completeness (in resolution range)	97.8 (19.97-4.00) 97.9 (19.97-4.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.40 (at 4.07Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.213 , 0.243 0.213 , 0.243	Depositor DCC
$R_{free}$ test set	1324 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	200.8	Xtrriage
Anisotropy	0.288	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.22 , 126.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.039 for -h,-k,l 0.053 for h,-h-k,-l 0.036 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8070	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	209.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/4043	0.41	0/5486
1	B	0.25	0/4105	0.40	0/5566
All	All	0.25	0/8148	0.41	0/11052

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4004	0	3922	62	0
1	B	4066	0	3988	50	0
All	All	8070	0	7910	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 109 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:376:ARG:HE	1:B:380:ARG:HH11	1.24	0.84
1:B:195:GLN:HB2	1:B:207:MET:HG2	1.72	0.71

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:563:TYR:OH	1:B:633:ARG:NH2	2.29	0.65
1:B:278:LEU:HD13	1:B:439:SER:HB3	1.78	0.64
1:A:243:LEU:HD21	1:A:462:PRO:HB2	1.81	0.61

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/806 (66%)	508 (96%)	22 (4%)	0	100	100
1	B	535/806 (66%)	511 (96%)	24 (4%)	0	100	100
All	All	1065/1612 (66%)	1019 (96%)	46 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/675 (61%)	396 (96%)	17 (4%)	30	57
1	B	422/675 (62%)	401 (95%)	21 (5%)	24	52
All	All	835/1350 (62%)	797 (95%)	38 (5%)	27	54

5 of 38 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	B	384	LYS
1	B	530	LEU
1	B	386	VAL
1	B	483	ASN
1	B	605	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	169	ASN
1	B	230	GLN
1	B	490	GLN
1	B	481	GLN
1	A	483	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	536/806 (66%)	-0.71	0 100 100	153, 204, 280, 494	0
1	B	541/806 (67%)	-0.68	0 100 100	147, 202, 269, 461	0
All	All	1077/1612 (66%)	-0.69	0 100 100	147, 203, 273, 494	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.