



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 7, 2024 – 04:44 pm GMT

PDB ID : 5O5Z  
Title : CRYSTAL STRUCTURE OF THERMOCOCCUS LITORALIS ADP-DEPENDENT GLUCOKINASE (GK)  
Authors : Herrera-Morande, A.; Castro-Fernandez, V.; Merino, F.; Ramirez-Sarmiento, C.A.; Fernandez, F.J.; Guixe, V.; Vega, M.C.  
Deposited on : 2017-06-02  
Resolution : 2.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	 93% 5%
1	B	464	 92%

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 7428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-dependent glucokinase,ADP-dependent glucokinase,AD P-dependent glucokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	454	Total	C	N	O	S	0	0	0
			3635	2327	618	682	8			
1	B	449	Total	C	N	O	S	0	0	0
			3591	2298	612	674	7			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	49	ALA	LYS	conflict	UNP Q7M537
A	59	LYS	ILE	conflict	UNP Q7M537
A	256	LYS	ARG	conflict	UNP Q7M537
A	363	GLY	-	linker	UNP Q7M537
A	364	GLY	-	linker	UNP Q7M537
A	365	GLY	-	linker	UNP Q7M537
A	399	GLY	-	linker	UNP Q7M537
A	400	GLY	-	linker	UNP Q7M537
A	401	GLY	-	linker	UNP Q7M537
A	402	GLY	-	linker	UNP Q7M537
A	403	GLY	-	linker	UNP Q7M537
B	49	ALA	LYS	conflict	UNP Q7M537
B	59	LYS	ILE	conflict	UNP Q7M537
B	256	LYS	ARG	conflict	UNP Q7M537
B	363	GLY	-	linker	UNP Q7M537
B	364	GLY	-	linker	UNP Q7M537
B	365	GLY	-	linker	UNP Q7M537
B	399	GLY	-	linker	UNP Q7M537
B	400	GLY	-	linker	UNP Q7M537
B	401	GLY	-	linker	UNP Q7M537
B	402	GLY	-	linker	UNP Q7M537
B	403	GLY	-	linker	UNP Q7M537

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>) (labeled as "Ligand

of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 5'-O-[(R)-HYDROXY(THIOPHOSPHONOXY)PHOSPHORYL]ADENOSINE (three-letter code: AT4) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>9</sub>P<sub>2</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	A	1	Total	C	N	O	P	S	0	0
			27	10	5	9	2	1		
4	B	1	Total	C	N	O	P	S	0	0
			27	10	5	9	2	1		

- Molecule 5 is alpha-D-glucopyranose (three-letter code: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	B	1	12	6	6	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	49	49	49	0	0
6	B	22	22	22	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ADP-dependent glucokinase,ADP-dependent glucokinase,ADP-dependent glucokinase



- Molecule 1: ADP-dependent glucokinase,ADP-dependent glucokinase,ADP-dependent glucokinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.28Å 97.83Å 113.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.73 – 2.44 58.73 – 2.44	Depositor EDS
% Data completeness (in resolution range)	99.9 (58.73-2.44) 99.9 (58.73-2.44)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 2.45Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, $R_{free}$	0.195 , 0.222 0.199 , 0.224	Depositor DCC
$R_{free}$ test set	2033 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.6	Xtrriage
Anisotropy	0.670	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 39.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.019 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7428	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, AT4, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/3705	0.45	1/5007 (0.0%)
1	B	0.28	0/3660	0.43	0/4948
All	All	0.29	0/7365	0.44	1/9955 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	363	GLY	N-CA-C	-5.22	100.04	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3635	0	3645	10	0
1	B	3591	0	3598	6	0
2	A	36	0	48	1	0
2	B	24	0	32	0	0
3	A	5	0	0	0	0
4	A	27	0	13	1	0
4	B	27	0	13	2	0
5	B	12	0	12	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	49	0	0	2	0
6	B	22	0	0	0	0
All	All	7428	0	7361	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:GLU:OE1	1:A:168:LYS:HD3	1.84	0.77
1:A:219:MET:O	1:A:224:ARG:NH1	2.19	0.76
1:B:45:LYS:NZ	1:B:111:TRP:O	2.19	0.76
1:A:229:GLU:OE1	1:A:232:ARG:NH1	2.26	0.69
1:A:405:ARG:NH2	6:A:601:HOH:O	2.32	0.61
1:A:342:ASP:OD1	1:A:366:ARG:NH2	2.38	0.55
1:A:191:ARG:NH2	1:B:343:GLU:OE2	2.35	0.54
4:B:506:AT4:S3B	4:B:506:AT4:H5'1	2.49	0.53
2:A:506:GOL:O3	2:A:506:GOL:O1	2.20	0.51
1:A:406:ASP:OD1	6:A:601:HOH:O	2.19	0.49
4:A:508:AT4:S3B	4:A:508:AT4:O5'	2.71	0.49
1:A:205:ARG:HH11	1:A:205:ARG:HG3	1.81	0.46
1:B:74:LYS:NZ	1:B:100:GLU:OE1	2.43	0.46
4:B:506:AT4:S3B	4:B:506:AT4:C5'	3.04	0.46
1:A:296:LEU:N	1:A:297:PRO:CD	2.80	0.44
1:A:205:ARG:HG3	1:A:205:ARG:NH1	2.33	0.43
1:B:318:ASP:OD1	1:B:318:ASP:N	2.53	0.41
1:B:141:VAL:O	1:B:220:ARG:HD2	2.21	0.40
1:B:30:VAL:HG22	1:B:399:GLY:HA3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	452/464 (97%)	439 (97%)	13 (3%)	0	100	100
1	B	447/464 (96%)	437 (98%)	10 (2%)	0	100	100
All	All	899/928 (97%)	876 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/394 (98%)	376 (98%)	9 (2%)	50	63
1	B	380/394 (96%)	370 (97%)	10 (3%)	46	58
All	All	765/788 (97%)	746 (98%)	19 (2%)	47	60

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	LYS
1	A	104	VAL
1	A	118	ILE
1	A	165	ASN
1	A	167	LEU
1	A	280	PHE
1	A	304	LEU
1	A	390	SER
1	A	405	ARG
1	B	70	ASN
1	B	132	VAL
1	B	202	ARG
1	B	204	ASN
1	B	223	PHE
1	B	280	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	294	GLU
1	B	304	LEU
1	B	405	ARG
1	B	451	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	502	-	5,5,5	0.31	0	5,5,5	0.26	0
2	GOL	A	505	-	5,5,5	0.35	0	5,5,5	0.22	0
2	GOL	B	501	-	5,5,5	0.38	0	5,5,5	0.22	0
4	AT4	A	508	-	22,29,29	0.97	1 (4%)	24,45,45	1.43	3 (12%)
5	GLC	B	505	-	12,12,12	0.69	0	17,17,17	0.82	0
2	GOL	A	504	-	5,5,5	0.37	0	5,5,5	0.19	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	B	502	-	5,5,5	0.35	0	5,5,5	0.24	0
4	AT4	B	506	-	22,29,29	1.22	1 (4%)	24,45,45	1.35	3 (12%)
2	GOL	A	506	-	5,5,5	0.39	0	5,5,5	0.13	0
2	GOL	A	503	-	5,5,5	0.36	0	5,5,5	0.14	0
3	SO4	A	507	-	4,4,4	0.13	0	6,6,6	0.17	0
2	GOL	A	501	-	5,5,5	0.33	0	5,5,5	0.16	0
2	GOL	B	504	-	5,5,5	0.37	0	5,5,5	0.28	0
2	GOL	B	503	-	5,5,5	0.35	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	502	-	-	3/4/4/4	-
2	GOL	A	505	-	-	2/4/4/4	-
2	GOL	B	501	-	-	4/4/4/4	-
4	AT4	A	508	-	-	4/10/32/32	0/3/3/3
5	GLC	B	505	-	-	0/2/22/22	0/1/1/1
2	GOL	A	504	-	-	2/4/4/4	-
2	GOL	B	502	-	-	2/4/4/4	-
4	AT4	B	506	-	-	6/10/32/32	0/3/3/3
2	GOL	A	506	-	-	2/4/4/4	-
2	GOL	A	503	-	-	0/4/4/4	-
2	GOL	A	501	-	-	4/4/4/4	-
2	GOL	B	504	-	-	0/4/4/4	-
2	GOL	B	503	-	-	1/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	506	AT4	PB-O2B	-3.54	1.47	1.56
4	A	508	AT4	C5-C4	2.06	1.46	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	508	AT4	N3-C2-N1	-3.90	122.58	128.68
4	B	506	AT4	N3-C2-N1	-3.64	122.99	128.68

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	506	AT4	C3'-C2'-C1'	3.37	106.05	100.98
4	A	508	AT4	C3'-C2'-C1'	3.12	105.68	100.98
4	A	508	AT4	C4-C5-N7	-2.75	106.53	109.40
4	B	506	AT4	C4-C5-N7	-2.65	106.64	109.40

There are no chirality outliers.

All (30) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	GOL	C1-C2-C3-O3
2	A	504	GOL	O1-C1-C2-C3
2	A	505	GOL	O1-C1-C2-C3
2	B	501	GOL	C1-C2-C3-O3
2	B	502	GOL	O1-C1-C2-O2
2	B	502	GOL	O1-C1-C2-C3
4	A	508	AT4	C3'-C4'-C5'-O5'
4	B	506	AT4	C5'-O5'-PA-O2A
4	B	506	AT4	C5'-O5'-PA-O1A
4	B	506	AT4	C5'-O5'-PA-O3A
4	B	506	AT4	C3'-C4'-C5'-O5'
2	A	502	GOL	O1-C1-C2-O2
2	A	505	GOL	O1-C1-C2-O2
4	B	506	AT4	O4'-C4'-C5'-O5'
2	A	502	GOL	O1-C1-C2-C3
2	A	506	GOL	O1-C1-C2-C3
2	B	501	GOL	O1-C1-C2-C3
4	B	506	AT4	PB-O3A-PA-O2A
2	A	501	GOL	O2-C2-C3-O3
2	B	501	GOL	O1-C1-C2-O2
4	A	508	AT4	O4'-C4'-C5'-O5'
2	A	504	GOL	O1-C1-C2-O2
2	A	506	GOL	O1-C1-C2-O2
4	A	508	AT4	C4'-C5'-O5'-PA
2	A	501	GOL	O1-C1-C2-O2
2	B	503	GOL	O1-C1-C2-O2
2	A	501	GOL	O1-C1-C2-C3
4	A	508	AT4	C5'-O5'-PA-O1A
2	B	501	GOL	O2-C2-C3-O3
2	A	502	GOL	C1-C2-C3-O3



There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	508	AT4	1	0
4	B	506	AT4	2	0
2	A	506	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
































## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	454/464 (97%)	0.01	5 (1%) 80 79	39, 56, 79, 100	0
1	B	449/464 (96%)	0.23	17 (3%) 40 37	43, 61, 87, 108	0
All	All	903/928 (97%)	0.12	22 (2%) 59 54	39, 58, 84, 108	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	401	GLY	5.3
1	B	5	LEU	4.9
1	B	3	GLU	4.2
1	B	165	ASN	3.3
1	A	402	GLY	3.2
1	A	363	GLY	2.9
1	B	14	ARG	2.6
1	B	57	THR	2.6
1	B	193	PHE	2.6
1	B	402	GLY	2.5
1	A	319	GLU	2.3
1	B	196	PHE	2.3
1	B	11	LEU	2.3
1	B	198	VAL	2.2
1	B	64	VAL	2.2
1	B	10	ARG	2.1
1	B	449	LEU	2.1
1	A	453	PHE	2.1
1	B	4	SER	2.1
1	B	197	ASP	2.1
1	B	52	LEU	2.0
1	B	448	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	AT4	B	506	27/27	0.73	0.31	72,74,85,86	0
4	AT4	A	508	27/27	0.74	0.28	58,59,71,71	0
2	GOL	B	501	6/6	0.81	0.24	69,71,71,72	0
2	GOL	A	501	6/6	0.82	0.24	71,71,71,72	0
2	GOL	A	505	6/6	0.84	0.38	70,71,71,71	0
2	GOL	A	502	6/6	0.85	0.31	56,58,58,59	0
2	GOL	B	503	6/6	0.85	0.15	71,73,73,73	0
3	SO4	A	507	5/5	0.86	0.30	57,58,58,58	5
2	GOL	B	504	6/6	0.87	0.24	58,60,60,61	0
2	GOL	B	502	6/6	0.87	0.30	64,66,66,66	0
2	GOL	A	506	6/6	0.88	0.18	55,55,55,55	0
2	GOL	A	504	6/6	0.93	0.23	65,68,69,70	0
2	GOL	A	503	6/6	0.93	0.12	60,61,61,61	0
5	GLC	B	505	12/12	0.94	0.12	58,61,62,63	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around AT4 B 506:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around AT4 A 508:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around GOL B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)













**Electron density around SO4 A 507:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)














**Electron density around GOL B 502:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)


**Electron density around GOL A 506:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









## 6.5 Other polymers [i](#)

There are no such residues in this entry.