



Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 08:07 pm BST

PDB ID : 1O6W
Title : Solution Structure of the Prp40 WW Domain Pair of the Yeast Splicing Factor Prp40
Authors : Wiesner, S.; Stier, G.; Sattler, M.; Macias, M.J.
Deposited on : 2002-10-16

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 82%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	75	

2 Ensemble composition and analysis

This entry contains 7 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:73 (72)	0.42	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	3, 4, 6, 7
2	1, 5
Single-model clusters	2

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1254 atoms, of which 625 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PRE-MRNA PROCESSING PROTEIN PRP40.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	75	1254	406	625	103	119	1	0

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: PRE-MRNA PROCESSING PROTEIN PRP40




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: PRE-MRNA PROCESSING PROTEIN PRP40




4.2.2 Score per residue for model 2

- Molecule 1: PRE-MRNA PROCESSING PROTEIN PRP40




4.2.3 Score per residue for model 3

- Molecule 1: PRE-MRNA PROCESSING PROTEIN PRP40




4.2.4 Score per residue for model 4

- Molecule 1: PRE-MRNA PROCESSING PROTEIN PRP40




4.2.5 Score per residue for model 5

- Molecule 1: PRE-MRNA PROCESSING PROTEIN PRP40




4.2.6 Score per residue for model 6

- Molecule 1: PRE-MRNA PROCESSING PROTEIN PRP40



4.2.7 Score per residue for model 7

- Molecule 1: PRE-MRNA PROCESSING PROTEIN PRP40



5 Refinement protocol and experimental data overview

The models were refined using the following method: *AMBIGUOUS DISTANCE RESTRAINTS SIMULATED ANNEALING WITH TORSION ANGLE DYNAMICS AS IMPLEMENTED IN ARIA1.0*.

Of the 20 calculated structures, 7 were deposited, based on the following criterion: *STRUCTURES WITH ACCEPTABLE COVALEN GEOMETRY, STRUCTURES WITH FAVORABLE NON-BOND ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	refinement	
XEASY	structure solution	
NMRView	structure solution	
PIPP/ CAPP	structure solution	CAPP
ARIA	structure solution	
CNS	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	882
Number of shifts mapped to atoms	858
Number of unparsed shifts	0
Number of shifts with mapping errors	24
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	82%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	602	590	590	14±1
All	All	4214	4130	4130	101

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:8:LYS:HG2	1:A:12:GLY:HA2	0.81	1.53	1	7
1:A:3:ILE:HA	1:A:20:LEU:HD21	0.79	1.52	4	1
1:A:38:LEU:O	1:A:38:LEU:HD13	0.75	1.81	6	2
1:A:29:PRO:HD2	1:A:32:LEU:HD23	0.72	1.60	7	1
1:A:32:LEU:HD12	1:A:32:LEU:O	0.70	1.86	7	1
1:A:29:PRO:HG2	1:A:32:LEU:HB2	0.70	1.64	6	1
1:A:3:ILE:HD13	1:A:20:LEU:HD12	0.69	1.64	6	2
1:A:29:PRO:HD2	1:A:32:LEU:CD2	0.69	2.18	7	1
1:A:30:LYS:HA	1:A:33:ILE:HD12	0.65	1.67	1	5
1:A:28:LYS:HD3	1:A:33:ILE:HG12	0.64	1.68	7	2
1:A:28:LYS:O	1:A:33:ILE:HD11	0.63	1.93	1	7
1:A:3:ILE:HG23	1:A:20:LEU:HG	0.61	1.71	5	4
1:A:5:LYS:HG3	1:A:6:GLU:N	0.61	2.09	5	3
1:A:50:THR:OG1	1:A:54:LYS:HB2	0.61	1.96	6	7

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:3:ILE:HG22	1:A:18:ASN:OD1	0.60	1.97	2	1
1:A:20:LEU:HD22	1:A:20:LEU:H	0.59	1.58	4	1
1:A:21:THR:OG1	1:A:23:LYS:HG2	0.59	1.97	2	1
1:A:49:LYS:HE2	1:A:55:VAL:HG22	0.57	1.75	1	4
1:A:3:ILE:HA	1:A:20:LEU:CD2	0.57	2.30	3	4
1:A:21:THR:HG1	1:A:23:LYS:HG2	0.55	1.61	2	1
1:A:28:LYS:HG2	1:A:32:LEU:HB3	0.55	1.79	1	3
1:A:8:LYS:HG2	1:A:12:GLY:CA	0.54	2.32	3	6
1:A:3:ILE:HG23	1:A:20:LEU:HD23	0.54	1.78	4	1
1:A:3:ILE:HG12	1:A:20:LEU:HG	0.54	1.78	3	3
1:A:43:ASN:HB2	1:A:45:TRP:CD1	0.52	2.39	3	2
1:A:20:LEU:HD23	1:A:20:LEU:N	0.51	2.20	7	1
1:A:4:TRP:CE2	1:A:29:PRO:HG3	0.51	2.40	6	1
1:A:3:ILE:HA	1:A:20:LEU:HG	0.50	1.84	6	3
1:A:4:TRP:CE2	1:A:29:PRO:HG2	0.49	2.42	7	2
1:A:4:TRP:CZ3	1:A:18:ASN:HB2	0.46	2.46	5	1
1:A:36:GLU:HA	1:A:39:LEU:HD12	0.46	1.86	7	2
1:A:16:TYR:O	1:A:25:THR:HG22	0.46	2.10	7	3
1:A:59:ASN:HB2	1:A:64:GLU:HB2	0.44	1.87	3	1
1:A:70:PRO:HB2	1:A:72:PHE:CE1	0.44	2.47	6	1
1:A:31:GLU:O	1:A:31:GLU:HG3	0.44	2.12	3	1
1:A:38:LEU:C	1:A:38:LEU:HD13	0.44	2.33	6	1
1:A:20:LEU:CD2	1:A:20:LEU:N	0.43	2.80	1	1
1:A:59:ASN:CB	1:A:64:GLU:HB2	0.43	2.44	3	2
1:A:20:LEU:N	1:A:20:LEU:CD2	0.43	2.82	3	1
1:A:3:ILE:HG23	1:A:20:LEU:CG	0.42	2.44	2	1
1:A:30:LYS:HG2	1:A:31:GLU:N	0.42	2.29	2	1
1:A:54:LYS:HB3	1:A:67:TRP:HZ3	0.42	1.75	1	1
1:A:18:ASN:HB3	1:A:21:THR:OG1	0.41	2.15	2	1
1:A:50:THR:HG23	1:A:54:LYS:O	0.40	2.16	3	1
1:A:17:TYR:CE2	1:A:19:THR:HG22	0.40	2.51	4	1
1:A:3:ILE:HA	1:A:20:LEU:HD23	0.40	1.93	5	1
1:A:39:LEU:O	1:A:43:ASN:ND2	0.40	2.53	6	1
1:A:70:PRO:HG2	1:A:72:PHE:CZ	0.40	2.52	5	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	72/75 (96%)	68±1 (95±1%)	3±1 (5±1%)	0±0 (0±0%)	50 82
All	All	504/525 (96%)	479 (95%)	24 (5%)	1 (0%)	50 82

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	2	SER	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	63/66 (95%)	40±2 (63±3%)	23±2 (37±3%)	1 7
All	All	441/462 (95%)	278 (63%)	163 (37%)	1 7

All 38 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	42	GLU	7
1	A	19	THR	7
1	A	62	THR	7
1	A	68	THR	7
1	A	37	GLU	7
1	A	40	LEU	7
1	A	32	LEU	7
1	A	8	LYS	6
1	A	11	SER	6
1	A	61	THR	5
1	A	52	ASP	5
1	A	41	ARG	5
1	A	17	TYR	5
1	A	38	LEU	5
1	A	35	GLN	5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	63	ARG	5
1	A	65	THR	5
1	A	5	LYS	5
1	A	49	LYS	4
1	A	43	ASN	4
1	A	20	LEU	4
1	A	73	GLU	4
1	A	30	LYS	4
1	A	66	SER	4
1	A	22	LYS	4
1	A	36	GLU	4
1	A	31	GLU	3
1	A	25	THR	3
1	A	23	LYS	3
1	A	6	GLU	3
1	A	64	GLU	3
1	A	2	SER	2
1	A	3	ILE	2
1	A	9	ASP	2
1	A	27	GLU	1
1	A	28	LYS	1
1	A	13	ARG	1
1	A	54	LYS	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation i

The completeness of assignment taking into account all chemical shift lists is 82% for the well-defined parts and 83% for the entire structure.

7.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping i

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	882
Number of shifts mapped to atoms	858
Number of unparsed shifts	0
Number of shifts with mapping errors	24
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	9

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Residue not found in structure. All 24 occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	9	PRO	CD	48.272	0.02	1
A	10	MET	CB	30.626	0.02	1
A	10	MET	N	120.437	0.02	1
A	9	PRO	HA	4.293	0.02	1
A	10	MET	H	8.664	0.02	1
A	9	PRO	CA	61.05	0.02	1
A	9	PRO	CG	25.078	0.02	1
A	10	MET	CA	53.165	0.02	1
A	10	MET	HB3	2.014	0.02	1
A	11	GLY	CA	43.094	0.02	1
A	9	PRO	HG2	1.869	0.02	2
A	11	GLY	HA2	3.866	0.001	1
A	10	MET	HB2	1.922	0.02	1
A	11	GLY	N	109.519	0.02	1

Continued on next page...

Continued from previous page...

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	10	MET	HG2	2.473	0.02	2
A	9	PRO	HD3	3.366	0.02	1
A	10	MET	CG	29.652	0.02	1
A	9	PRO	HD2	3.541	0.003	1
A	11	GLY	H	8.37	0.02	1
A	11	GLY	HA3	3.832	0.007	1
A	9	PRO	HB3	1.822	0.005	1
A	9	PRO	CB	29.772	0.02	1
A	10	MET	HA	4.398	0.02	1
A	9	PRO	HB2	2.179	0.02	1

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	78	2.55 ± 0.07	Should be applied
$^{13}\text{C}_\beta$	74	2.34 ± 0.12	Should be applied
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	74	0.56 ± 0.43	None needed (imprecise)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 82%, i.e. 767 atoms were assigned a chemical shift out of a possible 931. 4 out of 6 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	282/354 (80%)	141/141 (100%)	72/144 (50%)	69/69 (100%)
Sidechain	406/472 (86%)	254/279 (91%)	145/171 (85%)	7/22 (32%)
Aromatic	79/105 (75%)	41/53 (77%)	34/48 (71%)	4/4 (100%)
Overall	767/931 (82%)	436/473 (92%)	251/363 (69%)	80/95 (84%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 83%, i.e. 809 atoms were assigned a chemical shift out of a possible 980. 4 out of 6 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	294/369 (80%)	147/147 (100%)	75/150 (50%)	72/72 (100%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Sidechain	436/506 (86%)	274/300 (91%)	155/182 (85%)	7/24 (29%)
Aromatic	79/105 (75%)	41/53 (77%)	34/48 (71%)	4/4 (100%)
Overall	809/980 (83%)	462/500 (92%)	264/380 (69%)	83/100 (83%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	13	ARG	NE	114.11	92.63 – 76.73	18.5
1	A	63	ARG	NE	113.75	92.63 – 76.73	18.3
1	A	41	ARG	NE	113.29	92.63 – 76.73	18.0
1	A	59	ASN	HB3	-0.19	4.41 – 1.11	-9.0
1	A	18	ASN	HB3	0.57	4.41 – 1.11	-6.6
1	A	70	PRO	HG3	-0.18	3.56 – 0.26	-6.3
1	A	29	PRO	HG2	0.10	3.48 – 0.38	-5.9
1	A	70	PRO	HG2	0.21	3.48 – 0.38	-5.5
1	A	70	PRO	HB3	0.11	3.81 – 0.21	-5.3

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

