



wwPDB EM Validation Summary Report i

Jul 26, 2021 – 12:23 pm BST

PDB ID : 7O73
EMDB ID : EMD-12744
Title : Yeast RNA polymerase II transcription pre-initiation complex with closed distorted promoter DNA
Authors : Schilbach, S.; Aibara, S.; Dienemann, C.; Grabbe, F.; Cramer, P.
Deposited on : 2021-04-12
Resolution : 3.40 Å(reported)
Based on initial model : 5OQJ

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev84
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.22


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY


The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
9	A	1733		74%	8%	18%
10	B	1224		86%	10%	4%
11	C	347		69%	7%	23%
12	D	221		68%	5%	26%
13	E	215		94%	6%	0%
14	F	155		54%	45%	0%
15	G	177		86%	10%	4%
16	H	146		88%	8%	4%
17	I	122		84%	11%	5%
18	J	70		89%	10%	1%
19	K	120		88%	8%	0%
20	L	70		59%	6%	36%
21	M	352		75%	13%	13%
22	N	106		46%	15%	39%
23	O	247		66%	7%	27%
24	Q	738		28%	70%	0%
25	R	400		56%	11%	33%
26	T	106		49%	12%	39%
27	U	286		34%	63%	0%
28	V	129		73%	8%	19%
29	W	492		56%	7%	37%
30	X	328		54%	9%	37%

2 Entry composition i

There are 35 unique types of molecules in this entry. The entry contains 73768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called General transcription and DNA repair factor IIH helicase subunit XPD.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	0	752	6091	3882	1029	1142	38	0	0

- Molecule 2 is a protein called General transcription and DNA repair factor IIH subunit TFB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	1	522	4214	2660	734	798	22	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	-2	GLY	-	expression tag	UNP P32776
1	-1	GLY	-	expression tag	UNP P32776
1	0	SER	-	expression tag	UNP P32776

- Molecule 3 is a protein called General transcription and DNA repair factor IIH subunit TFB2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	2	445	3597	2327	591	663	16	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
2	-3	GLY	-	expression tag	UNP Q02939
2	-2	PRO	-	expression tag	UNP Q02939
2	-1	GLY	-	expression tag	UNP Q02939
2	0	SER	-	expression tag	UNP Q02939

- Molecule 4 is a protein called RNA polymerase II transcription factor B subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	3	131	1089	692	180	209	8	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
3	-2	GLY	-	expression tag	UNP Q03290
3	-1	PRO	-	expression tag	UNP Q03290
3	0	HIS	-	expression tag	UNP Q03290

- Molecule 5 is a protein called General transcription and DNA repair factor IIH subunit TFB4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	4	293	2274	1454	377	429	14	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
4	-2	SER	-	expression tag	UNP Q12004
4	-1	ASN	-	expression tag	UNP Q12004
4	0	ALA	-	expression tag	UNP Q12004

- Molecule 6 is a protein called General transcription and DNA repair factor IIH subunit TFB5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	5	65	514	326	90	95	3	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
5	-3	GLY	-	expression tag	UNP Q3E7C1
5	-2	PRO	-	expression tag	UNP Q3E7C1
5	-1	GLY	-	expression tag	UNP Q3E7C1
5	0	SER	-	expression tag	UNP Q3E7C1

- Molecule 7 is a protein called General transcription and DNA repair factor IIH subunit SSL1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	6	356	2794	1771	482	513	28	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
6	-2	GLY	-	expression tag	UNP Q04673
6	-1	GLY	-	expression tag	UNP Q04673
6	0	SER	-	expression tag	UNP Q04673

- Molecule 8 is a protein called General transcription and DNA repair factor IIH helicase subunit XPB.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	7	615	Total	C	N	O	S	0	0
			4954	3153	860	914	27		

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	A	1428	Total	C	N	O	S	0	0
			11238	7079	1962	2135	62		

- Molecule 10 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	B	1180	Total	C	N	O	S	0	0
			9404	5946	1643	1760	55		

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	C	266	Total	C	N	O	S	0	0
			2092	1315	348	416	13		

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-28	MET	-	initiating methionine	UNP P16370
C	-27	GLY	-	expression tag	UNP P16370
C	-26	SER	-	expression tag	UNP P16370
C	-25	HIS	-	expression tag	UNP P16370
C	-24	HIS	-	expression tag	UNP P16370
C	-23	HIS	-	expression tag	UNP P16370
C	-22	HIS	-	expression tag	UNP P16370
C	-21	HIS	-	expression tag	UNP P16370
C	-20	HIS	-	expression tag	UNP P16370

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-19	SER	-	expression tag	UNP P16370
C	-18	ASN	-	expression tag	UNP P16370
C	-17	SER	-	expression tag	UNP P16370
C	-16	GLY	-	expression tag	UNP P16370
C	-15	LEU	-	expression tag	UNP P16370
C	-14	ASN	-	expression tag	UNP P16370
C	-13	ASP	-	expression tag	UNP P16370
C	-12	ILE	-	expression tag	UNP P16370
C	-11	PHE	-	expression tag	UNP P16370
C	-10	GLU	-	expression tag	UNP P16370
C	-9	ALA	-	expression tag	UNP P16370
C	-8	GLN	-	expression tag	UNP P16370
C	-7	LYS	-	expression tag	UNP P16370
C	-6	ILE	-	expression tag	UNP P16370
C	-5	GLU	-	expression tag	UNP P16370
C	-4	TRP	-	expression tag	UNP P16370
C	-3	HIS	-	expression tag	UNP P16370
C	-2	GLU	-	expression tag	UNP P16370
C	-1	ASP	-	expression tag	UNP P16370
C	0	THR	-	expression tag	UNP P16370
C	1	GLY	-	expression tag	UNP P16370
C	2	SER	-	expression tag	UNP P16370
C	3	SER	-	expression tag	UNP P16370

- Molecule 12 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	D	163	Total	C	N	O	S	0	0
			1307	807	234	264	2		

- Molecule 13 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	E	214	Total	C	N	O	S	0	0
			1752	1111	309	321	11		

- Molecule 14 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	F	86	Total	C	N	O	S	0	0
			697	445	118	131	3		

- Molecule 15 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
15	G	171	1339	861	222	248	8	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	172	HIS	-	expression tag	UNP P34087
G	173	HIS	-	expression tag	UNP P34087
G	174	HIS	-	expression tag	UNP P34087
G	175	HIS	-	expression tag	UNP P34087
G	176	HIS	-	expression tag	UNP P34087
G	177	HIS	-	expression tag	UNP P34087

- Molecule 16 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
16	H	140	1120	704	188	224	4	0	0

- Molecule 17 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
17	I	116	944	581	172	181	10	0	0

- Molecule 18 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
18	J	69	569	362	101	100	6	0	0

- Molecule 19 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
19	K	115	924	593	157	172	2	0	0

- Molecule 20 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	L	45	Total	C 359	N 221	O 71	S 63	0	0

- Molecule 21 is a protein called Transcription initiation factor IIB.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	M	308	Total	C 2360	N 1489	O 406	S 447	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	346	LYS	-	expression tag	UNP P29055
M	347	HIS	-	expression tag	UNP P29055
M	348	HIS	-	expression tag	UNP P29055
M	349	HIS	-	expression tag	UNP P29055
M	350	HIS	-	expression tag	UNP P29055
M	351	HIS	-	expression tag	UNP P29055
M	352	HIS	-	expression tag	UNP P29055

- Molecule 22 is a DNA chain called Non-template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	N	65	Total	C 1332	N 637	O 236	P 394	0	0

- Molecule 23 is a protein called TATA-box-binding protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	O	181	Total	C 1422	N 925	O 243	S 248	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	241	LYS	-	expression tag	UNP P13393
O	242	HIS	-	expression tag	UNP P13393
O	243	HIS	-	expression tag	UNP P13393
O	244	HIS	-	expression tag	UNP P13393
O	245	HIS	-	expression tag	UNP P13393
O	246	HIS	-	expression tag	UNP P13393
O	247	HIS	-	expression tag	UNP P13393

- Molecule 24 is a protein called Transcription initiation factor IIF subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	Q	221	Total	C	N	O	S	0	0

1871 1179 346 339 7

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Q	-2	GLY	-	expression tag	UNP P41895
Q	-1	PRO	-	expression tag	UNP P41895
Q	0	GLY	-	expression tag	UNP P41895

- Molecule 25 is a protein called Transcription initiation factor IIF subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	R	268	Total	C	N	O	S	0	0

2230 1409 392 419 10

- Molecule 26 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	T	65	Total	C	N	O	P	0	0

1330 636 246 384 64

- Molecule 27 is a protein called Transcription initiation factor IIA large subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	U	107	Total	C	N	O	S	0	0

885 559 147 176 3

- Molecule 28 is a protein called Transcription initiation factor IIA subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	V	104	Total	C	N	O	S	0	0

815 511 136 164 4

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
V	123	LYS	-	expression tag	UNP P32774
V	124	HIS	-	expression tag	UNP P32774
V	125	HIS	-	expression tag	UNP P32774

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
V	126	HIS	-	expression tag	UNP P32774
V	127	HIS	-	expression tag	UNP P32774
V	128	HIS	-	expression tag	UNP P32774
V	129	HIS	-	expression tag	UNP P32774

- Molecule 29 is a protein called Transcription initiation factor IIE subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	W	312	Total	C 2532	N 1595	O 442	S 488	7	0


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
W	483	ALA	-	expression tag	UNP P36100
W	484	ALA	-	expression tag	UNP P36100
W	485	ALA	-	expression tag	UNP P36100
W	486	LEU	-	expression tag	UNP P36100
W	487	GLU	-	expression tag	UNP P36100
W	488	HIS	-	expression tag	UNP P36100
W	489	HIS	-	expression tag	UNP P36100
W	490	HIS	-	expression tag	UNP P36100
W	491	HIS	-	expression tag	UNP P36100
W	492	HIS	-	expression tag	UNP P36100

- Molecule 30 is a protein called Transcription initiation factor IIE subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	X	207	Total	C 1661	N 1061	O 281	S 313	6	0

- Molecule 31 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			AltConf
31	0	1	Total	Fe	S	0
			8	4	4	

- Molecule 32 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms			AltConf
32	3	2	Total	Zn		0
			2	2		
32	4	1	Total	Zn		0
			1	1		
32	6	4	Total	Zn		0
			4	4		
32	A	2	Total	Zn		0
			2	2		
32	B	1	Total	Zn		0
			1	1		
32	C	1	Total	Zn		0
			1	1		
32	I	2	Total	Zn		0
			2	2		
32	J	1	Total	Zn		0
			1	1		
32	L	1	Total	Zn		0
			1	1		
32	M	1	Total	Zn		0
			1	1		
32	W	1	Total	Zn		0
			1	1		

- Molecule 33 is BERYLLIUM TRIFLUORIDE ION (three-letter code: BEF) (formula: BeF₃).




Mol	Chain	Residues	Atoms	AltConf
33	7	1	Total Be F 4 1 3	0





- Molecule 34 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
34	7	1	Total Mg 1 1	0
34	A	1	Total Mg 1 1	0

- Molecule 35 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).





Mol	Chain	Residues	Atoms					AltConf
35	7	1	Total 27	C 10	N 5	O 10	P 2	0







- Molecule 9: DNA-directed RNA polymerase II subunit RPB1
- Chain A: ●



- Molecule 10: DNA-directed RNA polymerase II subunit RPB2
- Chain B: ●






- Molecule 11: DNA-directed RNA polymerase II subunit RPB3

Chain C:



- Molecule 12: DNA-directed RNA polymerase II subunit RPB4

Chain D:




- Molecule 13: DNA-directed RNA polymerases I, II, and III subunit RPABC1


Chain E:




- Molecule 14: DNA-directed RNA polymerases I, II, and III subunit RPABC2




- Molecule 15: DNA-directed RNA polymerase II subunit RPB7




- Molecule 16: DNA-directed RNA polymerases I, II, and III subunit RPABC3




- Molecule 17: DNA-directed RNA polymerase II subunit RPB9




- Molecule 18: DNA-directed RNA polymerases I, II, and III subunit RPABC5




- Molecule 19: DNA-directed RNA polymerase II subunit RPB11



- Molecule 20: DNA-directed RNA polymerases I, II, and III subunit RPABC4




- Molecule 25: Transcription initiation factor IIF subunit beta



- Molecule 26: Template DNA





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	10163	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	43.6	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	4000	Depositor
Magnification	130000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.189	Depositor
Minimum map value	-0.079	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.009	Depositor
Map size (Å)	398.99997, 398.99997, 398.99997	wwPDB
Map dimensions	380, 380, 380	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.05, 1.05, 1.05	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, BEF, MG, SF4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	0	0.24	0/6209	0.39	0/8384
2	1	0.23	0/4277	0.37	0/5755
3	2	0.23	0/3664	0.37	0/4952
4	3	0.23	0/1109	0.36	0/1492
5	4	0.24	0/2312	0.39	0/3127
6	5	0.22	0/520	0.37	0/701
7	6	0.23	0/2851	0.39	0/3856
8	7	0.24	0/5059	0.39	0/6841
9	A	0.24	0/11439	0.40	0/15469
10	B	0.25	0/9589	0.41	0/12934
11	C	0.24	0/2130	0.40	0/2887
12	D	0.24	0/1314	0.40	0/1761
13	E	0.24	0/1788	0.38	0/2406
14	F	0.25	0/709	0.39	0/956
15	G	0.25	0/1367	0.41	0/1844
16	H	0.25	0/1139	0.43	0/1544
17	I	0.25	0/962	0.43	0/1295
18	J	0.25	0/578	0.39	0/775
19	K	0.25	0/942	0.40	0/1272
20	L	0.23	0/361	0.42	0/478
21	M	0.24	0/2387	0.39	0/3211
22	N	0.56	0/1492	0.98	0/2301
23	O	0.25	0/1449	0.43	0/1952
24	Q	0.24	0/1907	0.40	0/2556
25	R	0.23	0/2270	0.41	0/3052
26	T	0.56	0/1493	0.93	0/2302
27	U	0.22	0/898	0.39	0/1212
28	V	0.22	0/822	0.40	0/1109
29	W	0.23	0/2575	0.37	0/3473
30	X	0.23	0/1692	0.38	0/2280
All	All	0.26	0/75304	0.44	0/102177

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	6091	0	6155	60	0
2	1	4214	0	4288	32	0
3	2	3597	0	3687	35	0
4	3	1089	0	1069	8	0
5	4	2274	0	2332	25	0
6	5	514	0	541	1	0
7	6	2794	0	2815	22	0
8	7	4954	0	4946	47	0
9	A	11238	0	11301	98	0
10	B	9404	0	9398	87	0
11	C	2092	0	2050	15	0
12	D	1307	0	1324	10	0
13	E	1752	0	1776	7	0
14	F	697	0	720	2	0
15	G	1339	0	1357	13	0
16	H	1120	0	1086	6	0
17	I	944	0	899	10	0
18	J	569	0	585	8	0
19	K	924	0	934	6	0
20	L	359	0	381	4	0
21	M	2360	0	2467	29	0
22	N	1332	0	737	13	0
23	O	1422	0	1500	12	0
24	Q	1871	0	1883	11	0
25	R	2230	0	2254	39	0
26	T	1330	0	734	9	0
27	U	885	0	866	7	0
28	V	815	0	822	7	0
29	W	2532	0	2529	27	0
30	X	1661	0	1694	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
31	0	8	0	0	0	0
32	3	2	0	0	0	0
32	4	1	0	0	0	0
32	6	4	0	0	0	0
32	A	2	0	0	0	0
32	B	1	0	0	0	0
32	C	1	0	0	0	0
32	I	2	0	0	0	0
32	J	1	0	0	0	0
32	L	1	0	0	0	0
32	M	1	0	0	0	0
32	W	1	0	0	0	0
33	7	4	0	0	1	0
34	7	1	0	0	0	0
34	A	1	0	0	0	0
35	7	27	0	12	1	0
All	All	73768	0	73142	583	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 583 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
30:X:235:THR:OG1	30:X:238:ASP:O	1.89	0.91
10:B:276:ILE:HG21	10:B:280:ILE:HD11	1.61	0.82
21:M:91:ASN:ND2	21:M:123:ASP:OD2	2.11	0.82
8:7:662:ILE:O	8:7:689:ARG:NH1	2.13	0.82
10:B:151:LEU:HD23	10:B:153:ALA:H	1.44	0.82

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	0	750/778 (96%)	733 (98%)	17 (2%)	0	100 100
2	1	508/645 (79%)	505 (99%)	3 (1%)	0	100 100
3	2	435/517 (84%)	432 (99%)	3 (1%)	0	100 100
4	3	129/324 (40%)	126 (98%)	3 (2%)	0	100 100
5	4	287/341 (84%)	286 (100%)	1 (0%)	0	100 100
6	5	63/76 (83%)	61 (97%)	2 (3%)	0	100 100
7	6	352/464 (76%)	348 (99%)	4 (1%)	0	100 100
8	7	609/843 (72%)	593 (97%)	16 (3%)	0	100 100
9	A	1420/1733 (82%)	1392 (98%)	27 (2%)	1 (0%)	51 82
10	B	1168/1224 (95%)	1129 (97%)	39 (3%)	0	100 100
11	C	264/347 (76%)	259 (98%)	5 (2%)	0	100 100
12	D	157/221 (71%)	154 (98%)	3 (2%)	0	100 100
13	E	212/215 (99%)	207 (98%)	5 (2%)	0	100 100
14	F	84/155 (54%)	83 (99%)	1 (1%)	0	100 100
15	G	169/177 (96%)	167 (99%)	2 (1%)	0	100 100
16	H	136/146 (93%)	134 (98%)	2 (2%)	0	100 100
17	I	114/122 (93%)	110 (96%)	4 (4%)	0	100 100
18	J	67/70 (96%)	66 (98%)	1 (2%)	0	100 100
19	K	113/120 (94%)	112 (99%)	1 (1%)	0	100 100
20	L	43/70 (61%)	42 (98%)	1 (2%)	0	100 100
21	M	302/352 (86%)	296 (98%)	6 (2%)	0	100 100
23	O	179/247 (72%)	176 (98%)	3 (2%)	0	100 100
24	Q	215/738 (29%)	207 (96%)	8 (4%)	0	100 100
25	R	264/400 (66%)	255 (97%)	9 (3%)	0	100 100
27	U	101/286 (35%)	97 (96%)	4 (4%)	0	100 100
28	V	100/129 (78%)	99 (99%)	1 (1%)	0	100 100
29	W	304/492 (62%)	294 (97%)	10 (3%)	0	100 100
30	X	203/328 (62%)	192 (95%)	11 (5%)	0	100 100
All	All	8748/11560 (76%)	8555 (98%)	192 (2%)	1 (0%)	100 100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
9	A	465	TYR

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	0	684/707 (97%)	684 (100%)	0	100 100
2	1	483/590 (82%)	483 (100%)	0	100 100
3	2	408/470 (87%)	408 (100%)	0	100 100
4	3	125/305 (41%)	125 (100%)	0	100 100
5	4	260/302 (86%)	260 (100%)	0	100 100
6	5	59/68 (87%)	59 (100%)	0	100 100
7	6	323/419 (77%)	323 (100%)	0	100 100
8	7	547/737 (74%)	547 (100%)	0	100 100
9	A	1250/1520 (82%)	1249 (100%)	1 (0%)	93 98
10	B	1024/1061 (96%)	1023 (100%)	1 (0%)	93 98
11	C	234/299 (78%)	233 (100%)	1 (0%)	91 95
12	D	145/200 (72%)	144 (99%)	1 (1%)	84 92
13	E	196/197 (100%)	196 (100%)	0	100 100
14	F	76/137 (56%)	76 (100%)	0	100 100
15	G	152/158 (96%)	152 (100%)	0	100 100
16	H	123/128 (96%)	123 (100%)	0	100 100
17	I	110/116 (95%)	110 (100%)	0	100 100
18	J	64/65 (98%)	64 (100%)	0	100 100
19	K	99/102 (97%)	99 (100%)	0	100 100
20	L	40/57 (70%)	40 (100%)	0	100 100
21	M	265/306 (87%)	265 (100%)	0	100 100
23	O	153/212 (72%)	153 (100%)	0	100 100
24	Q	204/642 (32%)	204 (100%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
25	R	252/363 (69%)	252 (100%)	0	100	100
27	U	99/260 (38%)	99 (100%)	0	100	100
28	V	94/115 (82%)	94 (100%)	0	100	100
29	W	281/436 (64%)	280 (100%)	1 (0%)	91	95
30	X	186/295 (63%)	186 (100%)	0	100	100
All	All	7936/10267 (77%)	7931 (100%)	5 (0%)	93	98

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
9	A	736	ASN
10	B	928	ARG
11	C	127	ARG
12	D	15	LEU
29	W	116	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	0	362	HIS
2	1	352	ASN
9	A	451	HIS
10	B	667	GLN
16	H	11	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 22 ligands modelled in this entry, 19 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
35	ADP	7	903	34,33	24,29,29	0.68	0	29,45,45	0.70	0
31	SF4	0	801	1	0,12,12	0.00	-	-	-	-
33	BEF	7	901	35	0,3,3	0.00	-	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
35	ADP	7	903	34,33	-	2/12/32/32	0/3/3/3
31	SF4	0	801	1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
35	7	903	ADP	PA-O3A-PB-O2B
35	7	903	ADP	PA-O3A-PB-O3B

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
35	7	903	ADP	1	0
33	7	901	BEF	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.


6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-12744. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [\(i\)](#)


6.2.1 Primary map



X Index: 190




Y Index: 190




Z Index: 190


6.2.2 Raw map



X Index: 190



Y Index: 190




Z Index: 190


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 183




Y Index: 234




Z Index: 124


6.3.2 Raw map



X Index: 196



Y Index: 223




Z Index: 131

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.009. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [\(i\)](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.5.1 emd_12744_msk_1.map [\(i\)](#)



X



Y




Z

7 Map analysis (i)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 685 nm³; this corresponds to an approximate mass of 619 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)




*Reported resolution corresponds to spatial frequency of 0.294 \AA^{-1}

8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.294 \AA^{-1}

8.2 Resolution estimates [\(i\)](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	3.42	4.15	3.57
Calculated*	3.87	7.35	3.99

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.87 differs from the reported value 3.4 by more than 10 %


9 Map-model fit [\(i\)](#)

This section contains information regarding the fit between EMDB map EMD-12744 and PDB model 7O73. Per-residue inclusion information can be found in section 3 on page 15.


9.1 Map-model overlay [\(i\)](#)



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.009 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Atom inclusion [\(i\)](#)



At the recommended contour level, 92% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.