



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 3, 2023 – 03:31 AM EDT

PDB ID : 6OC3
Title : Crystal structure of FluA-20 Fab in complex with the head domain of H1
(A/Solomon Islands/3/2006)
Authors : Wilson, I.A.; Lang, S.
Deposited on : 2019-03-21
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10038 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain of FluA-20 Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	224	Total 1674	C 1056	N 275	O 336	S 7	0	0	0
1	C	221	Total 1656	C 1047	N 272	O 330	S 7	0	0	0

- Molecule 2 is a protein called Light chain of FluA-20 Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	213	Total 1646	C 1031	N 281	O 329	S 5	0	0	0
2	D	213	Total 1646	C 1031	N 281	O 329	S 5	0	0	0

- Molecule 3 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	212	Total 1701	C 1084	N 293	O 320	S 4	0	0	0
3	F	212	Total 1701	C 1084	N 293	O 320	S 4	0	0	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	264	SER	-	expression tag	UNP A7Y8I1
E	265	GLY	-	expression tag	UNP A7Y8I1
E	266	LEU	-	expression tag	UNP A7Y8I1
E	267	VAL	-	expression tag	UNP A7Y8I1
E	268	PRO	-	expression tag	UNP A7Y8I1
E	269	ARG	-	expression tag	UNP A7Y8I1
E	270	GLY	-	expression tag	UNP A7Y8I1
E	271	SER	-	expression tag	UNP A7Y8I1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	272	GLY	-	expression tag	UNP A7Y8I1
E	273	HIS	-	expression tag	UNP A7Y8I1
E	274	HIS	-	expression tag	UNP A7Y8I1
E	275	HIS	-	expression tag	UNP A7Y8I1
E	276	HIS	-	expression tag	UNP A7Y8I1
E	277	HIS	-	expression tag	UNP A7Y8I1
E	278	HIS	-	expression tag	UNP A7Y8I1
F	264	SER	-	expression tag	UNP A7Y8I1
F	265	GLY	-	expression tag	UNP A7Y8I1
F	266	LEU	-	expression tag	UNP A7Y8I1
F	267	VAL	-	expression tag	UNP A7Y8I1
F	268	PRO	-	expression tag	UNP A7Y8I1
F	269	ARG	-	expression tag	UNP A7Y8I1
F	270	GLY	-	expression tag	UNP A7Y8I1
F	271	SER	-	expression tag	UNP A7Y8I1
F	272	GLY	-	expression tag	UNP A7Y8I1
F	273	HIS	-	expression tag	UNP A7Y8I1
F	274	HIS	-	expression tag	UNP A7Y8I1
F	275	HIS	-	expression tag	UNP A7Y8I1
F	276	HIS	-	expression tag	UNP A7Y8I1
F	277	HIS	-	expression tag	UNP A7Y8I1
F	278	HIS	-	expression tag	UNP A7Y8I1

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	E	1	14	8	1	5	14	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	100.47Å 109.78Å 146.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.17 – 2.85	Depositor
% Data completeness (in resolution range)	95.5 (48.17-2.85)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.86Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.237 , 0.257	Depositor
Wilson B-factor (Å ²)	65.1	Xtriage
Anisotropy	0.949	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	10038	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	E	301	3	14,14,15	0.29	0	17,19,21	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	E	301	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	301	NAG	O5-C5-C6-O6
4	E	301	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.