



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2023 – 06:24 AM EDT

PDB ID : 6OE3
Title : Crystal Structure of HIV-1 Reverse Transcriptase in Complex with 5-(2-(2-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)ethoxy)phenoxy)-7-fluoro-2-naphthonitrile (JLJ635), a Non-nucleoside Inhibitor
Authors : Bertoletti, N.; Kudalkar, S.N.; Anderson, K.S.; Cisneros Trigo, J.A.; Jorgensen, W.L.
Deposited on : 2019-03-27
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	
2	B	428	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 REVERSE TRANSCRIPTASE, P66 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	546	4279	2770	699	804	6	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP P03366
A	0	VAL	-	expression tag	UNP P03366
A	172	ALA	LYS	engineered mutation	UNP P03366
A	173	ALA	LYS	engineered mutation	UNP P03366
A	280	SER	CYS	engineered mutation	UNP P03366


- Molecule 2 is a protein called HIV-1 REVERSE TRANSCRIPTASE, P51 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	409	3271	2129	533	603	6	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is 5-{2-[2-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)ethoxy]phenoxy}-7-fluorona phthalene-2-carbonitrile (three-letter code: M9A) (formula: C₂₃H₁₆FN₃O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
3	A	1	31	23	1	3	4	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	A	1	1	1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	8	Total	O	0	0
			8	8		
6	B	5	Total	O	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HIV-1 REVERSE TRANSCRIPTASE, P66 SUBUNIT



• Molecule 2: HIV-1 REVERSE TRANSCRIPTASE, P51 SUBUNIT



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	224.31Å 69.57Å 104.83Å 90.00° 105.79° 90.00°	Depositor
Resolution (Å)	43.17 – 2.90 43.17 – 2.90	Depositor EDS
% Data completeness (in resolution range)	97.7 (43.17-2.90) 97.9 (43.17-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 2.90Å)	Xtrriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.228 , 0.273 0.228 , 0.275	Depositor DCC
R_{free} test set	1709 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	82.9	Xtrriage
Anisotropy	0.215	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7600	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, MG, M9A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/4393	0.52	0/6006
2	B	0.35	0/3363	0.53	0/4592
All	All	0.35	0/7756	0.52	0/10598

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	52	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4279	0	4136	66	0
2	B	3271	0	3199	58	0
3	A	31	0	0	0	0
4	A	1	0	0	0	0
5	A	5	0	0	0	0
6	A	8	0	0	0	0
6	B	5	0	0	0	0
All	All	7600	0	7335	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (115) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:85:GLN:HA	2:B:88:TRP:HB2	1.60	0.84
1:A:454:LYS:HB2	1:A:552:VAL:HG13	1.61	0.83
1:A:434:ILE:H	1:A:494:ASN:HD21	1.27	0.82
2:B:180:ILE:HG12	2:B:189:VAL:HG12	1.66	0.76
2:B:236:PRO:HA	2:B:238:LYS:HG3	1.69	0.74
1:A:73:LYS:NZ	1:A:146:TYR:OH	2.21	0.71
2:B:390:LYS:HB3	2:B:417:VAL:HG21	1.76	0.68
1:A:382:ILE:HA	2:B:136:ASN:HD22	1.61	0.65
1:A:343:GLN:HG3	1:A:349:LEU:HD11	1.80	0.63
1:A:362:THR:HG23	1:A:512:LYS:HD2	1.80	0.62
1:A:458:VAL:HB	1:A:548:VAL:HG22	1.81	0.62
2:B:134:SER:OG	2:B:139:THR:O	2.12	0.60
1:A:51:GLY:O	1:A:53:GLU:N	2.33	0.60
2:B:306:ASN:HA	2:B:309:ILE:HG22	1.85	0.58
2:B:369:THR:HG22	2:B:398:TRP:CH2	2.39	0.58
1:A:350:LYS:NZ	1:A:378:GLU:OE2	2.33	0.57
2:B:296:THR:HB	2:B:299:ALA:H	1.69	0.57
2:B:366:LYS:HD2	2:B:405:TYR:CE1	2.39	0.57
2:B:47:ILE:HG22	2:B:146:TYR:HA	1.86	0.56
1:A:405:TYR:CE2	1:A:407:GLN:HB2	2.41	0.56
2:B:254:VAL:HG22	2:B:293:ILE:HD11	1.88	0.56
1:A:317:VAL:HG11	1:A:347:LYS:HD2	1.87	0.56
1:A:478:GLU:HG2	1:A:499:SER:CB	2.36	0.56
2:B:10:VAL:HG13	2:B:88:TRP:CZ2	2.41	0.55
2:B:101:LYS:O	2:B:236:PRO:HB2	2.06	0.55
2:B:401:TRP:HE3	2:B:404:GLU:HG3	1.72	0.55
1:A:382:ILE:HG23	2:B:136:ASN:ND2	2.23	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:157:PRO:HG3	2:B:184:MET:HA	1.91	0.52
1:A:446:ALA:HB2	1:A:477:THR:HG21	1.92	0.52
2:B:125:ARG:HD3	2:B:147:ASN:HA	1.91	0.52
1:A:17:ASP:OD1	1:A:18:GLY:N	2.42	0.52
2:B:306:ASN:O	2:B:310:LEU:HB2	2.09	0.52
1:A:80:LEU:O	1:A:84:THR:HG23	2.09	0.52
1:A:171:PHE:CZ	1:A:205:LEU:HB2	2.45	0.52
2:B:10:VAL:HA	2:B:88:TRP:HZ2	1.75	0.52
2:B:199:ARG:NE	2:B:233:GLU:OE2	2.43	0.52
1:A:207:GLN:O	1:A:211:ARG:HG3	2.11	0.51
1:A:132:ILE:O	1:A:141:GLY:HA3	2.11	0.51
2:B:379:SER:OG	2:B:387:PRO:HD3	2.12	0.50
2:B:110:ASP:O	2:B:216:THR:HG22	2.11	0.50
1:A:110:ASP:O	1:A:217:PRO:HD3	2.10	0.50
1:A:235:HIS:HB2	1:A:238:LYS:O	2.12	0.50
2:B:240:THR:HG23	2:B:350:LYS:HG3	1.93	0.50
1:A:544:GLY:O	1:A:548:VAL:HG23	2.11	0.50
2:B:85:GLN:HG3	2:B:154:LYS:HB3	1.94	0.50
2:B:64:LYS:HD3	2:B:71:TRP:CD2	2.47	0.49
1:A:84:THR:HG21	1:A:153:TRP:HE1	1.76	0.49
2:B:88:TRP:HZ3	2:B:159:ILE:HG13	1.77	0.49
1:A:94:ILE:HG22	1:A:183:TYR:HE1	1.77	0.49
1:A:90:VAL:HG22	1:A:158:ALA:HB2	1.95	0.49
2:B:85:GLN:O	2:B:89:GLU:N	2.45	0.49
2:B:198:HIS:O	2:B:202:ILE:HG12	2.13	0.48
2:B:324:ASP:OD1	2:B:324:ASP:N	2.47	0.48
2:B:245:VAL:O	2:B:263:LYS:NZ	2.43	0.48
2:B:298:GLU:O	2:B:301:LEU:HB2	2.14	0.47
1:A:23:GLN:HE22	1:A:59:PRO:HA	1.77	0.47
1:A:269:GLN:HA	1:A:351:THR:O	2.14	0.47
1:A:410:TRP:HZ3	2:B:405:TYR:HE2	1.61	0.47
2:B:150:PRO:HG2	2:B:153:TRP:HB2	1.96	0.47
2:B:73:LYS:HG3	2:B:74:LEU:N	2.28	0.47
1:A:416:PHE:HZ	1:A:422:LEU:HD21	1.80	0.46
1:A:496:VAL:HG21	2:B:289:LEU:HD21	1.96	0.46
1:A:340:GLN:HG3	1:A:351:THR:HG22	1.97	0.46
1:A:516:GLU:O	1:A:520:GLN:HG3	2.16	0.45
1:A:88:TRP:CH2	2:B:140:PRO:HB2	2.51	0.45
1:A:478:GLU:HG2	1:A:499:SER:HB3	1.99	0.45
2:B:266:TRP:O	2:B:269:GLN:HG2	2.16	0.45
1:A:379:SER:CB	1:A:387:PRO:HD3	2.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:LEU:HD22	1:A:181:TYR:HE2	1.81	0.45
1:A:177:ASP:OD1	1:A:177:ASP:N	2.43	0.45
1:A:121:ASP:O	1:A:125:ARG:HG3	2.17	0.45
1:A:108:VAL:HG13	1:A:188:TYR:CE1	2.52	0.45
1:A:3:SER:N	1:A:5:ILE:HD12	2.32	0.45
2:B:108:VAL:O	2:B:109:LEU:HD23	2.16	0.45
1:A:198:HIS:O	1:A:202:ILE:HG12	2.17	0.44
2:B:422:LEU:HD21	2:B:426:TRP:CZ3	2.52	0.44
1:A:265:ASN:O	1:A:268:SER:OG	2.25	0.44
1:A:406:TRP:CE3	2:B:420:PRO:HD3	2.52	0.44
1:A:350:LYS:HE3	1:A:350:LYS:HB2	1.88	0.44
1:A:382:ILE:HG23	2:B:136:ASN:HD21	1.82	0.44
1:A:54:ASN:OD1	1:A:143:ARG:NH1	2.50	0.44
1:A:5:ILE:HG12	1:A:167:ILE:HD11	2.00	0.43
2:B:336:GLN:HE21	2:B:353:LYS:HD3	1.82	0.43
1:A:333:GLY:O	1:A:336:GLN:NE2	2.35	0.43
1:A:202:ILE:O	1:A:206:ARG:HG3	2.19	0.43
1:A:458:VAL:HG22	1:A:464:GLN:HG2	2.00	0.43
1:A:478:GLU:HG2	1:A:499:SER:HB2	2.00	0.43
1:A:186:ASP:HB3	1:A:188:TYR:CE1	2.53	0.43
1:A:434:ILE:N	1:A:494:ASN:HD21	2.05	0.43
2:B:63:ILE:HD13	2:B:74:LEU:HD22	2.00	0.43
1:A:60:VAL:HG21	1:A:130:PHE:CD2	2.54	0.42
1:A:94:ILE:HG22	1:A:183:TYR:CE1	2.54	0.42
1:A:19:PRO:HG3	1:A:80:LEU:HB2	2.01	0.42
2:B:372:VAL:HG13	2:B:389:PHE:CZ	2.55	0.42
2:B:335:GLY:HA2	2:B:367:GLN:HE22	1.84	0.42
2:B:179:VAL:O	2:B:189:VAL:HA	2.20	0.42
1:A:362:THR:OG1	1:A:363:ASN:N	2.51	0.41
1:A:491:LEU:O	1:A:529:GLU:N	2.52	0.41
2:B:365:VAL:O	2:B:369:THR:HG23	2.19	0.41
2:B:401:TRP:CE3	2:B:404:GLU:HG3	2.53	0.41
1:A:366:LYS:O	1:A:370:GLU:HG3	2.21	0.41
1:A:406:TRP:CE2	2:B:419:THR:HG22	2.56	0.41
1:A:105:SER:HB2	1:A:198:HIS:ND1	2.36	0.41
1:A:367:GLN:HE22	1:A:512:LYS:NZ	2.19	0.41
1:A:406:TRP:CZ2	2:B:419:THR:HG22	2.56	0.41
2:B:37:ILE:O	2:B:41:MET:HG3	2.21	0.41
1:A:120:LEU:HD12	1:A:121:ASP:H	1.86	0.40
2:B:156:SER:N	2:B:157:PRO:HD2	2.36	0.40
2:B:310:LEU:HA	2:B:310:LEU:HD12	1.85	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34:LEU:HD23	1:A:34:LEU:HA	1.83	0.40
2:B:270:ILE:HG12	2:B:346:PHE:HB3	2.02	0.40
2:B:76:ASP:OD2	2:B:78:ARG:NE	2.54	0.40
1:A:33:ALA:O	1:A:37:ILE:HG13	2.21	0.40
2:B:64:LYS:HD3	2:B:71:TRP:CE2	2.56	0.40
2:B:203:GLU:O	2:B:207:GLN:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	542/557 (97%)	529 (98%)	13 (2%)	0	100	100
2	B	403/428 (94%)	384 (95%)	17 (4%)	2 (0%)	29	61
All	All	945/985 (96%)	913 (97%)	30 (3%)	2 (0%)	47	78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	241	VAL
2	B	420	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	445/495 (90%)	432 (97%)	13 (3%)	42	76
2	B	348/390 (89%)	332 (95%)	16 (5%)	27	60
All	All	793/885 (90%)	764 (96%)	29 (4%)	34	68

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	THR
1	A	224	GLU
1	A	248	GLU
1	A	338	THR
1	A	353	LYS
1	A	402	TRP
1	A	452	LEU
1	A	459	THR
1	A	463	ARG
1	A	484	LEU
1	A	507	GLN
1	A	515	SER
1	A	551	LEU
2	B	55	PRO
2	B	73	LYS
2	B	85	GLN
2	B	139	THR
2	B	163	SER
2	B	189	VAL
2	B	200	THR
2	B	214	LEU
2	B	261	VAL
2	B	282	LEU
2	B	283	LEU
2	B	310	LEU
2	B	317	VAL
2	B	362	THR
2	B	414	TRP
2	B	422	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	367	GLN
1	A	373	GLN
1	A	428	GLN
1	A	494	ASN
2	B	85	GLN
2	B	91	GLN
2	B	269	GLN
2	B	336	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	A	603	-	4,4,4	0.14	0	6,6,6	0.08	0
3	M9A	A	601	-	34,34,34	0.20	0	46,47,47	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	M9A	A	601	-	-	0/12/12/12	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	546/557 (98%)	-0.03	6 (1%) 80 80	59, 89, 137, 167	0
2	B	409/428 (95%)	-0.04	3 (0%) 87 87	54, 86, 143, 210	0
All	All	955/985 (96%)	-0.03	9 (0%) 84 84	54, 88, 141, 210	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	61	PHE	3.5
2	B	232	TYR	3.2
2	B	360	ALA	3.0
1	A	3	SER	2.9
1	A	410	TRP	2.4
1	A	72	ARG	2.3
1	A	286	THR	2.1
2	B	301	LEU	2.0
1	A	291	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	A	603	5/5	0.84	0.12	137,138,138,140	0
4	MG	A	602	1/1	0.89	0.27	47,47,47,47	0
3	M9A	A	601	31/31	0.95	0.21	61,65,78,90	0

6.5 Other polymers [i](#)

There are no such residues in this entry.